

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 June 2006 (29.06.2006)

PCT

(10) International Publication Number
WO 2006/068867 A1

(51) International Patent Classification:
C07K 14/705 (2006.01) A61K 47/48 (2006.01)
C07K 16/28 (2006.01)

(21) International Application Number:
PCT/US2005/044926

(22) International Filing Date:
12 December 2005 (12.12.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
11/021,874 22 December 2004 (22.12.2004) US

(71) Applicant (for all designated States except US): GENEN-
TECH, INC. [US/US]; 1 DNA Way, South San Francisco,
CA 94080-4990 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CHAN, Andrew
[US/US]; 1201 Cloud Avenue, Menlo Park, CA 94025
(US). GONG, Qian [CN/US]; 755 Neptune Lane, Foster
City, CA 94404 (US). MARTIN, Flavius [RO/US]; 2735
Driftwood Street, Hayward, CA 94545 (US).

(74) Agent: KOWALCHYK, Katherine, M.; Merchant &
Gould P.C., P.O. Box 2903, Minneapolis, MN 55402-0903
(US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,
LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI,
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,
SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 2006/068867 A1

(54) Title: COMBINATION THERAPY FOR B CELL DISORDERS

(57) Abstract: The invention provides methods of treating B cell based malignancies and B-cell regulated autoimmune disorders using a combination therapy of anti-CD20 antibody with a BAFF antagonist.

5

10

COMBINATION THERAPY FOR B CELL DISORDERS

15

20

FIELD OF THE INVENTION

The invention relates to novel combination therapies for the treatment of B cell malignancies as well as autoimmune disorders.

BACKGROUND OF THE INVENTION

25

Lymphocytes are one of several populations of white blood cells; they specifically recognize and respond to foreign antigen. The three major classes of lymphocytes are B lymphocytes (B cells), T lymphocytes (T cells) and natural killer (NK) cells. B lymphocytes are the cells responsible for antibody production and provide humoral immunity. B cells mature within the bone marrow and leave the marrow expressing an antigen-binding antibody on their cell surface. When a naive B cell first encounters the antigen for which its membrane-bound antibody is specific, the cell begins to divide rapidly and its progeny differentiate into

30

memory B cells and effector cells called "plasma cells". Memory B cells have a longer life span and continue to express membrane-bound antibody with the same specificity as the original parent cell. Plasma cells do not produce membrane-bound antibody but instead produce secreted form of the antibody. Secreted antibodies are the major effector molecules of humoral immunity.

35

The CD20 antigen (also called human B-lymphocyte-restricted differentiation antigen, Bp35) is a hydrophobic transmembrane protein with a molecular weight of approximately 35 kD located on pre-B and mature B lymphocytes (Valentine *et al. J. Biol. Chem.* 264(19):11282-11287 (1989); and Einfeld *et al. EMBO J.* 7(3):711-717 (1988)). The antigen is also expressed on greater than 90% of B cell non-Hodgkin's lymphomas (NHL) (Anderson *et al. Blood* 63(6):1424-1433 (1984)), but is not found on hematopoietic stem cells, pro-B cells, normal plasma cells or other normal tissues (Tedder *et al. J. Immunol.* 135(2):973-979 (1985)). CD20 is thought to regulate an early step(s) in the activation process for cell cycle initiation and differentiation (Tedder *et al., supra*) and possibly functions as a calcium ion channel (Tedder *et al. J. Cell. Biochem.* 14D:195 (1990)).

40

Given the expression of CD20 in B cell lymphomas, this antigen has been a useful therapeutic target to treat such lymphomas. There are more than 300,000 people in the United States with B-cell NHL and more than 56,000 new cases are diagnosed each year. For example, the rituximab (RITUXAN®) antibody which is a genetically engineered chimeric murine/human monoclonal antibody directed against human CD20 antigen (commercially available from Genentech, Inc., South San Francisco, California, U.S.) is used for the treatment

45

5 of patients with relapsed or refractory low-grade or follicular, CD20 positive, B cell non-Hodgkin's lymphoma. Rituximab is the antibody referred to as "C2B8" in US Patent No. 5,736,137 issued April 7, 1998 (Anderson *et al.*). *In vitro* mechanism of action studies have demonstrated that RITUXAN® binds human complement and lyses lymphoid B cell lines through complement-dependent cytotoxicity (CDC) (Reff *et al. Blood* 83(2):435-445 (1994)). Additionally, it has significant activity in assays for antibody-dependent cellular cytotoxicity
 10 (ADCC). *In vivo* preclinical studies have shown that RITUXAN® depletes B cells from the peripheral blood, lymph nodes, and bone marrow of cynomolgus monkeys, presumably through complement and cell-mediated processes (Reff *et al. Blood* 83(2):435-445 (1994)). Other anti-CD20 antibodies indicated for the treatment of NHL include the murine antibody Zevalin™ which is linked to the radioisotope, Yttrium-90 (IDEC Pharmaceuticals, San Diego, CA), Bexxar™ which is a another fully murine antibody conjugated to I-131
 15 (Corixa, WA).

BLyS™ (also known as BAFF, TALL-1, THANK, TNFSF13B, or zTNF4) is a member of the TNF1 ligand superfamily that is essential for B cell survival and maturation. BAFF overexpression in transgenic mice leads to B cell hyperplasia and development of severe autoimmune disease (Mackay, *et al.* (1999) *J. Exp. Med.* 190, 1697-1710; Gross, *et al.* (2000) *Nature* 404, 995-999; Khare, *et al.* (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97, 3370-33752-4). BAFF levels are elevated in human patients with a variety of autoimmune disorders, such as systemic lupus erythematosus, rheumatoid arthritis, and Sjögren's syndrome (Cheema, G. S, *et al.*, (2001) *Arthritis Rheum.* 44, 1313-1319; Groom, J., *et al.* (2002) *J. Clin. Invest.* 109, 59-68; Zhang, J., *et al.*, (2001) *J. Immunol.* 166, 6-10). Furthermore, BAFF levels correlate with disease severity, suggesting that BAFF can play a direct role in the pathogenesis of these illnesses. BAFF acts on B cells by binding to three members of the
 25 TNF receptor superfamily, TACI, BCMA, and BR3 (also known as BAFF-R) (Gross, *et al.*, *supra*; Thompson, J. S., *et al.*, (2001) *Science* 293, 2108-2111; Yan, M., *et al.*, (2001) *Curr. Biol.* 11, 1547-1552; Yan, M., *et al.*, (2000) *Nat. Immunol.* 1, 37-41; Schiemann, B., *et al.*, (2001) *Science* 293, 2111-2114). Of the three, only BR3 is specific for BAFF; the other two also bind the related TNF family member, APRIL. Comparison of the phenotypes of BAFF and receptor knockout or mutant mice indicates that signaling through BR3 mediates the B
 30 cell survival functions of BAFF (Thompson, *et al.*, *supra*; Yan, (2001), *supra*; Schiemann, *supra*). In contrast, TACI appears to act as an inhibitory receptor (Yan, M., (2001) *Nat. Immunol.* 2, 638-643), while the role of BCMA is unclear (Schiemann, *supra*).

BR3 is a 184-residue type III transmembrane protein expressed on the surface of B cells (Thompson, *et al.*, *supra*; Yan, (2002), *supra*). The intracellular region bears no sequence similarity to known structural
 35 domains or protein-protein interaction motifs. Nevertheless, BAFF-induced signaling through BR3 results in processing of the transcription factor NF-B2/p100 to p52 (Claudio, E, *et al.*, (2002) *Nat. Immunol.* 3, 958-965; Kayagaki, N., *et al.*, (2002) *Immunity* 10, 515-524). The extracellular domain (ECD) of BR3 is also divergent. TNFR family members are usually characterized by the presence of multiple cysteine-rich domains (CRDs) in their extracellular region; each CRD is typically composed of ~40 residues stabilized by six cysteines in three
 40 disulfide bonds. Conventional members of this family make contacts with ligand through two CRDs interacting with two distinct patches on the ligand surface (reviewed in Bodmer, J.-L., *et al.*, (2002) *Trends Biochem. Sci.* 27, 19-26). However, the BR3 ECD contains only four cysteine residues, capable of forming a partial CRD at most, raising the question of how such a small receptor imparts high-affinity ligand binding.

5 Previously it has been shown that the BAFF-binding domain of BR3 resides within a 26-residue core region (Kayagaki, et al., *supra*). Six BR3 residues, when structured within a β -hairpin peptide (bhpBR3), were sufficient to confer BAFF binding and block BR3-mediated signaling. Others have reported polypeptides that have been purported to interact with BAFF (e.g., WO 02/24909, WO 03/035846, WO 02/16312, WO02/02641).

10 SUMMARY OF THE INVENTION

The invention provides a method of depleting B cells from a mixed population of cells comprising contacting the mixed population of cells with a BAFF antagonist and a CD20 binding antibody. This method is useful e.g., in a commercial in vitro assay to effectively and selectively deplete B cells from a mixed population of cells, by contacting the B cells with a BAFF antagonist and an anti-CD20 antibody. Another aspect of the
15 preceding method of B cell depletion is to specifically deplete certain subsets of B cells such as germinal center B cells and marginal zone B cells. In a specific embodiment, the germinal center B cells are in the spleen and Peyer's patches. Yet another aspect of the invention is a method of depleting all B cell subsets in vitro or in vivo by contacting the B cells with a BAFF antagonist and a CD20 binding antibody.

The invention also provides a method of depleting all populations of B cells in the spleen by
20 administering to a mammal, a BAFF antagonist and an anti-CD20 antibody in amounts effective to deplete all populations of B cells. In a specific embodiment, the method is effective to deplete marginal zone and germinal center B cells in the spleen, lymph node and Peyer's patches.

Also provided in the invention is a method of treating a B cell neoplasm or malignancy characterized by B cells expressing CD20, comprising administering to a patient suffering from the neoplasm or malignancy, a therapeutically effective amount of a CD20 binding antibody and of a BAFF antagonist. In one embodiment,
25 the CD20 binding antibody and BAFF antagonist are administered concurrently. In a different embodiment, the CD20 binding antibody and BAFF antagonist are administered sequentially. In a specific embodiment, the BAFF antagonist is administered before the CD20 binding antibody. In certain embodiments, the B cell neoplasm is non-Hodgkin's lymphoma (NHL), small lymphocytic (SL) NHL, lymphocyte predominant
30 Hodgkin's disease (LPHD), follicular center cell (FCC) lymphomas, acute lymphocytic leukemia (ALL), chronic lymphocytic leukemia (CLL), diffuse large B cell lymphoma, and Hairy cell leukemia. In this method of treatment, BR3-Fc and Rituxan are administered at dosages disclosed in the section under Dosing. In other embodiments, the BAFF antagonist and the CD20 binding antibody are administered in conjunction with chemotherapy.

Yet another aspect of the invention is a method of alleviating a B-cell regulated autoimmune disorders comprising administering to a patient suffering from the disorder, a therapeutically effective amount of a CD20
35 binding antibody and of a BAFF antagonist. In one embodiment, the autoimmune disorder is selected from the group consisting of rheumatoid arthritis, juvenile rheumatoid arthritis, systemic lupus erythematosus (SLE), lupus nephritis, Wegener's disease, inflammatory bowel disease, idiopathic thrombocytopenic purpura (ITP), thrombotic thrombocytopenic purpura (TTP), autoimmune thrombocytopenia, multiple sclerosis, psoriasis, IgA
40 nephropathy, IgM polyneuropathies, myasthenia gravis, vasculitis, diabetes mellitus, Reynaud's syndrome, Sjorgen's syndrome and glomerulonephritis. Wherein the autoimmune disorder is rheumatoid arthritis or systemic lupus erythematosus, in one embodiment, the BAFF antagonist and the CD20 binding antibody is

5 administered in conjunction with therapy using a drug selected from nonsteroidal anti-inflammatory drugs (NSAIDs), glucocorticoid, prednisone, and disease-modifying antirheumatic drug (DMARD).

In any of the methods of treatment or alleviation of a disorder where the CD20 binding antibody and BAFF antagonist are administered to a patient, the CD20 binding antibody and BAFF antagonist can be administered concurrently or sequentially. In a specific embodiment, the BAFF antagonist is administered
10 before the CD20 binding antibody.

A composition comprising a CD20 binding antibody and a BAFF antagonist is also provided.

Further provided by the invention is an article of manufacture comprising CD20 binding antibody, a BAFF antagonist, and a label wherein the label indicates that the composition is for treating a B cell neoplasm or a B cell regulated autoimmune disorder.

15 In any of the embodiments of the methods, compositions and articles of manufacture of the invention, the anti-CD20 antibody include chimeric and humanized antibody. Specific embodiments of the anti-CD20 antibody include rituximab (RITUXAN®), m2H7 (murine 2H7), hu2H7 (humanized 2H7) and all its functional variants, hu2H7.v16 (v stands for version), v31, v96, v114, v115, having the amino acid sequences. Intact hu2H7.v16 has the mature L chain sequence of SEQ ID NO. 15 and H chain of SEQ ID NO. 16.

20 In any of the embodiments of the methods, compositions and articles of manufacture of the invention, the BAFF antagonist, in one embodiment, is an immunoadhesin. In specific embodiments, the immunoadhesin selected from the group consisting of BR3 immunoadhesin comprising the extracellular domain of BR3 or the sequence of any one of SEQ ID NO.145-165 in FIG. 33 or a "minimal region" thereof; TACI immunoadhesin comprising the extracellular domain of TACI; and BCMA immunoadhesin comprising the extracellular domain
25 of BCMA. In other embodiments, the BAFF antagonist is an anti-BAFF antibody, in particular, an anti-BAFF antibody that binds BAFF within a region of BAFF comprising residues 162-275. In another embodiment, the BAFF antagonist is an anti-BR3 antibody including one that binds BR3 in a region comprising residues 23-38 of human BR3. The amino acid positions of human BR3 referred to in the claims is according to the sequence numbering under human BR3 or alternative human BR3 disclosed herein under the "BR3" definition.

30 In specific embodiments BAFF antagonist is selected from the group consisting of a 17-mer polypeptide having the sequence of ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCSEMLG (SEQ ID NO 7), ECFDLLVRSWVPCHMLR (SEQ ID NO 8), or ECFDLLVRHWVACGLLR (SEQ ID NO 9) as well as PEGylated forms of these 17mers;

a polypeptide having the sequence of

35 MLPGCKWDLLIKQWVCDPLGSGSATGGSGSTASSGSGSATHMLPGCKWDLLIKQWVCDPLG
GGGGVDKTHTCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCWWDVSHEDPEVKFNWYVDGVEV
HNAKTKPREEQYNSTYRWSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPRBPQVYTLPP
SRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN
VFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO. 10);

40 hBR3-Fc immunoadhesin having the sequence of

5 MSALLILALVGAAVASTRRGPRSLRGRDAPAPTPCVPAECFDLLVRHCVACGLLRTPRPKPAGASSPAP
 RTALQPQESQVTDKAAHYTLCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVAVVAVSHEDPEVK
 FNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKG
 QPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKL
 TVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO. 2).

10 In any of the preceding methods of the invention, in one embodiment, the BAFF antagonist and the
 anti-CD20 antibody act synergistically to deplete the B cells.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A-1B show a polynucleotide sequence encoding a native sequence human TACI (SEQ ID
 15 NO: 12) and its amino acid sequence (SEQ ID NO: 25).

Figure 2 shows a polynucleotide sequence encoding a native sequence human BCMA (SEQ ID NO:
 26) and its amino acid sequence (SEQ ID NO: 27).

20 Figure 3 shows a polynucleotide sequence encoding a native sequence human BAFF (SEQ ID NO: 28)
 and its amino acid sequence (SEQ ID NO: 29).

Figures 4A-4B show a polynucleotide sequence encoding a native sequence human APRIL (SEQ ID
 NO: 30) and its putative amino acid sequence (SEQ ID NO: 31).

25 Figure 5A shows a polynucleotide sequence (start and stop codons are underlined) encoding a native
 sequence human TACIs (SEQ ID NO: 52) and Figure 5B shows its amino acid sequence (SEQ ID NO: 53).

Figure 6A shows a polynucleotide sequence (start and stop codons are underlined) encoding a native
 30 sequence human BR3 (SEQ ID NO: 32), and Figure 6B shows its amino acid sequence (SEQ ID NO: 33);
 Figure 6C shows a polynucleotide sequence (start and stop codons are underlined) encoding murine BR3 (SEQ
 ID NO: 34), and Figure 9 shows its amino acid sequence (SEQ ID NO: 35).

Figures 7A-7B show exemplary methods for calculating the % amino acid sequence identity of the
 35 amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO". For
 purposes herein, the "PRO" sequence may be the TACI, BCMA, TALL-1, APRIL, TACIs, or BR3 sequences
 referred to in the Figures herein.

Figure 8 shows an alignment of two amino acid sequences for the TACI receptor, referred to as
 40 "hTACI (265)" (SEQ ID NO: 36), believed to be a spliced variant, and "hTACT", also referred to in Figures 1A-
 1B (SEQ ID NO: 25).

Figure 9 shows a sequence alignment of human (SEQ ID NO: 33) and murine BR3 (SEQ ID NO: 35)
 with identical amino acids indicated by letter and conserved amino acids indicated by a plus sign below.

5

Figure 10 shows the amino acid sequence of human CD20 (SEQ ID NO: 63) showing predicted transmembrane (boxed) and extracellular (underlined) regions. Potential Domains are 1-63: Cytoplasmic; 64-84: Transmembrane; 85-105: Transmembrane; 106-120: Cytoplasmic; 121-141: Transmembrane; 142-188: Extracellular; 189-209: Transmembrane; 210-297: Cytoplasmic; 81-167: Disulfide bond.

10

Figure 11 shows the nucleotide sequence for human CD20 (SEQ ID NO: 64).

FIG. 12 is a sequence alignment comparing the amino acid sequences of the light chain variable domain (V_L) of murine 2H7 (SEQ ID NO. 37), humanized 2H7 v16 variant (SEQ ID NO. 15), and human kappa light chain subgroup I (SEQ ID NO. 38). The CDRs of V_L of 2H7 and hu2H7.v16 are as follows: CDR1 (SEQ ID NO. 39), CDR2 (SEQ ID NO. 40), and CDR3 (SEQ ID NO. 41).

15

FIG. 13 is a sequence alignment which compares the V_H sequences of murine 2H7 (SEQ ID NO. 23), humanized 2H7 v16 variant (SEQ ID NO. 16), and the human consensus sequence of heavy chain subgroup III (SEQ ID NO. 42). The CDRs of V_H of 2H7 and hu2H7.v16 are as follow: CDR1 (SEQ ID NO. 43), CDR2 (SEQ ID NO. 44), and CDR3 (SEQ ID NO. 45).

20

In FIG. 12 and FIG. 13, the CDR1, CDR2 and CDR3 in each chain are enclosed within brackets, flanked by the framework regions, FR1-FR4, as indicated. The asterisks in between two rows of sequences indicate the positions that are different between the two sequences. Residue numbering is according to Kabat et al., Sequences of Immunological Interest. 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991), with insertions shown as a, b, c, d, and e.

25

Figure 14 shows human CD20 transgene expression in mouse B220⁺ cells (B cells) of hCD20 BAC Tg+ mice.

30

Figure 15 shows expression of human CD20 during B cell maturation in hCD20 BAC Tg mice. In Figures 15-19, the red line shows the negative control, staining of cells from transgene negative (Tg-) littermates with an anti-hCD20 mAb. Green line shows staining for human CD20 in Tg+ mice. Tg+ mice refers to hCD20 BAC transgenic mice.

35

Figure 16 shows FACS plots demonstrating expression of human CD20 in the B cells of different maturation/differentiation stages (mature, pre-B and immature B, pro-B and progenitor B) in Tg+ mouse bone marrow.

40

Figure 17 shows FACS plots demonstrating expression of human CD20 in Tg+ mouse splenic B cells. Cells were gated on B220+ to obtain B cells. IgM and CD21 allow delineation into the various B cell subsets of T2/follicular, marginal zone and T1.

5 Figure 18 shows FACS plots demonstrating expression of human CD20 in Tg+ mouse mesenteric lymph nodes (LN).

 Figure 19 shows FACS plots demonstrating expression of human CD20 in Tg+ mouse Peyer's Patches. Cells were gated for B220+. The CD38 marker distinguishes mature from germinal center B cells.

10

 Figure 20 outlines studies on the effects of anti-hCD20 mAb in the human CD20 Tg+ mice. Mice were injected with 1.0mg [equivalent to 50mg/kg] anti-CD20 antibody on day 0 (black arrow above horizontal line) and cells were analyzed on the days indicated by red arrows below the horizontal line. FACS analyses were done on peripheral blood, spleen, lymph node, bone marrow, and Peyer's Patches. Serum levels of anti-hCD20 mAb were monitored.

15

 Figure 21 shows FACS plots demonstrating depletion of peripheral blood B cells with anti-hCD20 mAbs. The left panel shows the IgG control, i.e., animals treated with non-specific, isotype matched antibody.

20

 Figure 22 shows FACS plots demonstrating depletion of mature peripheral LN B cells by anti-hCD20 mAb in the right panel. The left panel shows the IgG control, i.e., animals treated with non-specific, isotype matched antibody. CD21⁺CD23⁺ gates for all B cells.

25

 Figure 23 shows FACS plots demonstrating depletion of splenic T2 B cells, but not marginal zone B cells, by anti-hCD20 mAb.

 Figure 24 shows FACS plots demonstrating depletion of recirculating mature B cells, but not immature/pre-B or pro-B cells, by anti-hCD20 mAb. Red represents IgG-treated while green represents anti-hCD20 mAb treated mice expressing the hCD20. IgG treated mice (red) retained hCD20 expressing mature B cells, while anti-hCD20 mAb depleted hCD20 bearing cells. Human CD20 expression was monitored for detection of both unbound and Ab-bound CD20.

30

 Figure 25 shows FACS plots demonstrating resistance of Peyer's patches germinal center B cells to anti-hCD20 mAbs. The left panel shows cells from the control IgG treated mice. The right panel shows cells from anti-CD20 mAb treated Tg+ mice.

35

 Figure 26 shows FACS plots demonstrating depletion and recovery of B cells in peripheral blood following treatment of the Tg+ mice with anti-hCD20 mAb. The top row panels show staining of cells from control mAb treated mice. Mice were administered antibody at day 1. With time, precursor B cells which do not express hCD20 develop into CD20+ mature B cells (see staining at week 6 and 14).

40

 Figure 27 shows FACS plots demonstrating that resistance of splenic germinal center B cells to short-term (single injection) anti-CD20 mAb treatment. At day 8 following sheep red blood cell immunization to induce germinal center formation, one group of mice was treated with the m2H7 mAb to human CD20. The

5 control set of mice was treated with mIgG2a isotype control antibody. Spleen cells from the mice were analyzed at day 12. PNA (peanut agglutinin) stains for germinal center. No depletion of germinal center B cells was detected with anti-CD20 treatment.

10 Figure 28 shows FACS plots demonstrating that non-depleted marginal zone (MZ) and B1 B cells confer protection to T-independent antigens. On the 3 panels at the right, the spleen cells were stained for the streptococcus polysaccharide-phosphatidyl choline (the TI antigen). CD138 is a marker for plasma cells.

15 Figure 29 shows the distinct biological effects of the combination of a BAFF antagonist, BR3-Fc, and anti-CD20 mAb, m2H7, treatment in an animal model as described in Example 4. FACS analysis was performed on spleen, blood, lymph node B cells (gated on CD21⁺CD23⁺). The combination therapy clearly produced a synergistic effect in depleting B cells, and especially marginal zone, T2 and follicular B cells in the spleen.

20 Figure 30 shows the synergistic effects on B cell depletion of the combination of anti-hCD20 mAb and BR3-Fc in the human CD20 Tg⁺ mice, as described in Example 4. Mice were treated with control IgG_{2a}, BAFFR/BR3-Fc (100 µg/mouse IP daily for 12 days), anti-hCD20 mAb (100 µg/mouse IP on day 9) or the combination of BAFFR/BR3-Fc and anti-hCD20 mAb (same dosing as single treatment groups). B220⁺ splenocytes were isolated on day 13 and stained for CD21 and CD23. N=5 mice/group.

25 Figure 31 shows quantitation of depletion of the B220⁺ total spleen B cells (all subsets of MZ + FO + T1 + T2), marginal zone (MZ) and follicular (FO) B cells from hCD20 Tg⁺ mice as described in Example 4 and Fig. 30 except the mice were treated with single doses of 0.1 mg control IgG_{2a}, BAFF/BR3-Fc or anti-hCD20 mAb. Splenocytes were analyzed on day 4. N=5 mice/group.

30 Figure 32 A-C shows the amino acid sequence of 17mers selected from phage display libraries for high affinity BAFF binding (SEQ ID NOs: 65-142).

Figure 33 shows an alignment of the amino acid sequences of human and mouse BR3 (SEQ ID NO. 146 and 147) extracellular domain (ECD) as well as sequences of variants of BR3 ECD (SEQ ID NO. 148-165).
35 The "minimal region" is boxed.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

40 While anti-CD20 MAb treatment depletes certain subsets of B cells, we have previously observed that the marginal zone B cells, germinal center B cells and plasma cells are preserved. In contrast, blockade of B cell survival signals with BR3-Fc also depletes B cells or modulates B cell numbers, but to a different magnitude. It is believed that BR3 affects the survival of all B cells. It was discovered from the experiments described herein that administration of a combination of anti-CD20 antibody with a BAFF antagonist produced surprisingly synergistic results in depleting B cells in vivo. The combination of anti-CD20 antibody and

therapies directed against the BAFF pathway provides a novel method of treating B cell-mediated diseases including B cell based malignancies and B-cell regulated autoimmune disorders. The combination therapy of anti-CD20 antibody with BAFF antagonist may offer effective and less-toxic alternatives to existing treatments for certain diseases, e.g., chronic lymphocytic leukemia (CLL) and small lymphocytic lymphoma (SLL).

An "autoimmune disease" herein is a non-malignant disease or disorder arising from and directed against an individual's own (self) antigens and/or tissues.

As used herein, "B cell depletion" refers to a reduction in B cell levels in an animal or human after drug or antibody treatment, as compared to the level before treatment. B cell levels are measurable using well known assays such as by getting a complete blood count, by FACS analysis staining for known B cell markers, and by methods such as described in the Experimental Examples. B cell depletion can be partial or complete. In one embodiment, the depletion of CD20 expressing B cells is at least 25%. In a patient receiving a B cell depleting drug, B cells are generally depleted for the duration of time when the drug is circulating in the patient's body and the time for recovery of B cells.

The "CD20" antigen is a non-glycosylated, transmembrane phosphoprotein with a molecular weight of approximately 35 kD that is found on the surface of greater than 90% of B cells from peripheral blood or lymphoid organs. CD20 is expressed during early pre-B cell development and remains until plasma cell differentiation; it is not found on human stem cells, lymphoid progenitor cells or normal plasma cells. CD20 is present on both normal B cells as well as malignant B cells. Other names for CD20 in the literature include "B-lymphocyte-restricted differentiation antigen" and "Bp35". The CD20 antigen is described in, for example, Clark and Ledbetter, *Adv. Can. Res.* 52:81-149 (1989) and Valentine *et al. J. Biol. Chem.* 264(19):11282-11287 (1989).

CD20 binding antibody and anti-CD20 antibody are used interchangeably herein and encompass all antibodies that bind CD20 with sufficient affinity such that the antibody is useful as a therapeutic agent in targeting a cell expressing the antigen, and do not significantly cross-react with other proteins such as a negative control protein in the assays described below. Bispecific antibodies wherein one arm of the antibody binds CD20 are also contemplated. Also encompassed by this definition of CD20 binding antibody are functional fragments of the preceding antibodies. The CD20 binding antibody will bind CD20 with a Kd of < 10nM. In preferred embodiments, the binding is at a Kd of < 7.5nM, more preferably < 5nM, even more preferably at between 1-5nM, most preferably, <1nM.

Examples of antibodies which bind the CD20 antigen include: "C2B8" which is now called "Rituximab" ("RITUXAN®") (US Patent No. 5,736,137, expressly incorporated herein by reference); the yttrium-[90]-labeled 2B8 murine antibody designated "Y2B8" or "Ibritumomab Tiuxetan" ZEVALIN® (US Patent No. 5,736,137, expressly incorporated herein by reference); murine IgG2a "B1," also called "Tositumomab," (Beckman Coulter) optionally labeled with ¹³¹I to generate the "131I-B1" antibody (iodine I131 tositumomab, BEXXAR™) (US Patent No. 5,595,721, expressly incorporated herein by reference); murine monoclonal antibody "1F5" (Press *et al. Blood* 69(2):584-591 (1987) and variants thereof including "framework patched" or humanized 1F5 (WO03/002607, Leung, S.); ATCC deposit HB-96450); murine 2H7 and chimeric 2H7 antibody (US Patent No. 5,677,180, expressly incorporated herein by reference); humanized 2H7; huMax-CD20 (Genmab, Denmark); AME-133 (Applied Molecular Evolution); A20 antibody or variants thereof such as chimeric or humanized A20 antibody (cA20, hA20, respectively) (US 2003/0219433, Immunomedics); and

5 monoclonal antibodies L27, G28-2, 93-1B3, B-C1 or NU-B2 available from the International Leukocyte Typing Workshop (Valentine et al., In: Leukocyte Typing III (McMichael, Ed., p. 440, Oxford University Press (1987)).

The terms "rituximab" or "RITUXAN®" herein refer to the genetically engineered chimeric murine/human monoclonal antibody directed against the CD20 antigen and designated "C2B8" in US Patent No. 5,736,137, expressly incorporated herein by reference, including fragments thereof which retain the ability to
10 bind CD20.

In a specific embodiment, the anti-CD20 antibodies bind human and primate CD20. In specific embodiments, the antibodies that bind CD20 are humanized or chimeric. CD20 binding antibodies include rituximab (RITUXAN®), m2H7 (murine 2H7), hu2H7 (humanized 2H7) and all its functional variants, including without limitation, hu2H7.v16 (v stands for version), v31, v73, v75, as well as fucose deficient
15 variants. The sequences of some of the hu2H7 variant antibodies are provided below, with the sequences N-terminal sequence in bold being the leader sequence which is removed in the mature polypeptide:

hu2H7.v16 L chain [232 aa] (SEQ ID NO. 3)

20 **MGWSCIIILFLVATATGVHSDIQMTQSPSSLSASVGDRTITCRASSSVSYMHYQKPKGK**
APKPLIYAPSNLASGVPSRFSGSGSGTDFLTITSSLPEDFATYYCQQWSFNPPTFGQGT
KVEIKRTVAAPSVFI FPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES
VTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

hu2H7.v16 H chain [471 aa] (SEQ ID NO. 4)

30 **MGWSCIIILFLVATATGVHSEVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAP**
GKGLEWVGAIYPNGDTSYNQKFKGRFTISVDKSKNTLYLQMNSLRAEDTAVYYCARVVY
YSNSYWFYFDVWGQGTLLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT
SWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVE
PKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFN
WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKT
ISKAKGQPREPQVYTLPPSRREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTP
VLDSGDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

hu2H7.v31 H chain [471 aa] (SEQ ID NO. 11)

40 **MGWSCIIILFLVATATGVHSEVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAP**
GKGLEWVGAIYPNGDTSYNQKFKGRFTISVDKSKNTLYLQMNSLRAEDTAVYYCARVVY
YSNSYWFYFDVWGQGTLLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT
SWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVE
PKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFN
WYVDGVEVHNAKTKPREEQYNATYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIAATI
SKAKGQPREPQVYTLPPSRREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTP
VLDSGDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

The L chain of v31 is the same as that of v16 above, i.e., SEQ ID NO. 3.

50 Purely for the purposes herein, "humanized 2H7.v.16" refers to an intact antibody or antibody fragment comprising the variable light chain sequence:

DIQMTQSPSSLSASVGDRTITTCRASSSVSYMHYQKPKGKAPKPLIYAPSNLASGVPSRFSGSGSGTDF
TLTSSLPEDFATYYCQQWSFNPPTFGQGTKEIKR
55 (SEQ ID NO:13); and

variable heavy sequence:

5 EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPGNGDTSYNQKFKG
RFTISVDKSKNTLYLQMNSLRAEDTAVYYCARVVYYSNSYWFYFDVWGQGTLLTVSS
(SEQ ID NO: 14)

Where the humanized 2H7v.16 antibody is an intact antibody, preferably it comprises the v16 light chain amino
10 acid sequence:

DIQMTQSPSSLSASVGDRVTTCRASSSVSYMHVYQQKPGKAPKPLIYAPSNLASGVPSRFSGSGSGTDFT
LTISLQPEDFATYYCQQWSFNPPFTGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPR
EAKVQWKVDNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF
15 NRGE (SEQ ID NO: 15); and

v16 heavy chain amino acid sequence

EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPGNGDTSYNQKFKG
20 RFTISVDKSKNTLYLQMNSLRAEDTAVYYCARVVYYSNSYWFYFDVWGQGTLLTVSSASTKGPSVFPL
APSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT
YICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDV
SHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPI
EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDS
25 GSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK
(SEQ ID NO: 16).

The V region of all other variants based on version 16 will have the amino acid sequences of v16
except at the positions of amino acid substitutions which are indicated in the table below. Unless otherwise
30 indicated, the 2H7 variants will have the same L chain as that of v16.

2H7 version	Heavy chain (V _H) changes	Light chain (V _L) changes	Fc changes
31	-	-	S298A, E333A, K334A (SEQ ID NO. 17)
96	D56A, N100A SEQ ID NO: 46	S92A (SEQ ID NO. 18)	
114	D56A, N100A SEQ ID NO: 46	M32L, S92A (SEQ ID NO. 19)	S298A, E333A, K334A (SEQ ID NO. 20)
115	D56A, N100A SEQ ID NO: 46	M32L, S92A (SEQ ID NO. 21)	S298A, E333A, K334A, E356D, M358L (SEQ ID NO. 22)

A variant of the preceding humanized 2H7 mAb is 2H7v.31 having the same L chain sequence as SEQ
ID NO: 15 above, with the H chain amino acid sequence:

35 EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAP
GKGLEWVGAIYPGNGDTSYNQKFKGRFTISVDKSKNTLYLQMNSLRAEDTAVYYCARVVY
YNSYWFYFDVWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTV
SWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVE
40 PKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFN
WYVDGVEVHNAKTKPREEQYNATYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIAATI
SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTP
VLDSGGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO. 17).

45 The murine anti-human CD20 antibody, m2H7, has the VH sequence:

QAYLQQSGAELVRPGASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSY
NQKFKGKATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGVTGT'TVTV
S (SEQ ID NO: 23)

50

5 And VL sequence :
 QIVLSQSPAILASAPGKVTMTCRASSSVSYMHYQQKPGSSPKPWIYAPSNLASGVPAR
 FSGSGSGTSLTISRVEAEDAATYYCQQWSFNPTFGAGTKLELK (SEQ ID NO: 24)

Unless indicated, the sequences disclosed herein of the humanized 2H7 v.16 and variants thereof are of the
 10 mature polypeptide, i.e., without the leader sequence.

Patents and patent publications concerning CD20 antibodies include US Patent Nos. 5,776,456,
 5,736,137, 5,843,439, 6,399,061, and 6,682,734, as well as US patent appln nos. US 2002/0197255A1, US
 2003/0021781A1, US 2003/0082172 A1, US 2003/0095963 A1, US 2003/0147885 A1 (Anderson *et al.*); US
 Patent No. 6,455,043B1 and WO00/09160 (Grillo-Lopez, A.); WO00/27428 (Grillo-Lopez and White);
 15 WO00/27433 (Grillo-Lopez and Leonard); WO00/44788 (Braslawsky *et al.*); WO01/10462 (Rastetter, W.);
 WO01/10461 (Rastetter and White); WO01/10460 (White and Grillo-Lopez); US2001/0018041A1,
 US2003/0180292A1, WO01/34194 (Hanna and Hariharan); US appln no. US2002/0006404 and WO02/04021
 (Hanna and Hariharan); US appln no. US2002/0012665 A1 and WO01/74388 (Hanna, N.); US appln no. US
 2002/0058029 A1 (Hanna, N.); US appln no. US 2003/0103971 A1 (Hariharan and Hanna); US appln no.
 20 US2002/0009444A1, and WO01/80884 (Grillo-Lopez, A.); WO01/97858 (White, C.); US appln no.
 US2002/0128488A1 and WO02/34790 (Reff, M.); WO02/060955 (Braslawsky *et al.*); WO2/096948 (Braslawsky
et al.); WO02/079255 (Reff and Davies); US Patent No. 6,171,586B1, and WO98/56418 (Lam *et al.*);
 WO98/58964 (Raju, S.); WO99/22764 (Raju, S.); WO99/51642, US Patent No. 6,194,551B1, US Patent No.
 6,242,195B1, US Patent No. 6,528,624B1 and US Patent No. 6,538,124 (Idusogie *et al.*); WO00/42072 (Presta,
 25 L.); WO00/67796 (Curd *et al.*); WO01/03734 (Grillo-Lopez *et al.*); US appln no. US 2002/0004587A1 and
 WO01/77342 (Miller and Presta); US appln no. US2002/0197256 (Grewal, I.); US Appln no. US 2003/0157108
 A1 (Presta, L.); US Patent Nos. 6,565,827B1, 6,090,365B1, 6,287,537B1, 6,015,542, 5,843,398, and 5,595,721,
 (Kaminski *et al.*); US Patent Nos. 5,500,362, 5,677,180, 5,721,108, 6,120,767, 6,652,852B1 (Robinson *et al.*);
 US Pat No. 6,410,391B1 (Raubitschek *et al.*); US Patent No. 6,224,866B1 and WO00/20864 (Barbera-Guillem,
 30 E.); WO01/13945 (Barbera-Guillem, E.); WO00/67795 (Goldenberg); US Appl No. US 2003/0133930 A1 and
 WO00/74718 (Goldenberg and Hansen); WO00/76542 (Golay *et al.*); WO01/72333 (Wolin and Rosenblatt); US
 Patent No. 6,368,596B1 (Ghetie *et al.*); US Patent No. 6,306,393 and US Appln no. US2002/0041847 A1,
 (Goldenberg, D.); US Appln no. US2003/0026801A1 (Weiner and Hartmann); WO02/102312 (Engleman, E.);
 US Patent Application No. 2003/0068664 (Albiter *et al.*); WO03/002607 (Leung, S.); WO 03/049694,
 35 US2002/0009427A1, and US 2003/0185796 A1 (Wolin *et al.*); WO03/061694 (Sing and Siegall); US
 2003/0219818 A1 (Bohen *et al.*); US 2003/0219433 A1 and WO 03/068821 (Hansen *et al.*);
 US2003/0219818A1 (Bohen *et al.*); US2002/0136719A1 (Shenoy *et al.*); WO2004/032828 (Wahl *et al.*), each
 of which is expressly incorporated herein by reference. See, also, US Patent No. 5,849,898 and EP appln no.
 330,191 (Seed *et al.*); US Patent No. 4,861,579 and EP332,865A2 (Meyer and Weiss); USP 4,861,579 (Meyer
 40 *et al.*); WO95/03770 (Bhat *et al.*); US 2003/0219433 A1 (Hansen *et al.*).

The CD20 antibodies can be naked antibody or conjugated to a cytotoxic compound such as a
 radioisotope, or a toxin. Such antibodies include the antibody ZevalinTM which is linked to the radioisotope,
 Yttrium-90 (IDEC Pharmaceuticals, San Diego, CA), and BexxarTM which is conjugated to I-131 (Corixa, WA).
 The humanized 2H7 variants include those that have amino acid substitutions in the FR and affinity maturation
 45 variants with changes in the grafted CDRs. The substituted amino acids in the CDR or FR are not limited to

5 those present in the donor or acceptor antibody. In other embodiments, the anti-CD20 antibodies of the invention further comprise changes in amino acid residues in the Fc region that lead to improved effector function including enhanced CDC and/or ADCC function and B-cell killing (also referred to herein as B-cell depletion). In particular, three mutations have been identified for improving CDC and ADCC activity: S298A/E333A/K334A (also referred to herein as a triple Ala mutant or variant; numbering in the Fc region is
10 according to the EU numbering system; Kabat et al., *supra*) as described (Idusogie et al., *supra* (2001); Shields et al., *supra*).

Other anti-CD20 antibodies of the invention include those having specific changes that improve stability. In one embodiment, the chimeric anti-CD20 antibody has murine V regions and human C region. One such specific chimeric anti-CD20 antibody is Rituxan® (Rituximab®; Genentech, Inc.). Rituximab and hu2H7
15 can mediate lysis of B-cells through both complement-dependent cytotoxicity (CDC) and antibody-dependent cellular cytotoxicity (ADCC). Antibody variants with altered Fc region amino acid sequences and increased or decreased C1q binding capability are described in US patent No. 6,194,551B1 and WO99/51642. The contents of those patent publications are specifically incorporated herein by reference. See, also, Idusogie et al. *J. Immunol.* 164: 4178-4184 (2000).

20 WO00/42072 (Presta) describes polypeptide variants with improved or diminished binding to FcRs. The content of that patent publication is specifically incorporated herein by reference. See, also, Shields et al. *J. Biol. Chem.* 9(2): 6591-6604 (2001).

The N-glycosylation site in IgG is at Asn297 in the CH2 domain. Encompassed herein are humanized CD20-binding antibodies having a Fc region, wherein about 80-100% (and preferably about 90-99%) of the
25 antibody in the composition comprises a mature core carbohydrate structure which lacks fucose, attached to the Fc region of the glycoprotein. Such antibodies show improvement in binding to FcγRIIIA(F158), which is not as effective as FcγRIIIA (V158) in interacting with human IgG.

The term "antibody" is used in the broadest sense and specifically covers, for example, monoclonal antibodies, polyclonal antibodies, antibodies with polypeptidic specificity, single chain antibodies, and
30 fragments of antibodies. According to some embodiments, a polypeptide of this invention is fused into an antibody framework, for example, in the variable region or in a CDR such that the antibody can bind to and inhibit BAFF binding to BR3 or BAFF signaling. The antibodies comprising a polypeptide of this invention can be chimeric, humanized, or human. The antibodies comprising a polypeptide of this invention can be an antibody fragment. Such antibodies and methods of generating them are described in more detail below.
35 Alternatively, an antibody of this invention can be produced by immunizing an animal with a polypeptide of this invention. Thus, an antibody directed against a polypeptide of this invention is contemplated.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that can be present in minor amounts. Monoclonal antibodies
40 are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the

5 character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler et al., *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody
10 libraries using the techniques described in Clackson et al., *Nature*, 352:624-628 (1991) and Marks et al., *J. Mol. Biol.*, 222:581-597 (1991), for example.

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the
15 remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; Morrison et al., *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)). Methods of making chimeric antibodies are known in the art.

"Humanized" forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins,
20 immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementarity-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and
25 capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and maximize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the
30 hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence although the FR regions may include one or more amino acid substitutions that improve binding affinity. The number of these amino acid substitutions in the FR are typically no more than 6 in the H chain, and in the L chain, no more than 3. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a
35 human immunoglobulin. For further details, see Jones et al., *Nature*, 321:522-525 (1986); Reichmann et al., *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992). The humanized antibody includes a PRIMATIZED[®] antibody wherein the antigen-binding region of the antibody is derived from an antibody produced by, e.g., immunizing macaque monkeys with the antigen of interest. Methods of making humanized antibodies are known in the art.

40 Human antibodies can also be produced using various techniques known in the art, including phage-display libraries. Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991). The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies. Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985); Boerner et al., *J. Immunol.*, 147(1):86-95 (1991).

5 “Functional fragments” of the CD20 binding antibodies of the invention are those fragments that retain binding to CD20 with substantially the same affinity as the intact full chain molecule from which they are derived and are able to deplete B cells as measured by *in vitro* or *in vivo* assays such as those described herein. Antibody “effector functions” refer to those biological activities attributable to the Fc region (a native sequence Fc region or amino acid sequence variant Fc region) of an antibody, and vary with the antibody isotype. 10 Examples of antibody effector functions include: C1q binding and complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (*e.g.* B cell receptor); and B cell activation.

 “Antibody-dependent cell-mediated cytotoxicity” or “ADCC” refers to a form of cytotoxicity in which secreted Ig bound onto Fc receptors (FcRs) present on certain cytotoxic cells (*e.g.* Natural Killer (NK) cells, 15 neutrophils, and macrophages) enable these cytotoxic effector cells to bind specifically to an antigen-bearing target cell and subsequently kill the target cell with cytotoxins. The antibodies “arm” the cytotoxic cells and are absolutely required for such killing. The primary cells for mediating ADCC, NK cells, express FcγRIII only, whereas monocytes express FcγRI, FcγRII and FcγRIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991). To assess ADCC activity 20 of a molecule of interest, an *in vitro* ADCC assay, such as that described in US Patent No. 5,500,362 or 5,821,337 may be performed. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, *e.g.*, in a animal model such as that disclosed in Clynes *et al.* *PNAS (USA)* 95:652-656 (1998).

25 “Complement dependent cytotoxicity” or “CDC” refers to the lysis of a target cell in the presence of complement. Activation of the classical complement pathway is initiated by the binding of the first component of the complement system (C1q) to antibodies (of the appropriate subclass) which are bound to their cognate antigen. To assess complement activation, a CDC assay, *e.g.* as described in Gazzano-Santoro *et al.*, *J. Immunol. Methods* 202:163 (1996), may be performed.

30 An “isolated” antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 35 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody’s natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

40 The terms “BAFF,” “BAFF polypeptide,” “TALL-1” or “TALL-1 polypeptide,” “BLys”, “THANK” when used herein encompass “native sequence BAFF polypeptides” and “BAFF variants”. “BAFF” is a designation given to those polypeptides which are encoded by any one of the amino acid sequences shown below:

5

Human BAFF sequence (Fig. 3; SEQ ID NO: 29)

1 mddstereqs rltclckre emklkecvsi lprkespsvr sskdgkllaa tillallsc
 61 ltvvsfyqva alqgdlastr aelqghhaek lpagagapka gleeapavta gkifeppap
 121 gegnssqnsr nkravqgpee tvtdclqli adsetptiqk gsytfvpwll sfkrsgalee
 181 kenkilvket gyffiygqvl ytdktyamgh liqrkkvhvf gdelslvtlf rcinqmpetl
 241 pnnscysagi akleegdelq laiprenaqi sldgdvtffg alkll

10

Mouse BAFF sequence (SEQ ID NO: 47)

1 mdesaktlpp pclfcsekg edmkvgypdi tpqkeegawf gicrdgrlla atllallss
 61 sftamslyql aalqadlmnl rnelqsyrsg atpaaagape ltagvkltp aaprphnssr
 121 ghrnrrafqg peeteqdvdl sappapclpg crhsqhddng mnlrniidc lqliadsdtp
 181 tirkgtytfv pwllsfkrng aleekenkiv vrtqgyffiy sqvlytdpif amghviqrkk
 241 vhfvgdelsl vtlfricqnm pklpnnsy sagiarleeg deiqlaipre naqisrngdd
 301 tffgalkll

15

20

and in Figure 3 and homologs and fragments and variants thereof, which have the biological activity of the native sequence BAFF. A biological activity of BAFF can be selected from the group consisting of promoting B cell survival, promoting B cell maturation and binding to BR3. Variants of BAFF will preferably have at least 80% or any successive integer up to 100% including, more preferably, at least 90%, and even more preferably, at least 95% amino acid sequence identity with a native sequence of a BAFF polypeptide. A "native sequence" BAFF polypeptide comprises a polypeptide having the same amino acid sequence as the corresponding BAFF polypeptide derived from nature. For example, BAFF, exists in a soluble form following cleavage from the cell surface by furin-type proteases. Such native sequence BAFF polypeptides can be isolated from nature or can be produced by recombinant and/or synthetic means. The term "native sequence BAFF polypeptide" specifically encompasses naturally-occurring truncated or secreted forms (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. The term "BAFF" includes those polypeptides described in Shu et al., J. Leukocyte Biol., 65:680 (1999); GenBank Accession No. AF136293; WO98/18921 published May 7, 1998; EP 869,180 published October 7, 1998; WO98/27114 published June 25, 1998; WO99/12964 published March 18, 1999; WO99/33980 published July 8, 1999; Moore et al., Science, 285:260-263 (1999); Schneider et al., J. Exp. Med., 189:1747-1756 (1999); Mukhopadhyay et al., J. Biol. Chem., 274:15978-15981 (1999).

30

35

The term "BAFF antagonist" as used herein is used in the broadest sense, and includes any molecule that (1) binds a native sequence BAFF polypeptide or binds a native sequence BR3 polypeptide to partially or fully block BR3 interaction with BAFF polypeptide, and (2) partially or fully blocks, inhibits, or neutralizes native sequence BAFF signaling. Native sequence BAFF polypeptide signaling promotes, among other things, B cell survival and B cell maturation. The inhibition, blockage or neutralization of BAFF signaling results in, among other things, a reduction in the number of B cells. A BAFF antagonist according to this invention will partially or fully block, inhibit, or neutralize one or more biological activities of a BAFF polypeptide, *in vitro* or *in vivo*. In one embodiment, a biologically active BAFF potentiates any one or combination of the following events *in vitro* or *in vivo*: an increased survival of B cells, an increased level of IgG and/or IgM, an increased numbers of plasma cells, and processing of NF- κ B2/100 to p52 NF- κ B in splenic B cells (e.g., Batten, M et al., (2000) J. Exp. Med. 192:1453-1465; Moore, et al., (1999) Science 285:260-263; Kayagaki, et al., (2002) 10:515-524). Several assays useful for testing BAFF antagonists according to this invention are described herein.

40

45

5 As mentioned above, a BAFF antagonist can function in a direct or indirect manner to partially or fully block, inhibit or neutralize BAFF signaling, *in vitro* or *in vivo*. For instance, the BAFF antagonist can directly bind BAFF. For example, anti-BAFF antibodies that bind within a region of human BAFF comprising residues 162-275 and/or a neighboring residue of a residue selected from the group consisting of 162, 163, 206, 211, 231, 233, 264 and 265 of human BAFF such that the antibody sterically hinders BAFF binding to BR3 is

10 contemplated. In another example, a direct binder is a polypeptide comprising the extracellular domain of a BAFF receptor such as TACI, BR3 and BCMA, or comprising the boxed minimal region of the ECDs as shown in FIG. 33. In another example, BAFF antagonists include the polypeptides having a sequence of that of Formula I, Formula II, Formula III, ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCHMLR (SEQ ID NO 8),

15 ECFDLLVRHWVACGLLR (SEQ ID NO 9), or sequences listed in FIG. 32 and FIG. 33, as described herein. Alternatively, the BAFF antagonist can bind an extracellular domain of a native sequence BR3 at its BAFF binding region to partially or fully block, inhibit or neutralize BAFF binding to BR3 *in vitro*, *in situ*, or *in vivo*. For example, such indirect antagonist is an anti-BR3 antibody that binds in a region of BR3 comprising residues 23-38 of human BR3 or a neighboring region of those residues such that binding of human BR3 to BAFF is

20 sterically hindered.

In some embodiments, a BAFF antagonist according to this invention includes anti-BAFF antibodies, BAFF-binding polypeptides (including immunoadhesins and peptides), and BAFF-binding small molecules. BAFF antagonists include the BAFF binding antibodies described in WO 02/02641 (e.g., antibodies comprising the amino acid sequence of any of SEQ ID NOs. 1-46, 321-329, 834-872, 1563-1595, 1881-1905 of Table 1).

25 In a further embodiment, the immunoadhesin comprises a BAFF binding region of a BAFF receptor (e.g., an extracellular domain of BR3, BCMA or TACI, or a minimal region thereof of the sequences shown in FIG. 33 SEQ ID 146-165). In a still further embodiment, the immunoadhesin is BR3-Fc, or polypeptides having a sequence of that of Formula I, Formula II, Formula III, ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7),

30 ECFDLLVRSWVPCHMLR (SEQ ID NO 8), ECFDLLVRHWVACGLLR (SEQ ID NO 9), or comprising sequences listed in FIG.32 or FIG. 33 or minimal region thereof, optionally, fused or conjugated to an Fc portion of an immunoglobulin. Other examples of BAFF binding Fc proteins can be found in WO 02/66516, WO 00/40716, WO 01/87979, WO 03/024991, WO 02/16412, WO 02/38766, WO 02/092620, WO 01/12812.

According to one embodiment, the BAFF antagonist binds to a BAFF polypeptide or a BR3

35 polypeptide with a binding affinity of 100nM or less. According to another embodiment, the BAFF antagonist binds to a BAFF polypeptide or a BR3 polypeptide with a binding affinity of 10nM or less. According to yet another embodiment, the BAFF antagonist binds to a BAFF polypeptide or a BR3 polypeptide with a binding affinity of 1nM or less

The terms "BR3", "BR3 polypeptide" or "BR3 receptor" when used herein encompass "native

40 sequence BR3 polypeptides" and "BR3 variants" (which are further defined herein). "BR3" is a designation given to those polypeptides comprising any one of the following polynucleotide sequences and homologs thereof:

(a) human BR3 sequence (SEQ ID NO: 33)

5 1 MRRGPRSLRG RDAPAPTPCV PAECFDLLVR HCVACGLLRT PRPKPAGASS PAPRTALQPQ
 61 ESVGAGAGEA ALPLPGLLF APALLGLALV LALVLVGLVS WRRRQRRRLG ASSAEAPDGD
 121 KDAPEPLDKV IILSPGISDA TAPAWPPPGE DPGTTPPGHS VFPVPATELGS TELVTTKTAG
 181 PEQQ

10 (b) alternative human BR3 sequence (SEQ ID NO: 48)

 1 MRRGPRSLRG RDAPAPTPCV PAECFDLLVR HCVACGLLRT PRPKPAGAAS SPAPRTALQP
 61 QESVGAGAGE AALPLPGLLF GAPALLGLAL VLALVLVGLV SWRRRQRRRLR GASSAEAPDGD
 121 DKDAPEPLDK VIILSPGISD ATAPAWPPPGE EDPGTTPPGH SVPVPATELG STELVTTKTA
 181 GPEQQ

15

 (c) murine BR3 sequence (SEQ ID NO: 35)

 1 MGARRLRVRS QRSRDSSVPT QCNQTECFDP LVRNCVSCEL FHTPDTGHTS SLEPGTALQP
 61 QEGSALRPDV ALLVGAPALL GLILALTLVG LVSLVSWRWR QQLRTASPDY SEGVQQESLE
 121 NVFVPSSETP HASAPTWPFLL KEDADSALPR HSPVPATEL GSTELVTTKT AGPEQ

20

 (d) rat BR3 sequence (SEQ ID NO: 49)

 1 MGVRRLRVRS RRSRDSPVST QCNQTECFDP LVRNCVSCEL FYTPETRHAS SLEPGTALQP
 61 QEGSGLRPDV ALLFGAPALL GLVLALTLVG LVSLVGWRWR QQRRTASLDY SEGVQQESLE
 121 NVFVPPSETL HASAPNWPFLL KEDADNLSL HSPVPATEL GSTELVTTKT AGPEQ

25

and variants or fragments thereof. The BR3 polypeptides of the invention can be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant and/or synthetic methods. The term BR3, includes the BR3 polypeptides described in WO 02/24909, WO 03/14294, and Fig. 8A of US Publication No. 2003/0099990 and WO 2003/024991 (polypeptides encoded by clones 711, 713, 706 and 710).

A "native sequence" BR3 polypeptide comprises a polypeptide having the same amino acid sequence as the corresponding BR3 polypeptide derived from nature. Such native sequence BR3 polypeptides can be isolated from nature or can be produced by recombinant and/or synthetic means. The term "native sequence BR3 polypeptide" specifically encompasses naturally-occurring truncated, soluble or secreted forms (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. The BR3 polypeptides of the invention include the BR3 polypeptide comprising or consisting of the contiguous sequence of amino acid residues 1 to 184 of a human BR3.

40 A BR3 "extracellular domain" or "ECD" refers to a form of the BR3 polypeptide which is essentially free of the transmembrane and cytoplasmic domains. ECD forms of BR3 include those comprising any one of amino acids 1 to 77, 2 to 62, 2-71, 1-61 and 2-63 of BR3.

Mini-BR3 is a 26-residue core region of the BAFF-binding domain of BR3:

TPCVPAECFD LLVRHCVACG LLRTPR (SEQ. ID: 50)

45 "BR3 variant" means a BR3 polypeptide having at least about 80% amino acid sequence identity with the amino acid sequence of a native sequence full length BR3 or BR3 ECD and binds a native sequence BAFF polypeptide. Optionally, the BR3 variant includes a single cysteine rich domain. Such BR3 variant polypeptides include, for instance, BR3 polypeptides wherein one or more amino acid residues are added, or deleted, at the N- and/or C-terminus, as well as within one or more internal domains, of the full-length amino acid sequence. BR3 variant polypeptides also include native sequence BR3 that have one or more residues substituted with residues from the corresponding position from another species, e.g., human BR3 with amino

50

5 acid substitutions of mouse BR3 residues, that can improve biological or physical properties of human BR3 including binding to BAFF or reduction in aggregation. See e.g., Fig. 20 of WO 02/24909. Fragments of the BR3 ECD that bind a native sequence BAFF polypeptide, such as a portion of the ECD that is shorter in length than residues 2-71 of human BR3 and comprises a minimal region of one of SEQ ID NO. 146-165 shown in Figure 33, are also contemplated. ECD and ECD fragments of polypeptides described in WO 2003/024991 that
 10 bind BAFF are also contemplated. BR3 variants also include the TALL-1 binding peptides described in WO 02/092620. Ordinarily, a BR3 variant polypeptide will have at least about 80% amino acid sequence identity, more preferably at least about 81% amino acid sequence identity, more preferably at least about 82% amino acid sequence identity, more preferably at least about 83% amino acid sequence identity, more preferably at least about 84% amino acid sequence identity, more preferably at least about 85% amino acid sequence identity,
 15 more preferably at least about 86% amino acid sequence identity, more preferably at least about 87% amino acid sequence identity, more preferably at least about 88% amino acid sequence identity, more preferably at least about 89% amino acid sequence identity, more preferably at least about 90% amino acid sequence identity, more preferably at least about 91% amino acid sequence identity, more preferably at least about 92% amino acid sequence identity, more preferably at least about 93% amino acid sequence identity, more preferably at least about 94% amino acid sequence identity, more preferably at least about 95% amino acid sequence identity, more preferably at least about 96% amino acid sequence identity, more preferably at least about 97% amino acid sequence identity, more preferably at least about 98% amino acid sequence identity and yet more preferably at least about 99% amino acid sequence identity with a human BR3 polypeptide or a specified fragment thereof. BR3 variant polypeptides do not encompass the native BR3 polypeptide sequence. Ordinarily, BR3 variant
 25 polypeptides are at least about 10 amino acids in length, often at least about 20 amino acids in length, more often at least about 30 amino acids in length, more often at least about 40 amino acids in length, more often at least about 50 amino acids in length, more often at least about 60 amino acids in length, more often at least about 70 amino acids in length, more often at least about 80 amino acids in length, more often at least about 90 amino acids in length, more often at least about 100 amino acids in length, more often at least about 150 amino acids in length, more often at least about 200 amino acids in length, more often at least about 250 amino acids in length, more often at least about 300 amino acids in length, or more.

The terms "TACI" or "TACI polypeptide" or "TACI receptor" when used herein encompass "native sequence TACI polypeptides" and "TACI variants" (which are further defined herein). "TACI" is a designation given to those polypeptides comprising the amino acid sequences of Figures 1A-1B, amino acids 1-246 of
 35 Figure 5B and the amino acid sequences of Figure 8, polypeptides which are encoded by nucleic acid molecules comprising the polynucleotide sequence shown in Figures 1A-1B and 5A and homologs, variants and fragments thereof, nucleic acid molecules comprising the sequence shown in the Figures 1A-1B and 5A and variants thereof as well as fragments of the above. The TACI polypeptides of the invention can be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant and/or
 40 synthetic methods.

A "native sequence" TACI polypeptide comprises a polypeptide having the same amino acid sequence as the corresponding TACI polypeptide derived from nature. Such native sequence TACI polypeptides can be isolated from nature or can be produced by recombinant and/or synthetic means. The term "native sequence TACI polypeptide" specifically encompasses naturally-occurring truncated, soluble or secreted forms (e.g., an

extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. The TACI polypeptides of the invention include but are not limited to the polypeptides described in von Bulow et al., *supra* and WO98/39361 published September 11, 1998, the spliced variant (referred to as "hTACI(265)" above and shown in Figure 8, the TACI polypeptide comprising the contiguous sequence of amino acid residues 1-293 of Figure 8).

A TACI "extracellular domain" or "ECD" refers to a form of the TACI polypeptide which is essentially free of the transmembrane and cytoplasmic domains. ECD forms of TACI include those described in von Bulow et al., *supra*, WO 98/39361, WO 00/40716, WO 01/85782, WO 01/87979, WO 01/81417, amino acid residues 1-166 of Figure 1, amino acid residues 1-165 of Figure 1, amino acid residues 1-154 of Figure 1, amino acid residues 1-114 of Figure 1, amino acid residues 1-119 of Figure 5B, amino acid residues 1-120 of Figure 5B, and amino acid residues 1-126 of Figure 5B.

"TACI variant" means a TACI polypeptide having at least about 80% amino acid sequence identity with the amino acid sequence of a native sequence full length TACI or TACI ECD and binds a native sequence BAFF polypeptide. Such TACI variant polypeptides include, for instance, TACI polypeptides wherein one or more amino acid residues are added, or deleted, at the N- and/or C-terminus, as well as within one or more internal domains, of the full-length amino acid sequence. Fragments of the TACI ECD that bind a native sequence BAFF polypeptide are also contemplated. Ordinarily, a TACI variant polypeptide will have at least about 80% amino acid sequence identity, more preferably at least about 81% amino acid sequence identity, more preferably at least about 82% amino acid sequence identity, more preferably at least about 83% amino acid sequence identity, more preferably at least about 84% amino acid sequence identity, more preferably at least about 85% amino acid sequence identity, more preferably at least about 86% amino acid sequence identity, more preferably at least about 87% amino acid sequence identity, more preferably at least about 88% amino acid sequence identity, more preferably at least about 89% amino acid sequence identity, more preferably at least about 90% amino acid sequence identity, more preferably at least about 91% amino acid sequence identity, more preferably at least about 92% amino acid sequence identity, more preferably at least about 93% amino acid sequence identity, more preferably at least about 94% amino acid sequence identity, more preferably at least about 95% amino acid sequence identity, more preferably at least about 96% amino acid sequence identity, more preferably at least about 97% amino acid sequence identity, more preferably at least about 98% amino acid sequence identity and yet more preferably at least about 99% amino acid sequence identity with a TACI polypeptide encoded by a nucleic acid molecule shown in Figure 1A or a specified fragment thereof. TACI variant polypeptides do not encompass the native TACI polypeptide sequence. Ordinarily, TACI variant polypeptides are at least about 10 amino acids in length, often at least about 20 amino acids in length, more often at least about 30 amino acids in length, more often at least about 40 amino acids in length, more often at least about 50 amino acids in length, more often at least about 60 amino acids in length, more often at least about 70 amino acids in length, more often at least about 80 amino acids in length, more often at least about 90 amino acids in length, more often at least about 100 amino acids in length, more often at least about 150 amino acids in length, more often at least about 200 amino acids in length, more often at least about 250 amino acids in length, more often at least about 300 amino acids in length, or more.

The terms "BCMA" or "BCMA polypeptide" or "BCMA receptor" when used herein encompass "native sequence BCMA polypeptides" and "BCMA variants" (which are further defined herein). "BCMA" is a

5 designation given to those polypeptides which are encoded by the nucleic acid molecules comprising the polynucleotide sequences shown in Figure 2 and variants thereof, nucleic acid molecules comprising the sequence shown in the Figure 2 and variants thereof as well as fragments of the above. The BCMA polypeptides of the invention can be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant and/or synthetic methods.

10 A "native sequence" BCMA polypeptide comprises a polypeptide having the same amino acid sequence as the corresponding BCMA polypeptide derived from nature. Such native sequence BCMA polypeptides can be isolated from nature or can be produced by recombinant and/or synthetic means. The term "native sequence BCMA polypeptide" specifically encompasses naturally-occurring truncated or secreted forms (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and
15 naturally-occurring allelic variants of the polypeptide. The BCMA polypeptides of the invention include the polypeptides described in Laabi et al., EMBO J., 11:3897-3904 (1992); Laabi et al., Nucleic Acids Res., 22:1147-1154 (1994); Gras et al., Int. Immunology, 7:1093-1106 (1995); Madry et al., Int. Immunology, 10:1693-1702 (1998); and the BCMA polypeptide comprising the contiguous sequence of amino acid residues 1-184 of Fig. 2 (SEQ ID NO: 27).

20 A BCMA "extracellular domain" or "ECD" refers to a form of the BCMA polypeptide which is essentially free of the transmembrane and cytoplasmic domains. ECD forms of TACI include those described in Laabi et al., EMBO J., 11:3897-3904 (1992); Laabi et al., Nucleic Acids Res., 22:1147-1154 (1994); Gras et al., Int. Immunology, 7:1093-1106 (1995); Madry et al., Int. Immunology, 10:1693-1702 (1998), amino acid residues 4-55 of Figure 2, amino acid residues 1-48 of Figure 2, amino acid residues 8-41 of Figure 2, amino
25 acid residues 4-51 of Figure 2 or amino acid residues 21-53 of Figure 2.

"BCMA variant" means a BCMA polypeptide having at least about 80% amino acid sequence identity with the amino acid sequence of a native sequence BCMA or BCMA ECD and binds a native sequence BAFF polypeptide. Such BCMA variant polypeptides include, for instance, BCMA polypeptides wherein one or more amino acid residues are added, or deleted, at the N- and/or C-terminus, as well as within one or more internal
30 domains, of the full-length amino acid sequence. Fragments of the BCMA ECD that bind a native sequence BAFF polypeptide are also contemplated. Ordinarily, a BCMA variant polypeptide will have at least about 80% amino acid sequence identity, more preferably at least about 81% amino acid sequence identity, more preferably at least about 82% amino acid sequence identity, more preferably at least about 83% amino acid sequence identity, more preferably at least about 84% amino acid sequence identity, more preferably at least about 85%
35 amino acid sequence identity, more preferably at least about 86% amino acid sequence identity, more preferably at least about 87% amino acid sequence identity, more preferably at least about 88% amino acid sequence identity, more preferably at least about 89% amino acid sequence identity, more preferably at least about 90% amino acid sequence identity, more preferably at least about 91% amino acid sequence identity, more preferably at least about 92% amino acid sequence identity, more preferably at least about 93% amino acid sequence
40 identity, more preferably at least about 94% amino acid sequence identity, more preferably at least about 95% amino acid sequence identity, more preferably at least about 96% amino acid sequence identity, more preferably at least about 97% amino acid sequence identity, more preferably at least about 98% amino acid sequence identity and yet more preferably at least about 99% amino acid sequence identity with a BCMA polypeptide encoded by a nucleic acid molecule shown in Figure 2 or a specified fragment thereof. BCMA variant

5 polypeptides do not encompass the native BCMA polypeptide sequence. Ordinarily, BCMA variant polypeptides are at least about 10 amino acids in length, often at least about 20 amino acids in length, more often at least about 30 amino acids in length, more often at least about 40 amino acids in length, more often at least about 50 amino acids in length, more often at least about 60 amino acids in length, more often at least about 70 amino acids in length, more often at least about 80 amino acids in length, more often at least about 90 amino acids in length, more often at least about 100 amino acids in length, more often at least about 150 amino acids in length, more often at least about 200 amino acids in length, more often at least about 250 amino acids in length, more often at least about 300 amino acids in length, or more.

BAFF is expressed in spleen, lymph nodes, PBLs, monocytes, macrophages, DCs, T cells, K562, HL-60 and G361. APRIL is weakly expressed in spleen, pancreas, colon. APRIL is expressed in PBLs and various tumor cell lines and tissues. BCMA is expressed in spleen, lymph nodes, thymus, PBLs, liver, adrenals and B cells. TACI is expressed in spleen, thymus, PBLs, small intestine, B cells and activated T cells. BAFF-R is expressed in spleen, lymph nodes, thymus, PBLs, B cells. BAFF-R is expressed in low levels on resting T cells. (Mackay and Browning, July 2002, Nature Reviews, Immunology, 2: 465-475).

Amino acids may be grouped according to similarities in the properties of their side chains (in A. L. Lehninger, in Biochemistry, second ed., pp. 73-75, Worth Publishers, New York (1975)):

(1) non-polar: Ala (A), Val (V), Leu (L), Ile (I), Pro (P), Phe (F), Trp (W), Met (M)

(2) uncharged polar: Gly (G), Ser (S), Thr (T), Cys (C), Tyr (Y), Asn (N), Gln (Q)

(3) acidic: Asp (D), Glu (E)

(4) basic: Lys (K), Arg (R), His(H)

25 Alternatively, naturally occurring residues may be divided into groups based on common side-chain properties:

(1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;

(2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;

(3) acidic: Asp, Glu;

30 (4) basic: His, Lys, Arg;

(5) residues that influence chain orientation: Gly, Pro;

(6) aromatic: Trp, Tyr, Phe.

The term "conservative" amino acid substitution as used within this invention is meant to refer to amino acid substitutions which substitute functionally equivalent amino acids. Conservative amino acid changes result in silent changes in the amino acid sequence of the resulting peptide. For example, one or more amino acids of a similar polarity act as functional equivalents and result in a silent alteration within the amino acid sequence of the peptide. In general, substitutions within a group may be considered conservative with respect to structure and function. However, the skilled artisan will recognize that the role of a particular residue is determined by its context within the three-dimensional structure of the molecule in which it occurs. For example, Cys residues may occur in the oxidized (disulfide) form, which is less polar than the reduced (thiol) form. The long aliphatic portion of the Arg side chain may constitute a critical feature of its structural or functional role, and this may be best conserved by substitution of a nonpolar, rather than another basic residue. Also, it will be recognized that side chains containing aromatic groups (Trp, Tyr, and Phe) can participate in ionic-aromatic or "cation-pi" interactions. In these cases, substitution of one of these side chains with a member

5 of the acidic or uncharged polar group may be conservative with respect to structure and function. Residues such as Pro, Gly, and Cys (disulfide form) can have direct effects on the main chain conformation, and often may not be substituted without structural distortions.

"Percent (%) amino acid sequence identity" with respect to the ligand or receptor polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in such a ligand or receptor sequence identified herein, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in the table below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in the table below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or can be compiled from the source code provided in the table below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D.

25 All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

A useful method for identification of certain residues or regions in a protein that are preferred locations for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham and Wells *Science*, 244:1081-1085 (1989). A residue or group of target residues are identified (*e.g.*, charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with a binding target. Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* need not be predetermined. For example, to analyze the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed variants are screened for the desired activity.

The term, "dihedral angle" refers to a rotation about a bond. See *e.g.*, Creighton, T.E., (1993) *Protein: Structures and Molecular Properties*, 2 ed., W. H. Freeman and Company, New York, NY.

The term, "phi," is a dihedral angle that denotes a rotation about the N-C α bond of an amino acid. See *e.g.*, Creighton, T.E., (1993) *Protein: Structures and Molecular Properties*, 2 ed., W. H. Freeman and Company, New York, NY.

Type I beta turns are described in Hutchinson, E. G. & Thornton, J. M. (1994) A revised set of potentials for beta turn formation in proteins. *Protein Science* 3, 2207-2216.

A "fusion protein" and a "fusion polypeptide" refer to a polypeptide having two portions covalently linked together, where each of the portions is a polypeptide having a different property. The property may be a

5 biological property, such as activity *in vitro* or *in vivo*. The property may also be a simple chemical or physical property, such as binding to a target molecule, catalysis of a reaction, etc. The two portions may be linked directly by a single peptide bond or through a peptide linker containing one or more amino acid residues. Generally, the two portions and the linker will be in reading frame with each other.

A "conjugate" refers to any hybrid molecule, including fusion proteins and as well as molecules that contain both amino acid or protein portions and non-protein portions. Conjugates may be synthesized by a variety of techniques known in the art including, for example, recombinant DNA techniques, solid phase synthesis, solution phase synthesis, organic chemical synthetic techniques or a combination of these techniques. The choice of synthesis will depend upon the particular molecule to be generated. For example, a hybrid molecule not entirely "protein" in nature may be synthesized by a combination of recombinant techniques and solution phase techniques.

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin can be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM. For example, useful immunoadhesins according to this invention are polypeptides that comprise the BAFF binding portions of a BAFF receptor without the transmembrane or cytoplasmic sequences of the BAFF receptor. In one embodiment, the extracellular domain of BR3, TACI or BCMA is fused to a constant domain of an immunoglobulin sequence. For example, a mouse BR3-Fc immunoadhesin and human BR3-Fc immunoadhesin according to this invention can be represented by the formulae:

mBR3-Fc (SEQ ID NO. 1):

MSALLILALVGAAVASTGARRLRVRSQRSDSSVPTQCNCQTECFDPLVRNCVSCELFHTPDTGHTSSLE
PGTALQPQEGQVTGDKKIVPRDCGCKPCICTVPEVSSVFIFPPKPKDVLTTITLTPKVTCVVVDISKDDPEV
QFSWFVDDVEVHTAQTQPREEQFNSTFRSVSEL
PIMHQDWLNGKEFKCRVNSAAFPAPIEKTISKTKGRPKAPQVYTIPPPKEQMAKD KVS LTCMITDFFPE
DITVEWQWNGQPAENYKNTQPIMN TNGSYFVYSKLVNQKSNWEAGNTFTCSVLHEGLHNHHTKSL
HSPGK

hBR3-hIgG1 Fc (SEQ ID NO. 2)

MSALLILALVGAAVASTRRGPRSLRGRDAPAPTPCVPABCFDPLVRHCVACGLLRTPRKPAGASSPAP
RTALQPQESQVTDKAAHYTLCPPCAPELLGGPSVFLFPPKPKD/TLMISRTPVTCVVAVSHEDPEVK
FNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKG
QPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKL
TVDKSRWQQGNVFSQSVMEALHNHYTQKSLSLSPGK

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXANTM); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine,

- 5 triethylenephosphoramidate, triethylenethiophosphoramidate and trimethylolmelamine; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, anthramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, 10 detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine 15 analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone propionate, epitostanol, mepitiostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; 20 bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK®; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2, 2', 2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); 25 cyclophosphamide; thiotepe; taxoids, *e.g.* paclitaxel (TAXOL®, Bristol-Myers Squibb Oncology, Princeton, NJ) and doxetaxel (TAXOTERE®, Rhône-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 30 2000; difluoromethylornithine (DMFO); retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and toremifene (Fareston); and anti-androgens such as flutamide, nilutamide, 35 bicalutamide, leuprolide, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (*e.g.* At^{211} , I^{131} , I^{125} , Y^{90} , Re^{186} , Re^{188} , Sm^{153} , Bi^{212} , P^{32} and radioactive isotopes of Lu), chemotherapeutic agents *e.g.* methotrexate, 40 adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), doxorubicin, melphalan, mitomycin C, chlorambucil, daunorubicin or other intercalating agents, enzymes and fragments thereof such as nucleolytic enzymes, antibiotics, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof, and the various antitumor or anticancer agents disclosed below. Other cytotoxic agents are described below.

5 A "growth inhibitory agent" when used herein refers to a compound or composition which inhibits growth of a cell, especially a CD20 expressing cancer cell, either *in vitro* or *in vivo*. Thus, the growth inhibitory agent may be one which significantly reduces the percentage of PSCA expressing cells in S phase. Examples of growth inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and
10 vinblastine), taxanes, and topoisomerase II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in *The Molecular Basis of Cancer*, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogenes, and antineoplastic drugs" by Murakami *et al.* (WB
15 Saunders: Philadelphia, 1995), especially p. 13. The taxanes (paclitaxel and docetaxel) are anticancer drugs both derived from the yew tree. Docetaxel (TAXOTERE®, Rhone-Poulenc Rorer), derived from the European yew, is a semisynthetic analogue of paclitaxel (TAXOL®, Bristol-Myers Squibb). Paclitaxel and docetaxel promote the assembly of microtubules from tubulin dimers and stabilize microtubules by preventing depolymerization, which results in the inhibition of mitosis in cells.

20 The term "mammal" refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, etc. Preferably, the mammal herein is human.

The term "therapeutically effective amount" refers to an amount of an antibody or a antagonist drug effective to "alleviate" or "treat" a disease or disorder in a subject or mammal. In the case of cancer, the
25 therapeutically effective amount of the drug may reduce the number of cancer cells; reduce the tumor size; inhibit (*i.e.*, slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (*i.e.*, slow to some extent and preferably stop) tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the cancer. See the definition of "treated" below. To the extent the drug may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or
30 cytotoxic.

The anti-CD20 antibodies of the invention can be produced by transient or stable transfection eukaryotic host cells such as CHO cells.

"Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often
35 the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including
40 glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, polyethylene glycol (PEG), and PLURONICS™.

5 1. Polypeptide BAFF Antagonists

Polypeptides useful as antagonists of BAFF include, e.g., the polypeptide referred to as TALL-1 12-3 as SEQ ID No. 123 in WO 02/092620, the amino acid sequence provided here:

MLPGCKWDLLIKQWVCDPLGSGSATGGSGSTASSGSGSATHMLPGCKWDLLIKQWVCDPLGGGGGV
DKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCWWDVSHEDPEVKFNWYVDGVEVHNAKT
10 KPREEQYNSTYRWSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL
TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSC
VMHEALHNHYTQKSLSLSPGK (SEQ ID NO. 10)

This polypeptide binds BAFF and inhibits BR3 binding to BAFF.

In some embodiments, the 17-mer peptides are soluble (preferably not membrane bound), and may be used as
15 core sequences or otherwise combined or conjugated with a variety of structures as is described below. Some
amino acids in the 17-mer polypeptide were randomized and screened for conservative and non-conservative
substitutions. As is understood by one of skill in the art and described herein, additions and substitutions may
be accomplished on a limited basis without impairing the BAFF binding of the resulting 17mer peptide and
constructs including the resulting 17mer peptide. Guidance as to allowed substitutions that yield BAFF binding
20 function is provided below and in the examples. In some embodiments, residues implicated in structural or
binding affinity relationships are conserved, meaning that either the amino acid identity is retained or a
conservative substitution is made as described in the formulas and description below.

A BAFF polypeptide antagonist of this invention comprises a sequence selected from the group
consisting of: Formula I, Formula II, Formula III, a sequence recited in FIG.32, ECFDLLVRAWVPCSVLK
25 (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7),
ECFDLLVRSWVPCHMLR (SEQ ID NO 8), ECFDLLVRHWVACGLLR (SEQ ID NO 9), and mixtures
thereof.

In one aspect of the invention, the polypeptide comprises an amino acid sequence of Formula I:

$X_1-C_N-X_3-D-X_5-L-X_7-X_8-X_9-X_{10}-X_{11}-X_{12}-C_T-X_{14}-X_{15}-X_{16}-X_{17}$ (Formula I) (SEQ ID NO: 143)
30 wherein X_1 , X_3 , X_5 , X_7 , X_8 , X_9 , X_{10} , X_{11} , X_{12} , X_{14} , X_{15} and X_{17} are any amino acid except cysteine; and
wherein X_{16} is an amino acid selected from the group consisting of L, F, I and V; and
wherein the polypeptide does not comprise a cysteine within seven amino acid residues N-terminal to
 C_N (cysteine N terminal) and C-terminal to C_T (cysteine C terminal) of Formula I.

In some embodiments, a polypeptide comprising the sequence of Formula I has C_N and C_T joined by
35 disulfide bonding; $X_5LX_7X_8$ forming the conformation of a type I beta turn structure with the center of the turn
between L and X_7 ; and has a positive value for the dihedral angle phi of X_8 .

In some embodiments, X_{10} is selected from the group consisting of W, F, V, L, I, Y, M and a non-polar
amino acid. In some embodiments, X_{10} is W. In some embodiments, X_3 is an amino acid selected from the
group consisting of M, V, L, I, Y, F, W and a non-polar amino acid. In some embodiments, X_5 is selected from
40 the group consisting of V, L, P, S, I, A and R. In some embodiments, X_7 is selected from the group consisting
of V, T, I and L. In some embodiments, X_7 is not T or I. In some embodiments, X_8 is selected from the group
consisting of any R, K, G, N, H and a D-amino acid. In some embodiments, X_9 is selected from the group
consisting of H, K, A, R and Q. In some embodiments, X_{11} is I or V. In some embodiments, X_{12} is selected
from the group consisting of P, A, D, E and S. In some embodiments, X_{16} is L.

5 In specific embodiments, the sequence of Formula I is a sequence selected from the group consisting of ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCHMLR (SEQ ID NO 8), ECFDLLVRHWVACGLLR (SEQ ID NO 9).

10 In another aspect of the invention, the polypeptide comprises an amino acid sequence of Formula II:
 $X_1-C_N-X_3-D-X_5-L-V-X_8-X_9-W-X_{11}-X_{12}-C_T-X_{14}-X_{15}-L-X_{17}$ (Formula II) (SEQ ID NO: 144)
 wherein X_1 , X_3 , X_5 , X_8 , X_9 , X_{11} , X_{12} , X_{14} , X_{15} and X_{17} are any amino acid, except cysteine; and wherein the polypeptide does not comprise a cysteine within seven amino acid residues N-terminal to C_N (cysteine N terminal) and C-terminal to C_T (cysteine C terminal) of Formula II.

15 In some embodiments, a polypeptide comprising the sequence of Formula I has C_N and C_T joined by disulfide bonding; X_5LVX_8 forming the conformation of a type I beta turn structure with the center of the turn between L and X_7 ; and has a positive value for the dihedral angle phi of X_8 .

In some embodiments of Formula II, X_3 is an amino acid selected from the group consisting of M, A, V, L, I, Y, F, W and a non-polar amino acid. In some embodiments of Formula II, X_5 is selected from the group consisting of V, L, P, S, I, A and R. In some embodiments of Formula II, X_8 is selected from the group consisting of R, K, G, N, H and a D-amino acid. In some embodiments of Formula II, X_9 is selected from the group consisting of H, K, A, R and Q. In some embodiments of Formula II, X_{11} is selected from the group consisting of I and V. In some embodiments of Formula II, X_{12} is selected from the group consisting of P, A, D, E and S.

25 In other aspects, the polypeptide comprises a sequence selected from any one of the sequences described in FIG.32.

Another aspect of the invention includes a polypeptide comprising an amino acid sequence of Formula III:

30 $E-C_N-F-D-X_5-L-V-X_8-X_9-W-V-X_{12}-C_T-X_{14}-X_{15}-X_{16}-X_{17}$ (Formula III) (SEQ ID NO: 145)
 wherein X_5 , X_8 , X_9 , X_{12} , X_{14} , X_{15} and X_{17} are any amino acid except cysteine;
 wherein X_{16} is an amino acid selected from the group consisting of L, F, I and V;
 wherein the polypeptide does not comprise a cysteine within seven amino acid residues N-terminal to C_N (cysteine N terminal) and C-terminal to C_T (cysteine C terminal) of Formula III; and
 wherein C_N and C_T are joined by disulfide bonding.

35 In some embodiments of Formula III, the polypeptide comprising the contiguous sequence of Formula III has a disulfide bond between C_N and C_T and forms a type I beta turn structure with the center of the turn between L and V at X_5LVX_8 ; and has a positive value for the dihedral angle phi of X_8 .

40 In some embodiments of Formula III, X_5 , X_8 , X_9 , X_{12} , X_{14} , X_{15} and X_{17} are selected from the group consisting of L, P, H, R, I, T, N, S, V, A, D, and G. In some embodiments of Formula III, X_5 is L and X_8 is R. In some embodiments of Formula III, X_9 is selected from the group consisting of H, K, A, S, R and Q. In some embodiments of Formula III, X_{12} is selected from the group consisting of P, A, D, E and S. In some embodiments of Formula III, X_{12} is P. In some embodiments of Formula III, X_{16} is L.

In specific embodiments, the sequence of Formula III is selected from the group consisting of ECFDLLVRAWVPCSVLK (SEQ ID NO. 5), ECFDLLVRHWVPCGLLR (SEQ ID NO. 6),

5 ECFDLLVRRWVPCMLG (SEQ ID NO. 7), ECFDLLVRSWVPCHMLR (SEQ ID NO. 8) and ECFDLLVRHWVACGLLR (SEQ ID NO. 9).

Also included is a polypeptide comprising a contiguous polypeptide sequence selected from the group consisting of ECFDLLVRAWVPCSVLK, ECFDLLVRHWVPCGLLR, ECFDLLVRRWVPCMLG, ECFDLLVRSWVPCHMLR, and ECFDLLVRHWVACGLLR (SEQ ID NO. 5 TO 9). The present invention
10 also relates to a polypeptide comprising a sequence selected from any one of the sequences described in FIG. 32. Polypeptides comprising any one of the sequences described in FIG. 32 preferably join the cysteines of the sequence by disulfide bonding. In some embodiments, the sequence between the fifth and eighth residues of the sequence forms a conformation of a type I beta turn structure with the center of the turn between L and X₇ and the eighth residue has a positive value for the dihedral angle phi.

15 In some embodiments, the BAFF polypeptides of this invention are contiguous sequences. The present invention also relates to a polypeptide comprising at least 88% sequence identity with a contiguous 17mer polypeptide sequence selected from the group consisting of: ECFDLLVRAWVPCSVLK, ECFDLLVRHWVPCGLLR, ECFDLLVRRWVPCMLG, ECFDLLVRSWVPCHMLR, and ECFDLLVRHWVACGLLR (SEQ ID NOs 5-9). In other embodiments sequence identity is at least 64 %, and
20 each successive integer to 100% after aligning to provide maximum homology. Homology is reduced for sequence gaps and sequences shorter than the 17mers of the present invention after aligning to provide maximum homology. Neither N-nor C-terminal extensions nor insertions shall be construed as reducing homology.

According to some embodiments of this invention, the polypeptide is less than 50 amino acids in
25 length, less than 25 amino acids in length, or is a 17-mer.

In some embodiments, the polypeptides of this invention comprise additional polypeptide sequences N-terminal, C-terminal or both N-terminal and C-terminal to the sequence of Formula I, Formula II, Formula III, ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCHMLR (SEQ ID NO 8),
30 ECFDLLVRHWVACGLLR (SEQ ID NO 9), or sequences listed in FIG.32. The additional polypeptide sequences are heterologous to a native sequence BR3 polypeptide, and include, for example, Fc portion of immunoglobulins.

In another aspect of the invention, the BAFF antagonist polypeptides comprise at least one and more preferably, more than one of a polypeptide comprising a sequence of Formula I, Formula II, Formula III,
35 ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCHMLR (SEQ ID NO 8), ECFDLLVRHWVACGLLR (SEQ ID NO 9), or sequences listed in FIG. 32. The polypeptides that are linked together can have the same sequence or have different sequences. In some embodiments, these polypeptides can be joined to one another, optionally, through the use of a linker. The linker serves as a spacer and can be made
40 of a variety of chemical compounds. In some embodiments, the linker is a polypeptide that has about 1 to 50 amino acids, more preferably about 1 to 30 amino acids.. Linker sequences are known to those of skill in the art. For example, linker sequences include GGGKGGGG and GGGNSSGG and the like. In specific embodiments, the polypeptides linked together have the same sequence and comprise a formula: PP1-L1-PP1-L2-PP1, wherein PP1 comprises an amino acid sequence selected from the group consisting of Formula I, Formula II, Formula

5 III, ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6),
ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCMLR (SEQ ID NO 8),
ECFDLLVRHWVACGLLR (SEQ ID NO 9), and sequences listed in FIG.32, and L1 and L2 are linker
sequences that are different in sequence.

Antagonists for BAFF binding to BR3, such as the polypeptides described herein, preferably bind to
10 BAFF with an affinity the same as or greater than a native BR3 sequence, such as BR3 ECD of SEQ ID NO: 51
or mini-BR3 of SEQ ID NO: 50. In some embodiments, the polypeptides having a sequence of that of Formula
I, Formula II, Formula III, ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID
NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCMLR (SEQ ID NO 8),
ECFDLLVRHWVACGLLR (SEQ ID NO 9), or sequences listed in FIG.32 have a binding affinity for BAFF of
15 about 100nM or less, preferably 10 nM or less, or 1 nM or less. One method of measuring binding affinity is
provided in the Examples.

A method used in the present invention to find BAFF antagonists involves identifying, modifying and
selectively randomizing a core sequence of 17 residues of BR3. Specific techniques used are described further
below and in the examples. In some embodiments, the N terminal cysteine residue (C_N) at position X_2 and C-
20 terminal cysteine (C_T) at position X_{13} are conserved and preferably form a disulfide bridge. In some
embodiments, C_N and C_T are separated by 10 contiguous amino acids. Preferably, the 17mer sequence does not
contain any cysteine residues other than at positions X_2 and X_{13} . Additionally, if the 17mer is included in a
larger structure, sequences flanking the 17mer will preferably not include any cysteine residues within 7 amino
acids of C_N or C_T . X_{10} is substituted with any non-polar amino acid except for cysteine; for example: W, F, V,
25 L, I, Y or M. In some embodiments, X_{10} is W.

In some embodiments, the motif D- X_5 -L- X_7 is conserved due to demonstrated contribution to BAFF
binding. In some embodiments, a beta-turn located between C_N and C_T , is formed between X_4 and X_7 . In some
embodiments, the center of the beta-turn is positioned between L- X_7 . In some embodiments, the structure of the
17mer peptides of the present invention is generally two beta-strands linked by a type I beta-turn, forming a
30 beta-hairpin connected by a disulfide bond between C_N and C_T . Additionally, in some embodiments, the residue
at X_8 adopts a positive value for the dihedral angle phi of X_8 to accommodate the type I beta turn in the beta
hairpin structure.

Additional structural information indicates that D at X_4 and L at X_6 are buried in the BAFF-BR3
interface of BAFF-BR3 complex. In some embodiments, these residues are conserved. The residue at X_7 is also
35 buried in the BAFF-BR3 interface. In some embodiments, X_7 may be selected from the group consisting of V,
T, I and L. In some embodiments, X_7 is preferably V. In some embodiments, the motif from X_4 to X_7 is DLLV.

In some embodiments, the length of the binding region of the BAFF antagonist is 17 amino acids. In
some embodiments, the polypeptide BAFF antagonist is 17 amino acids. In some embodiments, four amino
acids, X_{14} - X_{17} , follow C_T at the C-terminal end. In some embodiments, X_{16} forms a hydrophobic contact with
40 BAFF when the 17mer is bound and is conserved. In some embodiments X_{16} is L.

5 2. Polynucleotides, Vectors, Host Cells

According to some embodiments, the BAFF antagonist polypeptides of this invention are selected from the group consisting of: 17mer peptides described herein, polypeptides incorporating one or more 17mer peptides as core regions, and covalently modified forms of the 17mer peptides and polypeptides (e.g., immunoadhesins, labeled polypeptides, protected polypeptides, conjugated polypeptides, fusion proteins, etc.).
10 Various techniques that are employed for making these forms of polypeptides are described herein. Methods for labeling polypeptides and conjugating molecules to polypeptides are known in the art.

Compositions of the invention can be prepared using recombinant techniques known in the art. The description below relates to methods of producing such polypeptides by culturing host cells transformed or transfected with a vector containing the encoding nucleic acid and recovering the polypeptide from the cell
15 culture. (See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual* (New York: Cold Spring Harbor Laboratory Press, 1989); Dieffenbach et al., *PCR Primer: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1995)).

The nucleic acid (e.g., cDNA or genomic DNA) encoding the desired polypeptide may be inserted into a replicable vector for further cloning (amplification of the DNA) or for expression. Various vectors are
20 publicly available. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence, each of which is described below. Optional signal sequences, origins of replication, marker genes, enhancer elements and transcription terminator sequences that may be employed are known in the art and described in further detail in WO97/25428.

25 Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the encoding nucleic acid sequence. Promoters are untranslated sequences located upstream (5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of a particular nucleic acid sequence, to which they are operably linked. Such promoters typically fall into two classes, inducible and constitutive. Inducible promoters are promoters that
30 initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, e.g., the presence or absence of a nutrient or a change in temperature. At this time a large number of promoters recognized by a variety of potential host cells are well known. These promoters are operably linked to the encoding DNA by removing the promoter from the source DNA by restriction enzyme digestion and inserting the isolated promoter sequence into the vector.

35 Construction of suitable vectors containing one or more of the above-listed components employs standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and re-ligated in the form desired to generate the plasmids required. For analysis to confirm correct sequences in plasmids constructed, the ligation mixtures can be used to transform *E. coli* K12 strain 294 (ATCC 31,446) and successful transformants selected by ampicillin or tetracycline resistance where appropriate. Plasmids from the
40 transformants are prepared, analyzed by restriction endonuclease digestion, and/or sequenced using standard techniques known in the art. [See, e.g., Messing et al., *Nucleic Acids Res.*, 9:309 (1981); Maxam et al., *Methods in Enzymology*, 65:499 (1980)].

5 Expression vectors that provide for the transient expression in mammalian cells of the encoding DNA may be employed. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of a desired polypeptide encoded by the expression vector [Sambrook et al., supra]. Transient expression systems, comprising a suitable expression vector and a host cell, allow for the
10 convenient positive identification of polypeptides encoded by cloned DNAs, as well as for the rapid screening of such polypeptides for desired biological or physiological properties.

 Other methods, vectors, and host cells suitable for adaptation to the synthesis of the desired polypeptide in recombinant vertebrate cell culture are described in Gething et al., *Nature*, 293:620-625 (1981); Mantei et al., *Nature*, 281:40-46 (1979); EP 117,060; and EP 117,058.

15 Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes for this purpose include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as Bacilli such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P
20 disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. Preferably, the host cell should secrete minimal amounts of proteolytic enzymes.

 In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for vectors. Suitable host cells for the expression of glycosylated polypeptide are derived from multicellular organisms. Examples of all such host cells are described further in WO97/25428.

25 Host cells are transfected and preferably transformed with the above-described expression or cloning vectors and cultured in nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

 Transfection refers to the taking up of an expression vector by a host cell whether or not any coding sequences are in fact expressed. Numerous methods of transfection are known to the ordinarily skilled artisan, for example, CaPO₄ and electroporation. Successful transfection is generally recognized when any indication
30 of the operation of this vector occurs within the host cell.

 Transformation means introducing DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integrant. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as
35 described in Sambrook et al., supra, or electroporation is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., *Gene*, 23:315 (1983) and WO 89/05859 published 29 June 1989. In addition, plants may be transfected using ultrasound treatment as described in WO 91/00358 published 10
January 1991.

40 For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, *Virology*, 52:456-457 (1978) may be employed. General aspects of mammalian cell host system transformations have been described in U.S. Pat. No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., *J. Bact.*, 130:946 (1977) and Hsiao et al., *Proc. Natl. Acad. Sci. (USA)*, 76:3829 (1979). However, other methods for introducing DNA into cells, such as

5 by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., *Methods in Enzymology*, 185:527-537 (1990) and Mansour et al., *Nature*, 336:348-352 (1988).

Prokaryotic cells can be cultured in suitable culture media as described generally in Sambrook et al., *supra*. Examples of commercially available culture media include Ham's F10 (Sigma), Minimal Essential
 10 Medium ("MEM", Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ("DMEM", Sigma). Any such media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleosides (such as adenosine and thymidine), antibiotics (such as gentamycin), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and
 15 glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

In general, principles, protocols, and practical techniques for maximizing the productivity of
 20 mammalian cell cultures can be found in *Mammalian Cell Biotechnology: A Practical Approach*, M. Butler, ed. (IRL Press, 1991).

The expressed polypeptides may be recovered from the culture medium as a secreted polypeptide, although may also be recovered from host cell lysates when directly produced without a secretory signal. If the polypeptide is membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g.
 25 Triton-X 100) or its extracellular region may be released by enzymatic cleavage.

When the polypeptide is produced in a recombinant cell other than one of human origin, it is free of proteins or polypeptides of human origin. However, it is usually necessary to recover or purify the polypeptide from recombinant cell proteins or polypeptides to obtain preparations that are substantially homogeneous. As a first step, the culture medium or lysate may be centrifuged to remove particulate cell debris. The following are
 30 procedures exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; and protein A Sepharose columns to remove contaminants such as IgG.

35 Phage Display

According to some embodiments, the polypeptides of this invention selected from the group consisting of: Formula I, Formula II, Formula III, ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCMLR (SEQ ID NO 8), ECFDLLVRHWVACGLLR (SEQ ID NO 9), and sequences
 40 listed in FIG.32, may be utilized in phage display.

Using the techniques of phage display allows the generation of large libraries of protein variants which can be rapidly sorted for those sequences that bind to a target molecule with high affinity. Nucleic acids encoding variant polypeptides are fused to a nucleic acid sequence encoding a viral coat protein, such as the gene III protein or the gene VIII protein. Monovalent phage display systems where the nucleic acid sequence

5 encoding the protein or polypeptide is fused to a nucleic acid sequence encoding a portion of the gene III protein have been developed. (Bass, S., *Proteins*, 8:309 (1990); Lowman and Wells, *Methods: A Companion to Methods in Enzymology*, 3:205 (1991)). In a monovalent phage display system, the gene fusion is expressed at low levels and wild type gene III proteins are also expressed so that infectivity of the particles is retained. Methods of generating peptide libraries and screening those libraries have been disclosed in many patents (e.g. 10 U.S. Patent No. 5,723,286, U.S. Patent No. 5,432, 018, U.S. Patent No. 5,580,717, U.S. Patent No. 5,427,908 and U.S. Patent No. 5,498,530).

In some embodiments, Formula I, Formula II or Formula III are expressed as peptide libraries on phage. The phage expressing the library of polypeptides of Formula I, Formula II or Formula III are then subjected to selection based on BAFF binding. In some embodiments, the selection process involves allowing 15 some phage bind to biotinylated BAFF which is subsequently bound to a neutravidin plate. Phage bound to the plate through the BAFF-biotin-neutravidin binding are recovered and propagated. In some embodiments, the phage are subject to several rounds of selection. In some embodiments, the phage is incubated with BAFF-biotin, followed by the addition of unbiotinylated BAFF as a competitive binder. Additional guidance of use of phage display in the context of the present invention is provided in the Examples.

20

Polypeptides fused or conjugated to Heterologous polypeptides

Immunoadhesin molecules comprising the polypeptides of this invention are further contemplated for use in the methods herein. In some embodiments, the molecule comprises a fusion of a polypeptide of this invention with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the 25 immunoadhesin, such a fusion usefully comprises the Fc region of an IgG molecule. In a further embodiment, the Fc region is from a human IgG1 molecule. In some embodiments, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions, see also US Patent No. 5,428,130 issued June 27, 1995 and Chamow et al., *TIBTECH*, 14:52-60 (1996).

30 The simplest and most straightforward immunoadhesin design often combines the binding domain(s) of the adhesin (e.g. antagonist polypeptide of this invention) with the Fc region of an immunoglobulin heavy chain. For example, a polypeptide comprising a sequence of Formula I, Formula II, Formula III, ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCHMLR (SEQ ID NO 8), 35 ECFDLLVRHWVACGLLR (SEQ ID NO 9), or sequences listed in FIG.32 can be covalently linked to an Fc portion of an immunoglobulin. In addition, one or more of these polypeptides can be linked to one another and linked to an Fc portion of an immunoglobulin.

Ordinarily, when preparing the immunoadhesins of the present invention, nucleic acid encoding the binding domain of the adhesin will be fused C-terminally to nucleic acid encoding the N-terminus of an 40 immunoglobulin constant domain sequence, however N-terminal fusions are also possible.

Typically, in such fusions the encoded chimeric polypeptide will retain at least functionally active hinge, CH2 and CH3 domains of the constant region of an immunoglobulin heavy chain. Fusions are also made to the C-terminus of the Fc portion of a constant domain, or immediately N-terminal to the CH1 of the heavy

5 chain or the corresponding region of the light chain. The precise site at which the fusion is made is not critical; particular sites are well known and may be selected in order to optimize the biological activity, secretion, or binding characteristics of the immunoadhesin.

In a preferred embodiment, the adhesin sequence is fused to the N-terminus of the Fc region of immunoglobulin G1 (IgG1). It is possible to fuse the entire heavy chain constant region to the adhesin
 10 sequence. However, more preferably, a sequence beginning in the hinge region just upstream of the papain cleavage site which defines IgG Fc chemically (i.e. residue 216, taking the first residue of heavy chain constant region to be 114), or analogous sites of other immunoglobulins is used in the fusion. In a particularly preferred embodiment, the adhesin amino acid sequence is fused to (a) the hinge region and CH2 and CH3 or (b) the CH1, hinge, CH2 and CH3 domains, of an IgG heavy chain.

15 For bispecific immunoadhesins, the immunoadhesins are assembled as multimers, and particularly as heterodimers or heterotetramers. Generally, these assembled immunoglobulins will have known unit structures. A basic four chain structural unit is the form in which IgG, IgD, and IgE exist. A four chain unit is repeated in the higher molecular weight immunoglobulins; IgM generally exists as a pentamer of four basic units held together by disulfide bonds. IgA globulin, and occasionally IgG globulin, may also exist in multimeric form in
 20 serum. In the case of multimer, each of the four units may be the same or different.

Various exemplary assembled immunoadhesins within the scope herein are schematically diagrammed below:

- (a) ACL-ACL;
- (b) ACH-(ACH, ACL-ACH, ACL-VHCH, or VLCL-ACH);
- 25 (c) ACL-ACH-(ACL-ACH, ACL-VHCH, VLCL-ACH, or VLCL-VHCH)
- (d) ACL-VHCH-(ACH, or ACL-VHCH, or VLCL-ACH);
- (e) VLCL-ACH-(ACL-VHCH, or VLCL-ACH); and
- (f) (A-Y)_n-(VLCL-VHCH)₂,

wherein each A represents identical or different polypeptides comprising an amino acid sequence of Formula I,
 30 Formula II, Formula III, ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCHMLR (SEQ ID NO 8), ECFDLLVRHWVACGLLR (SEQ ID NO 9), or sequences listed in FIG.32 or combinations thereof;

VL is an immunoglobulin light chain variable domain;

VH is an immunoglobulin heavy chain variable domain;

35 CL is an immunoglobulin light chain constant domain;

CH is an immunoglobulin heavy chain constant domain;

n is an integer greater than 1;

Y designates the residue of a covalent cross-linking agent.

In the interests of brevity, the foregoing structures only show key features; they do not indicate joining
 40 (J) or other domains of the immunoglobulins, nor are disulfide bonds shown. However, where such domains are required for binding activity, they shall be constructed to be present in the ordinary locations which they occupy in the immunoglobulin molecules.

Alternatively, the adhesin sequences can be inserted between immunoglobulin heavy chain and light chain sequences, such that an immunoglobulin comprising a chimeric heavy chain is obtained. In this

embodiment, the adhesin sequences are fused to the 3' end of an immunoglobulin heavy chain in each arm of an immunoglobulin, either between the hinge and the CH2 domain, or between the CH2 and CH3 domains.

Similar constructs have been reported by Hoogenboom et al., *Mol. Immunol.*, 28:1027-1037 (1991).

Although the presence of an immunoglobulin light chain is not required in the immunoadhesins of the present invention, an immunoglobulin light chain might be present either covalently associated to an adhesin-immunoglobulin heavy chain fusion polypeptide, or directly fused to the adhesin. In the former case, DNA encoding an immunoglobulin light chain is typically coexpressed with the DNA encoding the adhesin-immunoglobulin heavy chain fusion protein. Upon secretion, the hybrid heavy chain and the light chain will be covalently associated to provide an immunoglobulin-like structure comprising two disulfide-linked immunoglobulin heavy chain-light chain pairs. Methods suitable for the preparation of such structures are, for example, disclosed in U.S. Patent No. 4,816,567, issued 28 March 1989.

Immunoadhesins are most conveniently constructed by fusing the cDNA sequence encoding the adhesin portion in-frame to an immunoglobulin cDNA sequence. However, fusion to genomic immunoglobulin fragments can also be used (see, e.g. Aruffo et al., *Cell*, 61:1303-1313 (1990); and Stamenkovic et al., *Cell*, 66:1133-1144 (1991)). The latter type of fusion requires the presence of Ig regulatory sequences for expression. cDNAs encoding IgG heavy-chain constant regions can be isolated based on published sequences from cDNA libraries derived from spleen or peripheral blood lymphocytes, by hybridization or by polymerase chain reaction (PCR) techniques. The cDNAs encoding the "adhesin" and the immunoglobulin parts of the immunoadhesin are inserted in tandem into a plasmid vector that directs efficient expression in the chosen host cells.

Leucine zipper forms of these molecules are also contemplated by the invention. "Leucine zipper" is a term in the art used to refer to a leucine rich sequence that enhances, promotes, or drives dimerization or trimerization of its fusion partner (e.g., the sequence or molecule to which the leucine zipper is fused or linked to). Various leucine zipper polypeptides have been described in the art. See, e.g., Landschulz et al., *Science*, 240:1759 (1988); US Patent 5,716,805; WO 94/10308; Hoppe et al., *FEBS Letters*, 344:1991 (1994); Maniatis et al., *Nature*, 341:24 (1989). Those skilled in the art will appreciate that a leucine zipper sequence may be fused at either the 5' or 3' end of the polypeptide of this invention.

The polypeptides of the present invention can also be modified in a way to form chimeric molecules by fusing the polypeptide to another, heterologous polypeptide or amino acid sequence. According to some embodiments, such heterologous polypeptide or amino acid sequence is one which acts to oligomerize the chimeric molecule. In some embodiments, such a chimeric molecule comprises a fusion of the polypeptide with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the polypeptide. The presence of such epitope-tagged forms of the polypeptide can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the polypeptide to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., *Mol. Cell. Biol.*, 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., *Molecular and Cellular Biology*, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., *Protein Engineering*, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide

- 5 [Hopp et al., *BioTechnology*, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., *Science*, 255:192-194 (1992)]; an "-tubulin epitope peptide [Skinner et al., *J. Biol. Chem.*, 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., *Proc. Natl. Acad. Sci. USA*, 87:6393-6397 (1990)].

Construction of Peptide-Polymer Conjugates

- 10 In some embodiments the strategy for the conjugation of a polymer, (e.g. PEGylation) of synthetic peptides consists of combining, through forming a conjugate linkage in solution, a peptide and a PEG moiety, each bearing a special functionality that is mutually reactive toward the other. The peptides can be easily prepared with conventional solid phase synthesis. The peptides are "preactivated" with an appropriate functional group at a specific site. The precursors are purified and fully characterized prior to reacting with the PEG
- 15 moiety. Ligation of the peptide with PEG usually takes place in aqueous phase and can be easily monitored by reverse phase analytical HPLC. The PEGylated peptides can be easily purified by preparative HPLC and characterized by analytical HPLC, amino acid analysis and laser desorption mass spectrometry.

a. Peptide reactive sites

- In some embodiments, a peptide is covalently bonded via one or more of the amino acid residues of the peptide to a terminal reactive group on the polymer, depending mainly on the reaction conditions, the molecular weight of the polymer, etc. The polymer with the reactive group(s) is designated herein as activated polymer. The reactive group selectively reacts with free amino or other reactive groups on the peptide. Potential reactive sites include: N-terminal amino group, epsilon amino groups on lysine residues, as well as other amino, imino, carboxyl, sulfhydryl, hydroxyl, and other hydrophilic groups. It will be understood, however, that the type and
- 25 amount of the reactive group chosen, as well as the type of polymer employed, to obtain optimum results, will depend on the particular peptide employed to avoid having the reactive group react with too many particularly active groups on the peptide. In some embodiments, a reactive residue, (e.g., lysine (K), a modified, non-natural amino acid, or other small molecule) may be substituted at a position suitable for conjugation.

- In some embodiments, the peptide comprises the sequence of Formula I, Formula II, Formula III,
- 30 ECFDLLVRAWVPCSVLK (SEQ ID NO 5), ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCHMLR (SEQ ID NO 8), ECFDLLVRHWVACGLLR (SEQ ID NO 9), or sequences listed in FIG. 32 have a terminal reactive group. In some embodiments, the peptide comprises at least one and more preferably, more than one of a polypeptide comprising a sequence of Formula I, Formula II, Formula III, ECFDLLVRAWVPCSVLK (SEQ ID NO 5),
- 35 ECFDLLVRHWVPCGLLR (SEQ ID NO 6), ECFDLLVRRWVPCMLG (SEQ ID NO 7), ECFDLLVRSWVPCHMLR (SEQ ID NO 8), ECFDLLVRHWVACGLLR (SEQ ID NO 9), or sequences listed in FIG.32. The polypeptides that are linked together can have the same sequence or have different sequences and a terminal reactive group. In some embodiments, these polypeptides can be joined to one another, optionally, through the use of a linker.

- 40 While conjugation may occur at any reactive amino acid on the polypeptide, in some embodiments, the reactive amino acid is lysine, which is linked to the reactive group of the activated polymer through its free epsilon-amino group, or glutamic or aspartic acid, which is linked to the polymer through an amide bond. In some embodiments, the reactive amino acids of the peptide are not cysteine residues at positions X₂ and X₁₂.

5 The degree of polymer conjugation with each peptide will vary depending upon the number of reactive sites on the peptide, the molecular weight, hydrophilicity and other characteristics of the polymer, and the particular peptide derivatization sites chosen. In some embodiments, the conjugate has a final molar ratio of 1 to 10 polymer molecules per peptide molecule, but greater numbers of polymer molecules attached to the peptides of the invention are also contemplated. In some embodiments, each conjugate contains one polymer molecule.

10 The desired amount of derivatization is easily achieved by using an experimental matrix in which the time, temperature and other reaction conditions are varied to change the degree of substitution, after which the level of polymer substitution of the conjugates is determined by size exclusion chromatography or other means known in the art.

b. Activated polymers

15 In some embodiments, the polymer contains only a single group which is reactive. This helps to avoid cross-linking of protein molecules. However, it is within the scope herein to maximize reaction conditions to reduce cross-linking, or to purify the reaction products through gel filtration or ion exchange chromatography to recover substantially homogenous derivatives. In other embodiments, the polymer contains two or more reactive groups for the purpose of linking multiple peptides to the polymer backbone. Again, gel filtration or ion

20 exchange chromatography can be used to recover the desired derivative in substantially homogeneous form. In some embodiments, the polymer is covalently bonded directly to the peptide without the use of a multifunctional (ordinarily bifunctional) crosslinking agent. In some embodiments, there is a 1:1 molar ratio of PEG chain to peptide.

 The covalent modification reaction may take place by any appropriate method generally used for

25 reacting biologically active materials with inert polymers, preferably at about pH 5-9, more preferably 7-9 if the reactive groups on the peptide are lysine groups. Generally, the process involves preparing an activated polymer (the polymer typically having at least one terminal hydroxyl group to be activated), preparing an active substrate from this polymer, and thereafter reacting the peptide with the active substrate to produce the peptide suitable for formulation. The above modification reaction can be performed by several methods, which may involve one

30 or more steps. Examples of modifying agents that can be used to produce the activated polymer in a one-step reaction include cyanuric acid chloride (2,4,6-trichloro-S-triazine) and cyanuric acid fluoride.

 In some embodiments, the modification reaction takes place in two steps wherein the polymer is reacted first with an acid anhydride such as succinic or glutaric anhydride to form a carboxylic acid, and the carboxylic acid is then reacted with a compound capable of reacting with the carboxylic acid to form an

35 activated polymer with a reactive ester group that is capable of reacting with the peptide. Examples of such compounds include N-hydroxysuccinimide, 4-hydroxy-3-nitrobenzene sulfonic acid, and the like, and preferably N-hydroxysuccinimide or 4-hydroxy-3-nitrobenzene sulfonic acid is used. For example, monomethyl substituted PEG may be reacted at elevated temperatures, preferably about 100-110°C for four hours, with glutaric anhydride. The monomethyl PEG-glutaric acid thus produced is then reacted with N-hydroxysuccinimide in the

40 presence of a carbodiimide reagent such as dicyclohexyl or isopropyl carbodiimide to produce the activated polymer, methoxypolyethylene glycolyl-N-succinimidyl glutarate, which can then be reacted with the GH. This method is described in detail in Abuchowski et al., Cancer Biochem. Biophys., 7: 175-186 (1984). In another example, the monomethyl substituted PEG may be reacted with glutaric anhydride followed by reaction with 4-hydroxy-3-nitrobenzene sulfonic acid (HNSA) in the presence of dicyclohexyl carbodiimide to produce the

5 activated polymer. HNSA is described by Bhatnagar et al., *Peptides: Synthesis-Structure-Function. Proceedings of the Seventh American Peptide Symposium*, Rich et al. (eds.) (Pierce Chemical Co., Rockford Ill., 1981), p. 97-100, and in Nitecki et al., *High-Technology Route to Virus Vaccines* (American Society for Microbiology: 1986) entitled "Novel Agent for Coupling Synthetic Peptides to Carriers and Its Applications."

In some embodiments, covalent binding to amino groups is accomplished by known chemistries based
 10 upon cyanuric chloride, carbonyl diimidazole, aldehyde reactive groups (PEG alkoxide plus diethyl acetal of bromoacetaldehyde; PEG plus DMSO and acetic anhydride, or PEG chloride plus the phenoxide of 4-hydroxybenzaldehyde, activated succinimidyl esters, activated dithiocarbonate PEG, 2,4,5-trichlorophenylchloroformate or P-nitrophenylchloroformate activated PEG.). Carboxyl groups are derivatized by coupling PEG-amine using carbodiimide. Sulfhydryl groups are derivatized by coupling to maleimido-
 15 substituted PEG (e.g. alkoxy-PEG amine plus sulfosuccinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate) as described in WO 97/10847 published Mar. 27, 1997, or PEG-maleimide commercially available from Nektar Technologies, San Carlos, CA (formerly Shearwater Polymers, Inc.). Alternatively, free amino groups on the peptide (e.g. epsilon amino groups on lysine residues) may be coupled to N-hydroxysuccinimidyl substituted PEG (PEG-NHS available from Nektar Technologies;) or can be thiolated with 2-imino-thiolane
 20 (Traut's reagent) and then coupled to maleimide-containing derivatives of PEG as described in Pedley et al., *Br. J. Cancer*, 70: 1126-1130 (1994).

Many inert polymers, including but not limited to PEG, are suitable for use in pharmaceuticals. See, e.g., Davis et al., *Biomedical Polymers: Polymeric Materials and Pharmaceuticals for Biomedical Use*, pp.441-451 (1980). In some embodiments of the invention, a non-proteinaceous polymer is used. The nonproteinaceous
 25 polymer is typically a hydrophilic synthetic polymer, i.e., a polymer not otherwise found in nature. However, polymers which exist in nature and are produced by recombinant or in vitro methods are also useful, as are polymers which are isolated from native sources. Hydrophilic polyvinyl polymers fall within the scope of this invention, e.g. polyvinylalcohol and polyvinylpyrrolidone. Particularly useful are polyalkylene ethers such as polyethylene glycol (PEG); polyoxyalkylenes such as polyoxyethylene, polyoxypropylene, and block
 30 copolymers of polyoxyethylene and polyoxypropylene (Pluronic); polymethacrylates; carbomers; branched or unbranched polysaccharides which comprise the saccharide monomers D-mannose, D- and L-galactose, fucose, fructose, D-xylose, L-arabinose, D-glucuronic acid, sialic acid, D-galacturonic acid, D-mannuronic acid (e.g. polymannuronic acid, or alginic acid), D-glucosamine, D-galactosamine, D-glucose and neuraminic acid including homopolysaccharides and heteropolysaccharides such as lactose, amylopectin, starch, hydroxyethyl
 35 starch, amylose, dextrane sulfate, dextran, dextrans, glycogen, or the polysaccharide subunit of acid mucopolysaccharides, e.g. hyaluronic acid; polymers of sugar alcohols such as polysorbitol and polymannitol; heparin or heparon.

The polymer prior to conjugation need not be, but preferably is, water soluble, but the final conjugate is preferably water-soluble. Preferably, the conjugate exhibits a water solubility of at least about 0.01 mg/ml, and
 40 more preferably at least about 0.1 mg/ml, and still more preferably at least about 1 mg/ml. In addition, the polymer should not be highly immunogenic in the conjugate form, nor should it possess viscosity that is incompatible with intravenous infusion, injection, or inhalation if the conjugate is intended to be administered by such routes.

5 The molecular weight of the polymer can range up to about 100,000 D, and preferably is at least about 500 D, or at least about 1,000 D, or at least about 5,000 D. In some embodiments, the PEG or other polymer has a molecular weight in the range of 5000 to 20,000 D. The molecular weight chosen can depend upon the effective size of the conjugate to be achieved, the nature (e.g. structure, such as linear or branched) of the polymer, and the degree of derivatization, i.e. the number of polymer molecules per peptide, and the polymer attachment site or sites on the peptide. In some embodiments, branched PEG's may be used to induce a large increase in effective size of the peptides. PEG or other polymer conjugates may be utilized to increase half-life, increase solubility, stabilize against proteolytic attack, and reduce immunogenicity.

 Functionalized PEG polymers to modify the peptides of the invention are available from Nektar Technologies of San Carlos, CA (formerly Shearwater Polymers, Inc.). Such commercially available PEG derivatives include, but are not limited to, amino-PEG, PEG amino acid esters, PEG- N-hydroxysuccinamide chemistry (NHS), PEG-hydrazide, PEG-thiol, PEG-succinate, carboxymethylated PEG, PEG-propionic acid, PEG amino acids, PEG succinimidyl succinate, PEG succinimidyl propionate, succinimidyl ester of carboxymethylated PEG, succinimidyl carbonate of PEG, succinimidyl esters of amino acid PEGs, PEG-xycarbonylimidazole, PEG-nitrophenyl carbonate, PEG tresylate, PEG-glycidyl ether, PEG-aldehyde, PEG vinylsulfone, PEG-maleimide, PEG-orthopyridyl-disulfide, heterofunctional PEGs, PEG vinyl derivatives, PEG silanes, and PEG phospholides. The reaction conditions for coupling these PEG derivatives will vary depending on the protein, the desired degree of PEGylation, and the PEG derivative utilized. Some factors involved in the choice of PEG derivatives include: the desired point of attachment (such as lysine or cysteine R-groups), hydrolytic stability and reactivity of the derivatives, stability, toxicity and antigenicity of the linkage, suitability for analysis, etc. Specific instructions for the use of any particular derivative are available from the manufacturer.

c. Characterization of conjugates.

 The conjugates may be characterized by SDS-PAGE, gel filtration, NMR, tryptic mapping, liquid chromatography-mass spectrophotometry, and in vitro biological assays. For example, the extent of PEG conjugation may be shown by SDS-PAGE and gel filtration, and then analyzed by NMR, which has a specific resonance peak for the methylene hydrogens of PEG. The number of PEG groups on each molecule can be calculated from the NMR spectrum or mass spectrometry. Polyacrylamide gel electrophoresis in 10% SDS is appropriately run in 10 mM Tris-HCl pH 8.0, 100 mM NaCl as elution buffer. To demonstrate which residue is PEGylated, tryptic mapping can be performed. Thus, PEGylated peptides are digested with trypsin at the protein/enzyme ratio of 100 to 1 in mg basis at 37°C for 4 hours in 100 mM sodium acetate, 10 mM Tris-HCl, 1 mM calcium chloride, pH 8.3, and acidified to pH<4 to stop digestion before separating on HPLC Nucleosil C-18 (4.6 mm.times.150 mm, 5.mu., 100A). The chromatogram is compared to that of non-PEGylated starting material. Each peak can then be analyzed by mass spectrometry to verify the size of the fragment in the peak. The fragment(s) that carried PEG groups are usually not retained on the HPLC column after injection and disappear from the chromatograph. Such disappearance from the chromatograph is an indication of PEGylation on that particular fragment that should contain at least one lysine residue. PEGylated peptides may then be assayed for ability to bind to the BAFF by conventional methods.

 In some embodiments, conjugates are purified by ion-exchange chromatography, (e.g. ion exchange HPLC). The chemistry of many of the electrophilically activated PEG's results in a reduction of amino group

5 charge of the PEGylated product. Thus, high resolution ion exchange chromatography can be used to separate the free and conjugated proteins, and to resolve species with different levels of PEGylation. In fact, the resolution of different species (e.g. containing one or two PEG residues) is also possible due to the difference in the ionic properties of the unreacted amino acids. In one embodiment, species with difference levels of PEGylation are resolved according to the methods described in WO 96/34015 (International Application No. PCT/US96/05550 published Oct. 31, 1996). Heterologous species of the conjugates are purified from one another in the same fashion.

In some embodiments, PEG-N-hydroxysuccinamide (NHS) reacts with a primary amine (e.g. lysines and the N-terminus). In some embodiments, PEG-NHS reacts with a C-terminal lysine (K) of the polypeptide. In some embodiments, the lysine residue is added to the C-terminus of the 17-mer polypeptide, while in other 15 embodiments, X₁₇ is substituted with lysine. In some embodiments, the polymer reacts with the N-terminus. In a preferred embodiment, the conjugate is generated by utilizing the derivatization and purification methods described in the Examples below.

In one aspect, the invention provides any of the above-described conjugates formed by its component parts, i.e. one or more peptide(s) covalently attached to one or more polymer molecule(s), without any 20 extraneous matter in the covalent molecular structure of the conjugate.

Production of antibodies

The methods and articles of manufacture of the present invention use, or incorporate, an antibody which binds to CD20. Accordingly, methods for generating such antibodies will be described here and in the 25 Examples.

The CD20 to be used for production of, or screening for, antibodies may be, e.g., a soluble form of the antigen or a portion thereof, containing the desired epitope. The sequence of human CD20 is known, see Figure 10 and Figure 11. Cloning and the sequences for human CD20 are described in at least the following references: Stamenkovic I., Seed B., J. Exp. Med. 167, 1975-1980, 1988. Analysis of two cDNA clones encoding the B lymphocyte antigen CD20 (B1, Bp35), a type III integral membrane protein."; Tedder T.F., Streuli M., Schlossman S.F., Saito H., Proc. Natl. Acad. Sci. U.S.A. 85, 208-212, 1988. "Isolation and structure of a cDNA encoding the B1 (CD20) cell-surface antigen of human B lymphocytes"; Tedder T.F., Klejman G., Schlossman S.F., Saito H., J. Immunol. 142, 2560-2568, 1989. "Structure of the gene encoding the human B lymphocyte differentiation antigen CD20 (B1)"; Einfeld D.A., Brown J.P., Valentine M.A., Clark E.A., 35 Ledbetter J.A., EMBO J. 7, 711-717, 1988.: "Molecular cloning of the human B cell CD20 receptor predicts a hydrophobic protein with multiple transmembrane domains. "Peptide fragments of the extracellular domain (ECD) can be used as immunogens. Based on these known sequences and domain delineations, one of skill in the art can express the CD20 polypeptide and fragments thereof for use to produce antibodies.

To generate antibodies to human CD20, the extracellular domain amino acid residues 142-188 and 40 peptide fragments of 6 of greater residues in length can be used as immunogens to raise antibodies in rodents including mice, hamsters, and rats, in rabbit, goat, or other suitable animal. Soluble CD20 polypeptide or immunogenic fragments thereof can be expressed in suitable host cells such as bacteria or eukaryotic cells. In one embodiment, human and murine detergent-solubilized full-length CD20 are produced in E. coli (see Examples below) and used to immunize and screen for hybridomas producing anti-CD20 antibodies.

5 Alternatively, or additionally, B cells or cell lines expressing CD20 at their cell surface can be used to generate, and/or screen for, antibodies. One such cell line is the human lymphoblastoid cell line SB (ATCC accession no. ATCC CCL 120, from ATCC, Rockville, Md). The antibodies are generated to human CD20 for treatment of humans. Other forms of CD20 useful for generating antibodies will be apparent to those skilled in the art.

10 Phage display methodology can also be used to produce CD20 binding antibody.

The antibodies that bind CD20 may be chimeric, humanized, or human. Such antibodies and methods of generating them are described in more detail below.

(i) *Polyclonal antibodies*

15 Polyclonal antibodies are preferably raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to conjugate the relevant antigen to a protein that is immunogenic in the species to be immunized, e.g., keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are different alkyl groups.

Animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining, e.g., 100 μg or 5 μg of the protein or conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of peptide or conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later the animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate of the same antigen, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are suitably used to enhance the immune response.

(ii) *Monoclonal antibodies*

Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete antibodies.

For example, the monoclonal antibodies may be made using the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as a hamster, is immunized as hereinabove described to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)).

5 The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

10 Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Maryland USA.

15 Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

 Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as
20 radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

 The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson *et al.*, *Anal. Biochem.*, 107:220 (1980).

 After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or
25 activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal.

 The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharese, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

 DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells serve as a preferred source of such DNA.
35 Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese Hamster Ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra *et al.*, *Curr. Opinion in Immunol.*, 5:256-262 (1993) and Plückthun, *Immunol. Revs.*, 130:151-188 (1992).

40 In a further embodiment, antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in McCafferty *et al.*, *Nature*, 348:552-554 (1990). Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991) describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks *et al.*, *Bio/Technology*,

5 10:779-783 (1992)), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage libraries (Waterhouse *et al.*, *Nuc. Acids. Res.*, 21:2265-2266 (1993)). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

10 The DNA also may be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, *et al.*, *Proc. Natl Acad. Sci. USA*, 81:6851 (1984)), or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide.

15 Typically such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody, or they are substituted for the variable domains of one antigen-combining site of an antibody to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for an antigen and another antigen-combining site having specificity for a different antigen.

(iii) *Humanized antibodies*

Methods for humanizing non-human antibodies have been described in the art. Preferably, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. 20 These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, *Nature*, 321:522-525 (1986); Riechmann *et al.*, *Nature*, 332:323-327 (1988); Verhoeven *et al.*, *Science*, 239:1534-1536 (1988)), by substituting hypervariable region sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric 25 antibodies (U.S. Patent No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some hypervariable region residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized 30 antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework region (FR) for the humanized antibody (Sims *et al.*, *J. Immunol.*, 151:2296 (1993); Chothia *et al.*, *J. Mol. Biol.*, 196:901 (1987)). Another method uses a particular framework region derived from the consensus 35 sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta *et al.*, *J. Immunol.*, 151:2623 (1993)).

It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized 40 antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the

5 residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

10 (iv) *Human antibodies*

As an alternative to humanization, human antibodies can be generated. For example, it is now possible to produce transgenic animals (*e.g.*, mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, *e.g.*, Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:2551 (1993); Jakobovits *et al.*, *Nature*, 362:255-258 (1993); Bruggermann *et al.*, *Year in Immuno.*, 7:33 (1993); and US Patent Nos. 5,591,669, 5,589,369 and 5,545,807.

20 Alternatively, phage display technology (McCafferty *et al.*, *Nature* 348:552-553 (1990)) can be used to produce human antibodies and antibody fragments *in vitro*, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors. According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and displayed as functional antibody fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties. Thus, the phage mimics some of the properties of the B cell. Phage display can be performed in a variety of formats; for their review see, *e.g.*, Johnson, Kevin S. and Chiswell, David J., *Current Opinion in Structural Biology* 3:564-571 (1993). Several sources of V-gene segments can be used for phage display. Clackson *et al.*, *Nature*, 352:624-628 (1991) isolated a diverse array of anti-oxazolone antibodies from a small random combinatorial library of V genes derived from the spleens of immunized mice. A repertoire of V genes from unimmunized human donors can be constructed and antibodies to a diverse array of antigens (including self-antigens) can be isolated essentially following the techniques described by Marks *et al.*, *J. Mol. Biol.* 222:581-597 (1991), or Griffith *et al.*, *EMBO J.* 12:725-734 (1993). See, also, US Patent Nos. 5,565,332 and 5,573,905.

35 Human antibodies may also be generated by *in vitro* activated B cells (see US Patents 5,567,610 and 5,229,275).

5 (v) *Antibody fragments*

Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, *e.g.*, Morimoto *et al.*, *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992) and Brennan *et al.*, *Science*, 229:81 (1985)).

However, these fragments can now be produced directly by recombinant host cells. For example, the antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter *et al.*, *Bio/Technology* 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv).
 10 See WO 93/16185; US Patent No. 5,571,894; and US Patent No. 5,587,458. The antibody fragment may also be a "linear antibody", *e.g.*, as described in US Patent 5,641,870 for example. Such linear antibody fragments may be monospecific or bispecific.

(vi) *Bispecific antibodies*

Bispecific antibodies are antibodies that have binding specificities for at least two different epitopes.

20 Exemplary bispecific antibodies may bind to two different epitopes of the B cell surface marker. Other such antibodies may bind a first B cell marker and further bind a second B cell surface marker. Alternatively, an anti-B cell marker binding arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (*e.g.* CD2 or CD3), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the B cell.
 25 Bispecific antibodies may also be used to localize cytotoxic agents to the B cell. These antibodies possess a B cell marker-binding arm and an arm which binds the cytotoxic agent (*e.g.* saporin, anti-interferon-γ, vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten). Bispecific antibodies can be prepared as full length antibodies or antibody fragments (*e.g.* F(ab')₂ bispecific antibodies).

Methods for making bispecific antibodies are known in the art. Traditional production of full length
 30 bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein *et al.*, *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather
 35 cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker *et al.*, *EMBO J.*, 10:3655-3659 (1991).

According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2,
 40 and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the

5 mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

10 In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For
15 further details of generating bispecific antibodies see, for example, Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986). According to another approach described in US Patent No. 5,731,168, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the
20 first antibody molecule are replaced with larger side chains (*e.g.* tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (*e.g.* alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the
25 antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (US Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

30 Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, *Science*, 229: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are
35 then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be
40 chemically coupled to form bispecific antibodies. Shalaby *et al.*, *J. Exp. Med.*, 175: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

5 Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, *J. Immunol.*, 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber *et al.*, *J. Immunol.*, 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.* *J. Immunol.* 147: 60 (1991).

20 Amino acid sequence modification(s) of protein or peptide antagonists and antibodies described herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the CD20 binding antibody or antagonist. Amino acid sequence variants of the antagonist are prepared by introducing appropriate nucleotide changes into the antagonist nucleic acid, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the amino acid sequences of the antagonist. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the antagonist, such as changing the number or position of glycosylation sites.

30 A useful method for identification of certain residues or regions of the antagonist that are preferred locations for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham and Wells *Science*, 244:1081-1085 (1989). Here, a residue or group of target residues are identified (*e.g.*, charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with antigen. Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* need not be predetermined. For example, to analyze the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed antagonist variants are screened for the desired activity.

40 Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antagonist with an N-terminal methionyl residue or the antagonist fused to a cytotoxic polypeptide. Other insertional variants of the antagonist molecule include the fusion to the N- or C-terminus of the antagonist of an enzyme, or a polypeptide which increases the serum half-life of the antagonist.

- 5 Another type of variant is an amino acid substitution variant. These variants have at least one amino acid residue in the antagonist molecule replaced by different residue. The sites of greatest interest for substitutional mutagenesis of antibody antagonists include the hypervariable regions, but FR alterations are also contemplated. Conservative substitutions are shown in Table 1 under the heading of "preferred substitutions".
- 10 If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions" in Table 1, or as further described below in reference to amino acid classes, may be introduced and the products screened.

Table 1

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp, Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu
Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val; Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

15

Substantial modifications in the biological properties of the antagonist are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity

5 of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- 10 (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

Any cysteine residue not involved in maintaining the proper conformation of the antagonist also may
15 be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antagonist to improve its stability (particularly where the antagonist is an antibody fragment such as an Fv fragment).

A particularly preferred type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody. Generally, the resulting variant(s) selected for further development will
20 have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants is affinity maturation using phage display. Briefly, several hypervariable region sites (e.g. 6-7 sites) are mutated to generate all possible amino substitutions at each site. The antibody variants thus generated are displayed in a monovalent fashion from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants
25 are then screened for their biological activity (e.g. binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or in addition, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the antibody and antigen. Such contact residues and neighboring residues are candidates
30 for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

Another type of amino acid variant of the antagonist alters the original glycosylation pattern of the antagonist. By altering is meant deleting one or more carbohydrate moieties found in the antagonist, and/or
35 adding one or more glycosylation sites that are not present in the antagonist.

Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of
40 these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylglactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the antagonist is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked

5 glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antagonist (for O-linked glycosylation sites). Nucleic acid molecules encoding amino acid sequence variants of the antagonist are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-
10 directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the antagonist.

It may be desirable to modify the antagonist of the invention with respect to effector function, *e.g.* so as to enhance antigen-dependent cell-mediated cytotoxicity (ADCC) and/or complement dependent cytotoxicity (CDC) of the antagonist. This may be achieved by introducing one or more amino acid substitutions in an Fc
15 region of an antibody antagonist. Alternatively or additionally, cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, *J. Exp Med.* 176:1191-1195 (1992) and Shopes, B. *J. Immunol.* 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may
20 also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.* *Cancer Research* 53:2560-2565 (1993). Alternatively, an antibody can be engineered which has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al.* *Anti-Cancer Drug Design* 3:219-230 (1989).

To increase the serum half life of the antagonist, one may incorporate a salvage receptor binding
25 epitope into the antagonist (especially an antibody fragment) as described in US Patent 5,739,277, for example. As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (*e.g.*, IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule.

30 Assays

Peripheral B-cell concentrations are determined by a FACS method that count CD3-/CD40+ cells. The percent of CD3-CD40+ B cells of total lymphocytes in samples can be obtained by the following gating strategy. The lymphocyte population is marked on the forward scatter/ side scatter scattergram to define Region 1 (R1). Using events in R1, fluorescence intensity dot plots are displayed for CD40 and CD3 markers.
35 Fluorescently labeled isotype controls are used to determine respective cutoff points for CD40 and CD3 positivity.

FACS analysis

Half million cells are washed and resuspended in 100 µl of FACS buffer, which is phosphate buffered
40 saline with 1% BSA, containing 5 µl of staining or control antibody. All the staining antibodies, including isotype controls, are obtained from PharMingen, San Diego, CA. Human CD20 expression is assessed by staining with Rituxan® along with FITC-conjugated anti-human IgG1 secondary antibody. FACS analysis is conducted using FACScan and Cell Quest (Becton Dickinson Immunocytometry Systems, San Jose, CA). All

5 the lymphocytes are defined in the forward and side light scatterings; while all the B lymphocytes are defined with the expression of B220 on the cell surface.

B cell depletion and recovery are assessed by analyzing peripheral B cell counts and analysis of hCD20+ B cells by FACS in the spleen, lymph node and bone marrow on a daily basis for the first week after injection and thereafter on a weekly basis. Serum levels of the injected 2H7 variant antibody are monitored.

10 **Pharmaceutical Formulations**

Therapeutic formulations of the CD20-binding antibodies used in accordance with the present invention are prepared for storage by mixing an antibody having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as TWEENTM, PLURONICSTM or polyethylene glycol (PEG).

Exemplary anti-CD20 antibody formulations are described in WO98/56418, expressly incorporated herein by reference. Another formulation is a liquid multidose formulation comprising the anti-CD20 antibody at 40 mg/mL, 25 mM acetate, 150 mM trehalose, 0.9% benzyl alcohol, 0.02% polysorbate 20 at pH 5.0 that has a minimum shelf life of two years storage at 2-8°C. Another anti-CD20 formulation of interest comprises 10mg/mL antibody in 9.0 mg/mL sodium chloride, 7.35 mg/mL sodium citrate dihydrate, 0.7mg/mL polysorbate 80, and Sterile Water for Injection, pH 6.5. Yet another aqueous pharmaceutical formulation comprises 10-30 mM sodium acetate from about pH 4.8 to about pH 5.5, preferably at pH5.5, polysorbate as a surfactant in an amount of about 0.01-0.1% v/v, trehalose at an amount of about 2-10% w/v, and benzyl alcohol as a preservative (U.S. 6,171,586). Lyophilized formulations adapted for subcutaneous administration are described in WO97/04801. Such lyophilized formulations may be reconstituted with a suitable diluent to a high protein concentration and the reconstituted formulation may be administered subcutaneously to the mammal to be treated herein.

One formulation for the humanized 2H7 variants is antibody at 12-14 mg/mL in 10 mM histidine, 6% sucrose, 0.02% polysorbate 20, pH 5.8. In a specific embodiment, 2H7 variants and in particular 2H7.v16 is formulated at 20mg/mL antibody in 10mM histidine sulfate, 60mg/ml sucrose., 0.2 mg/ml polysorbate 20, and Sterile Water for Injection, at pH5.8.

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect

5 each other. For example, it may be desirable to further provide a cytotoxic agent, chemotherapeutic agent, cytokine or immunosuppressive agent (e.g. one which acts on T cells, such as cyclosporin or an antibody that binds T cells, e.g. one which binds LFA-1). The effective amount of such other agents depends on the amount of antibody present in the formulation, the type of disease or disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as described herein or
 10 about from 1 to 99% of the heretofore employed dosages.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions.
 15 Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980).

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic polymers containing the antagonist, which matrices are in the form of shaped articles, e.g. films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides
 20 (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by
 25 filtration through sterile filtration membranes.

Disease Treatment

Diseases

The CD20 binding antibodies and BAFF antagonists of the invention are useful to treat B cell
 30 malignancies and B-cell regulated autoimmune disorders.

B-cell regulated autoimmune diseases include but are not limited to arthritis (rheumatoid arthritis such as acute arthritis, chronic rheumatoid arthritis, gouty arthritis, acute gouty arthritis, chronic inflammatory arthritis, degenerative arthritis, infectious arthritis, Lyme arthritis, proliferative arthritis, psoriatic arthritis, vertebral arthritis, and juvenile-onset rheumatoid arthritis, osteoarthritis, arthritis chronica progrediente, arthritis
 35 deformans, polyarthritis chronica primaria, reactive arthritis, and ankylosing spondylitis), inflammatory hyperproliferative skin diseases, psoriasis such as plaque psoriasis, guttate psoriasis, pustular psoriasis, and psoriasis of the nails, atopy including atopic diseases such as hay fever and Job's syndrome, dermatitis including contact dermatitis, chronic contact dermatitis, allergic dermatitis, allergic contact dermatitis, dermatitis herpetiformis, and atopic dermatitis, x-linked hyper IgM syndrome, urticaria such as chronic allergic urticaria
 40 and chronic idiopathic urticaria, including chronic autoimmune urticaria, polymyositis/dermatomyositis, juvenile dermatomyositis, toxic epidermal necrolysis, scleroderma (including systemic scleroderma), sclerosis such as systemic sclerosis, multiple sclerosis (MS) such as spino-optical MS, primary progressive MS (PPMS), and relapsing remitting MS (RRMS), progressive systemic sclerosis, atherosclerosis, arteriosclerosis, sclerosis disseminata, and ataxic sclerosis, inflammatory bowel disease (IBD) (for example, Crohn's disease,

5 autoimmune-mediated gastrointestinal diseases, colitis such as ulcerative colitis, colitis ulcerosa, microscopic colitis, collagenous colitis, colitis polyposa, necrotizing enterocolitis, and transmural colitis, and autoimmune inflammatory bowel disease), pyoderma gangrenosum, erythema nodosum, primary sclerosing cholangitis, episcleritis), respiratory distress syndrome, including adult or acute respiratory distress syndrome (ARDS), meningitis, inflammation of all or part of the uvea, iritis, choroiditis, an autoimmune hematological disorder,

10 rheumatoid spondylitis, sudden hearing loss, IgE-mediated diseases such as anaphylaxis and allergic and atopic rhinitis, encephalitis such as Rasmussen's encephalitis and limbic and/or brainstem encephalitis, uveitis, such as anterior uveitis, acute anterior uveitis, granulomatous uveitis, nongranulomatous uveitis, phacoantigenic uveitis, posterior uveitis, or autoimmune uveitis, glomerulonephritis (GN) with and without nephrotic syndrome such as chronic or acute glomerulonephritis such as primary GN, immune-mediated GN, membranous GN

15 (membranous nephropathy), idiopathic membranous GN or idiopathic membranous nephropathy, membrano- or membranous proliferative GN (MPGN), including Type I and Type II, and rapidly progressive GN, allergic conditions, allergic reaction, eczema including allergic or atopic eczema, asthma such as asthma bronchiale, bronchial asthma, and auto-immune asthma, conditions involving infiltration of T cells and chronic inflammatory responses, chronic pulmonary inflammatory disease, autoimmune myocarditis, leukocyte

20 adhesion deficiency, systemic lupus erythematosus (SLE) or systemic lupus erythematoses such as cutaneous SLE, subacute cutaneous lupus erythematosus, neonatal lupus syndrome (NLE), lupus erythematosus disseminatus, lupus (including nephritis, cerebritis, pediatric, non-renal, extra-renal, discoid, alopecia), juvenile onset (Type I) diabetes mellitus, including pediatric insulin-dependent diabetes mellitus (IDDM), adult onset diabetes mellitus (Type II diabetes), autoimmune diabetes, idiopathic diabetes insipidus, immune responses

25 associated with acute and delayed hypersensitivity mediated by cytokines and T-lymphocytes, tuberculosis, sarcoidosis, granulomatosis including lymphomatoid granulomatosis, Wegener's granulomatosis, agranulocytosis, vasculitides, including vasculitis (including large vessel vasculitis (including polymyalgia rheumatica and giant cell (Takayasu's) arteritis), medium vessel vasculitis (including Kawasaki's disease and polyarteritis nodosa), microscopic polyarteritis, CNS vasculitis, necrotizing, cutaneous, or hypersensitivity

30 vasculitis, systemic necrotizing vasculitis, and ANCA-associated vasculitis, such as Churg-Strauss vasculitis or syndrome (CSS)), temporal arteritis, aplastic anemia, autoimmune aplastic anemia, Coombs positive anemia, Diamond Blackfan anemia, hemolytic anemia or immune hemolytic anemia including autoimmune hemolytic anemia (AIHA), pernicious anemia (anemia perniciosa), Addison's disease, pure red cell anemia or aplasia (PRCA), Factor VIII deficiency, hemophilia A, autoimmune neutropenia, pancytopenia, leukopenia, diseases

35 involving leukocyte diapedesis, CNS inflammatory disorders, multiple organ injury syndrome such as those secondary to septicemia, trauma or hemorrhage, antigen-antibody complex-mediated diseases, anti-glomerular basement membrane disease, anti-phospholipid antibody syndrome, allergic neuritis, Bechet's or Behcet's disease, Castleman's syndrome, Goodpasture's syndrome, Reynaud's syndrome, Sjogren's syndrome, Stevens-Johnson syndrome, pemphigoid such as pemphigoid bullous and skin pemphigoid, pemphigus (including

40 pemphigus vulgaris, pemphigus foliaceus, pemphigus mucus-membrane pemphigoid, and pemphigus erythematosus), autoimmune polyendocrinopathies, Reiter's disease or syndrome, immune complex nephritis, antibody-mediated nephritis, neuromyelitis optica, polyneuropathies, chronic neuropathy such as IgM polyneuropathies or IgM-mediated neuropathy, thrombocytopenia (as developed by myocardial infarction patients, for example), including thrombotic thrombocytopenic purpura (TTP), post-transfusion purpura (PTP),

5 heparin-induced thrombocytopenia, and autoimmune or immune-mediated thrombocytopenia such as idiopathic
 thrombocytopenic purpura (ITP) including chronic or acute ITP, autoimmune disease of the testis and ovary
 including autoimmune orchitis and oophoritis, primary hypothyroidism, hypoparathyroidism, autoimmune
 endocrine diseases including thyroiditis such as autoimmune thyroiditis, Hashimoto's disease, chronic
 thyroiditis (Hashimoto's thyroiditis), or subacute thyroiditis, autoimmune thyroid disease, idiopathic
 10 hypothyroidism, Grave's disease, polyglandular syndromes such as autoimmune polyglandular syndromes (or
 polyglandular endocrinopathy syndromes), paraneoplastic syndromes, including neurologic paraneoplastic
 syndromes such as Lambert-Eaton myasthenic syndrome or Eaton-Lambert syndrome, stiff-man or stiff-person
 syndrome, encephalomyelitis such as allergic encephalomyelitis or encephalomyelitis allergica and experimental
 allergic encephalomyelitis (EAE), myasthenia gravis such as thymoma-associated myasthenia gravis, cerebellar
 15 degeneration, neuromyotonia, opsoclonus or opsoclonus myoclonus syndrome (OMS), and sensory neuropathy,
 multifocal motor neuropathy, Sheehan's syndrome, autoimmune hepatitis, chronic hepatitis, lupoid hepatitis,
 giant cell hepatitis, chronic active hepatitis or autoimmune chronic active hepatitis, lymphoid interstitial
 pneumonitis (LIP), bronchiolitis obliterans (non-transplant) vs NSIP, Guillain-Barré syndrome, Berger's disease
 (IgA nephropathy), idiopathic IgA nephropathy, linear IgA dermatosis, primary biliary cirrhosis,
 20 pneumonocirrhosis, autoimmune enteropathy syndrome, Celiac disease, Coeliac disease, celiac sprue (gluten
 enteropathy), refractory sprue, idiopathic sprue, cryoglobulinemia, amyotrophic lateral sclerosis (ALS; Lou
 Gehrig's disease), coronary artery disease, autoimmune ear disease such as autoimmune inner ear disease
 (AIED), autoimmune hearing loss, opsoclonus myoclonus syndrome (OMS), polychondritis such as refractory
 or relapsed polychondritis, pulmonary alveolar proteinosis, amyloidosis, scleritis, a non-cancerous
 25 lymphocytosis, a primary lymphocytosis, which includes monoclonal B cell lymphocytosis (e.g., benign
 monoclonal gammopathy and monoclonal gammopathy of undetermined significance, MGUS), peripheral
 neuropathy, paraneoplastic syndrome, channelopathies such as epilepsy, migraine, arrhythmia, muscular
 disorders, deafness, blindness, periodic paralysis, and channelopathies of the CNS, autism, inflammatory
 myopathy, focal segmental glomerulosclerosis (FSGS), endocrine ophthalmopathy, uveoretinitis, chorioretinitis,
 30 autoimmune hepatological disorder, fibromyalgia, multiple endocrine failure, Schmidt's syndrome, adrenalitis,
 gastric atrophy, presenile dementia, demyelinating diseases such as autoimmune demyelinating diseases and
 chronic inflammatory demyelinating polyneuropathy, diabetic nephropathy, Dressler's syndrome, alopecia
 areata, CREST syndrome (calcinosis, Raynaud's phenomenon, esophageal dysmotility, sclerodactyly, and
 telangiectasia), male and female autoimmune infertility, mixed connective tissue disease, Chagas' disease,
 35 rheumatic fever, recurrent abortion, farmer's lung, erythema multiforme, post-cardiotomy syndrome, Cushing's
 syndrome, bird-fancier's lung, allergic granulomatous angiitis, benign lymphocytic angiitis, Alport's syndrome,
 alveolitis such as allergic alveolitis and fibrosing alveolitis, interstitial lung disease, transfusion reaction,
 leprosy, malaria, leishmaniasis, kypanosomiasis, schistosomiasis, ascariasis, aspergillosis, Sampter's syndrome,
 Caplan's syndrome, dengue, endocarditis, endomyocardial fibrosis, diffuse interstitial pulmonary fibrosis,
 40 interstitial lung fibrosis, pulmonary fibrosis, idiopathic pulmonary fibrosis, cystic fibrosis, endophthalmitis,
 erythema elevatum et diutinum, erythroblastosis fetalis, eosinophilic fasciitis, Shulman's syndrome, Felty's
 syndrome, filariasis, cyclitis such as chronic cyclitis, heterochronic cyclitis, iridocyclitis, or Fuch's cyclitis,
 Henoch-Schönlein purpura, human immunodeficiency virus (HIV) infection, echovirus infection,
 cardiomyopathy, Alzheimer's disease, parvovirus infection, rubella virus infection, post-vaccination syndromes,

- 5 congenital rubella infection, Epstein-Barr virus infection, mumps, Evan's syndrome, autoimmune gonadal failure, Sydenham's chorea, post-streptococcal nephritis, thromboangitis obliterans, thyrotoxicosis, tabes dorsalis, chorioiditis, giant cell polymyalgia, endocrine ophthalmopathy, chronic hypersensitivity pneumonitis, keratoconjunctivitis sicca, epidemic keratoconjunctivitis, idiopathic nephritic syndrome, minimal change nephropathy, benign familial and ischemia-reperfusion injury, retinal autoimmunity, joint inflammation,
- 10 bronchitis, chronic obstructive airway disease, silicosis, aphthae, aphthous stomatitis, arteriosclerotic disorders, aspermiogenesis, autoimmune hemolysis, Boeck's disease, cryoglobulinemia, Dupuytren's contracture, endophthalmitis phacoanaphylactica, enteritis allergica, erythema nodosum leprosum, idiopathic facial paralysis, chronic fatigue syndrome, febris rheumatica, Hamman-Rich's disease, sensorineural hearing loss, haemoglobinuria paroxysmalica, hypogonadism, ileitis regionalis, leucopenia, mononucleosis infectiosa,
- 15 traverse myelitis, primary idiopathic myxedema, nephrosis, ophthalmia sympathica, orchitis granulomatosa, pancreatitis, polyradiculitis acuta, pyoderma gangrenosum, Quervain's thyroiditis, acquired splenic atrophy, infertility due to antispermatozoan antibodies, non-malignant thymoma, vitiligo, SCID and Epstein-Barr virus-associated diseases, acquired immune deficiency syndrome (AIDS), parasitic diseases such as Leishmania, toxic-shock syndrome, food poisoning, conditions involving infiltration of T cells, leukocyte-adhesion
- 20 deficiency, immune responses associated with acute and delayed hypersensitivity mediated by cytokines and T-lymphocytes, diseases involving leukocyte diapedesis, multiple organ injury syndrome, antigen-antibody complex-mediated diseases, antiglomerular basement membrane disease, allergic neuritis, autoimmune polyendocrinopathies, oophoritis, primary myxedema, autoimmune atrophic gastritis, sympathetic ophthalmia, rheumatic diseases, mixed connective tissue disease, nephrotic syndrome, insulinitis, polyendocrine failure,
- 25 peripheral neuropathy, autoimmune polyglandular syndrome type I, adult-onset idiopathic hypoparathyroidism (AOIH), alopecia totalis, dilated cardiomyopathy, epidermolysis bullosa acquisita (EBA), hemochromatosis, myocarditis, nephrotic syndrome, primary sclerosing cholangitis, purulent or nonpurulent sinusitis, acute or chronic sinusitis, ethmoid, frontal, maxillary, or sphenoid sinusitis, an eosinophil-related disorder such as eosinophilia, pulmonary infiltration eosinophilia, eosinophilia-myalgia syndrome, Löffler's syndrome, chronic
- 30 eosinophilic pneumonia, tropical pulmonary eosinophilia, bronchopneumonic aspergillosis, aspergilloma, or granulomas containing eosinophils, anaphylaxis, seronegative spondyloarthritides, polyendocrine autoimmune disease, sclerosing cholangitis, sclera, episclera, chronic mucocutaneous candidiasis, Bruton's syndrome, transient hypogammaglobulinemia of infancy, Wiskott-Aldrich syndrome, ataxia telangiectasia, autoimmune disorders associated with collagen disease, rheumatism, neurological disease, ischemic re-perfusion disorder,
- 35 reduction in blood pressure response, vascular dysfunction, angiectasis, tissue injury, cardiovascular ischemia, hyperalgesia, cerebral ischemia, and disease accompanying vascularization, allergic hypersensitivity disorders, glomerulonephritides, reperfusion injury, reperfusion injury of myocardial or other tissues, dermatoses with acute inflammatory components, acute purulent meningitis or other central nervous system inflammatory disorders, ocular and orbital inflammatory disorders, granulocyte transfusion-associated syndromes, cytokine-
- 40 induced toxicity, acute serious inflammation, chronic intractable inflammation, pyelitis, pneumonocirrhosis, diabetic retinopathy, diabetic large-artery disorder, endarterial hyperplasia, peptic ulcer, vulvitis, and endometriosis.

The B cell neoplasms include CD20-positive Hodgkin's disease including lymphocyte predominant Hodgkin's disease (LPHD); non-Hodgkin's lymphoma (NHL); follicular center cell (FCC) lymphomas; acute

- 5 lymphocytic leukemia (ALL); chronic lymphocytic leukemia (CLL); Hairy cell leukemia. The non-Hodgkins lymphoma include low grade/follicular non-Hodgkin's lymphoma (NHL), small lymphocytic (SL) NHL, intermediate grade/follicular NHL, intermediate grade diffuse NHL, high grade immunoblastic NHL, high grade lymphoblastic NHL, high grade small non-cleaved cell NHL, bulky disease NHL, plasmacytoid lymphocytic lymphoma, mantle cell lymphoma, AIDS- related lymphoma and Waldenstrom's macroglobulinemia.
- 10 Treatment of relapses of these cancers are also contemplated. LPHD is a type of Hodgkin's disease that tends to relapse frequently despite radiation or chemotherapy treatment and is characterized by CD20-positive malignant cells. CLL is one of four major types of leukemia. A cancer of mature B-cells called lymphocytes, CLL is manifested by progressive accumulation of cells in blood, bone marrow and lymphatic tissues. Indolent lymphoma is a slow-growing, incurable disease in which the average patient survives between six and 10 years
- 15 following numerous periods of remission and relapse.

In specific embodiments, the BAFF antagonists and CD20 binding antibodies are used to treat non-Hodgkin's lymphoma (NHL), lymphocyte predominant Hodgkin's disease (LPHD), chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL) which is a type of non-Hodgkin's lymphoma (NHL), diffuse large B cell lymphoma, rheumatoid arthritis and juvenile rheumatoid arthritis, systemic lupus

20 erythematosus (SLE) including lupus nephritis, Wegener's disease, inflammatory bowel disease, idiopathic thrombocytopenic purpura (ITP), thrombotic thrombocytopenic purpura (TTP), autoimmune thrombocytopenia, multiple sclerosis, psoriasis, IgA nephropathy, IgM polyneuropathies, myasthenia gravis, vasculitis, diabetes mellitus, Reynaud's syndrome, Sjorgen's syndrome and glomerulonephritis.

The desired level of B cell depletion will depend on the disease. For the treatment of a CD20 positive

25 cancer, it may be desirable to maximize the depletion of the B cells which are the target of the anti-CD20 antibodies of the invention. Thus, for the treatment of a CD20 positive B cell neoplasm, it is desirable that the B cell depletion be sufficient to at least prevent progression of the disease which can be assessed by the physician of skill in the art, e.g., by monitoring tumor growth (size), proliferation of the cancerous cell type, metastasis, other signs and symptoms of the particular cancer. Preferably, the B cell depletion is sufficient to prevent

30 progression of disease for at least 2 months, more preferably 3 months, even more preferably 4 months, more preferably 5 months, even more preferably 6 or more months. In even more preferred embodiments, the B cell depletion is sufficient to increase the time in remission by at least 6 months, more preferably 9 months, more preferably one year, more preferably 2 years, more preferably 3 years, even more preferably 5 or more years. In a most preferred embodiment, the B cell depletion is sufficient to cure the disease. In preferred embodiments,

35 the B cell depletion in a cancer patient is at least about 75% and more preferably, 80%, 85%, 90%, 95%, 99% and even 100% of the baseline level before treatment.

For treatment of an autoimmune disease, it may be desirable to modulate the extent of B cell depletion depending on the disease and/or the severity of the condition in the individual patient, by adjusting the dosage of CD20 binding antibody. Thus, B cell depletion can but does not have to be complete. Or, total B cell depletion

40 may be desired in initial treatment but in subsequent treatments, the dosage may be adjusted to achieve only partial depletion. In one embodiment, the B cell depletion is at least 20%, i.e., 80% or less of CD20 positive B cells remain as compared to the baseline level before treatment. In other embodiments, B cell depletion is 25%, 30%, 40%, 50%, 60%, 70% or greater. Preferably, the B cell depletion is sufficient to halt progression of the

5 disease, more preferably to alleviate the signs and symptoms of the particular disease under treatment, even more preferably to cure the disease.

Publications concerning therapy with Rituximab include: Perrotta and Abuel "Response of chronic relapsing ITP of 10 years duration to Rituximab" Abstract # 3360 *Blood* 10(1)(part 1-2): p. 88B (1998); Stasi *et al.* "Rituximab chimeric anti-CD20 monoclonal antibody treatment for adults with chronic idiopathic thrombocytopenic purpura" *Blood* 98(4):952-957 (2001); Matthews, R. "Medical Heretics" *New Scientist* (7 April, 2001); Leandro *et al.* "Clinical outcome in 22 patients with rheumatoid arthritis treated with B lymphocyte depletion" *Ann Rheum Dis* 61:833-888 (2002); Leandro *et al.* "Lymphocyte depletion in rheumatoid arthritis: early evidence for safety, efficacy and dose response. *Arthritis and Rheumatism* 44(9): S370 (2001); Leandro *et al.* "An open study of B lymphocyte depletion in systemic lupus erythematosus", *Arthritis & Rheumatism* 46(1):2673-2677 (2002); Edwards and Cambridge "Sustained improvement in rheumatoid arthritis following a protocol designed to deplete B lymphocytes" *Rheumatology* 40:205-211 (2001); Edwards *et al.* "B-lymphocyte depletion therapy in rheumatoid arthritis and other autoimmune disorders" *Biochem. Soc. Trans.* 30(4):824-828 (2002); Edwards *et al.* "Efficacy and safety of Rituximab, a B-cell targeted chimeric monoclonal antibody: A randomized, placebo controlled trial in patients with rheumatoid arthritis. *Arthritis and Rheumatism* 46(9): S197 (2002); Levine and Pestronk "IgM antibody-related polyneuropathies: B-cell depletion chemotherapy using Rituximab" *Neurology* 52: 1701-1704 (1999); DeVita *et al.* "Efficacy of selective B cell blockade in the treatment of rheumatoid arthritis" *Arthritis & Rheum* 46:2029-2033 (2002); Higashida *et al.* "Treatment of DMARD-Refractory rheumatoid arthritis with rituximab." Presented at the *Annual Scientific Meeting of the American College of Rheumatology*; Oct 24-29; New Orleans, LA 2002; Tusciano, J. "Successful treatment of Infliximab-refractory rheumatoid arthritis with rituximab" Presented at the *Annual Scientific Meeting of the American College of Rheumatology*; Oct 24-29; New Orleans, LA 2002.

For therapeutic applications, the anti-CD20 antibody and BAFF antagonist compositions of the invention can be used in combination therapy with, e.g., chemotherapeutic agents, hormones, antiangiogens, radiolabelled compounds, or with surgery, cryotherapy, and/or radiotherapy. The preceding treatment methods can be administered in conjunction with other forms of conventional therapy, either consecutively with, pre- or post-conventional therapy. The anti-CD20 antibody and BAFF antagonist will be administered with a therapeutically effective dose of the chemotherapeutic agent. In another embodiment, the anti-CD20 antibody and BAFF antagonist are administered in conjunction with chemotherapy to enhance the activity and efficacy of the chemotherapeutic agent. The Physicians' Desk Reference (PDR) discloses dosages of chemotherapeutic agents that have been used in the treatment of various cancers. The dosing regimen and dosages of these aforementioned chemotherapeutic drugs that are therapeutically effective will depend on the particular cancer being treated, the extent of the disease and other factors familiar to the physician of skill in the art and can be determined by the physician.

A patient is alleviated or successfully treated of a B cell neoplasm or a B cell regulated autoimmune diseases by the present methods of the invention if there is a measurable improvement in the symptoms or other applicable criteria after administration of the compositions of the invention compared to before treatment. The effect of treatment may be apparent within 3-10 weeks after administration of the compositions of the invention. The applicable criteria for each disease will be well known to the physician of skill in the appropriate art. For example, the physician can monitor the treated patient for clinical, or serologic evidence of disease such as

5 serologic markers of disease, complete blood count including B cell count, and serum immunoglobulin levels. Serum levels of IgG and IgM are reduced in BR3-Fc treated mice. It is expected that human patients responding to BR3-Fc or anti-CD20 antibody treatment or both would likewise show a reduction in serum IgG and IgM levels. The patient may show observable and/or measurable reduction in or absence of one or more of the following: reduction in the number of cancer cells or absence of the cancer cells; reduction in the tumor size; inhibition (*i.e.*, slow to some extent and preferably stop) of cancer cell infiltration into organs; inhibition (*i.e.*, slow to some extent and preferably stop) of tumor metastasis; inhibition, to some extent, of tumor growth; and/or relief to some extent, one or more of the symptoms associated with the specific cancer; reduced morbidity and mortality, and improvement in quality of life issues. Preferably, after administration of the compositions of the invention, the improvement is at least 20 % over the baseline for a particular symptom or criterion taken before treatment by the methods of the invention, more preferably, 25-30%, even more preferably 30-35%, most preferably 40% and above.

The parameters for assessing efficacy or success of treatment of the neoplasm will be known to the physician of skill in the appropriate disease. Generally, the physician of skill will look for reduction in the signs and symptoms of the specific disease. Parameters can include median time to disease progression, time in remission and stable disease. For B cell neoplasms, measurable criteria may include, *e.g.*, time to disease progression, an increase in duration of overall and/or progression-free survival. In the case of leukemia, a bone marrow biopsy can be conducted to determine the degree of remission. Complete remission can be defined as the leukemia cells making up less than 5 percent of all cells found in a patient's bone marrow 30 days following treatment.

25 The following references describe lymphomas and CLL, their diagnoses, treatment and standard medical procedures for measuring treatment efficacy. Canellos GP, Lister, TA, Sklar JL: *The Lymphomas*. W.B. Saunders Company, Philadelphia, 1998; van Besien K and Cabanillas, F: Clinical Manifestations, Staging and Treatment of Non-Hodgkin's Lymphoma, Chap. 70, pp 1293-1338, in: *Hematology , Basic Principles and Practice*, 3rd ed. Hoffman et al. (editors). Churchill Livingstone, Philadelphia, 2000; and Rai, K and Patel, D: Chronic Lymphocytic Leukemia, Chap. 72, pp 1350-1362, in: *Hematology , Basic Principles and Practice*, 3rd ed. Hoffman et al. (editors). Churchill Livingstone, Philadelphia, 2000.

The parameters for assessing efficacy or success of treatment of an autoimmune or autoimmune related disease will be known to the physician of skill in the appropriate disease. Generally, the physician of skill will look for reduction in the signs and symptoms of the specific disease. The following are by way of examples.

35 Rheumatoid arthritis (RA) is an autoimmune disorder of unknown etiology. Most RA patients suffer a chronic course of disease that, even with therapy, may result in progressive joint destruction, deformity, disability and even premature death. The goals of RA therapy are to prevent or control joint damage, prevent loss of function and decrease pain. Initial therapy of RA usually involves administration of one or more of the following drugs: nonsteroidal anti-inflammatory drugs (NSAIDs), glucocorticoid (via joint injection), and low-dose prednisone. See "Guidelines for the management of rheumatoid arthritis" *Arthritis & Rheumatism* 46(2): 328-346 (February, 2002). The majority of patients with newly diagnosed RA are started with disease-modifying antirheumatic drug (DMARD) therapy within 3 months of diagnosis. DMARDs commonly used in RA are hydroxycloquine, sulfasalazine, methotrexate, leflunomide, etanercept, infliximab (plus oral and

5 subcutaneous methotrexate), azathioprine, D-penicillamine, Gold (oral), Gold (intramuscular), minocycline, cyclosporine, Staphylococcal protein A immunoadsorption.

Because the body produces tumor necrosis factor alpha (TNF α) during RA, TNF α inhibitors have used for therapy of that disease. Etanercept (ENBREL®) is an injectable drug approved in the US for therapy of active RA. Etanercept binds to TNF α and serves to remove most TNF α from joints and blood, thereby
10 preventing TNF α from promoting inflammation and other symptoms of rheumatoid arthritis. Etanercept is an "immunoadhesin" fusion protein consisting of the extracellular ligand binding portion of the human 75 kD (p75) tumor necrosis factor receptor (TNFR) linked to the Fc portion of a human IgG1. Infliximab, sold under the trade name REMICADE®, is an immune-suppressing drug prescribed to treat RA and Crohn's disease. Infliximab is a chimeric monoclonal antibody that binds to TNF α and reduces inflammation in the body by
15 targeting and binding to TNF α which produces inflammation.

Adalimumab (HUMIRA™, Abbott Laboratories), previously known as D2E7, is a human monoclonal antibody that binds to TNF α and is approved for reducing the signs and symptoms and inhibiting the progression of structural damage in adults with moderately to severely active RA who have had insufficient response to one or more traditional disease modifying DMARDs.

20 Treatment of rheumatoid arthritis by administering an anti-CD20 antibody and a BAFF antagonist can be preformed in conjunction with therapy with one or more of the aforementioned drugs for RA.

For rheumatoid arthritis, for example, measurements for progress in treatment may include the number of swollen and tender joints and the length of morning stiffness. Patients may be examined for how much the joint in the hands and feet have eroded by using X-rays and a scoring system known as the Sharp score.

25 Another scoring system is based on the American College of Rheumatology criteria for assessing response to therapies.

One method of evaluating treatment efficacy in RA is based on American College of Rheumatology (ACR) criteria, which measures the percentage of improvement in tender and swollen joints, among other things. The RA patient can be scored at for example, ACR 20 (20 percent improvement) compared with no
30 antibody treatment (e.g., baseline before treatment) or treatment with placebo. Other ways of evaluating the efficacy of antibody treatment include X-ray scoring such as the Sharp X-ray score used to score structural damage such as bone erosion and joint space narrowing. Patients can also be evaluated for the prevention of or improvement in disability based on Health Assessment Questionnaire [HAQ] score, AIMS score, SF-36 at time periods during or after treatment. The ACR 20 criteria may include 20% improvement in both tender (painful)
35 joint count and swollen joint count plus a 20% improvement in at least 3 of 5 additional measures:

1. patient's pain assessment by visual analog scale (VAS),
2. patient's global assessment of disease activity (VAS),
3. physician's global assessment of disease activity (VAS),
4. patient's self-assessed disability measured by the Health Assessment Questionnaire, and
- 40 5. acute phase reactants, CRP or ESR.

The ACR 50 and 70 are defined analogously. Preferably, the patient is administered an amount of a CD20 binding antibody of the invention effective to achieve at least a score of ACR 20, preferably at least ACR 30, more preferably at least ACR50, even more preferably at least ACR70, most preferably at least ACR 75 and
45 higher.

5 Psoriatic arthritis has unique and distinct radiographic features. For psoriatic arthritis, joint erosion and joint space narrowing can be evaluated by the Sharp score as well. The humanized CD20 binding antibodies disclosed herein can be used to prevent the joint damage as well as reduce disease signs and symptoms of the disorder.

10 Yet another aspect of the invention is a method of treating Lupus or SLE by administering to the patient suffering from SLE, a therapeutically effective amount of a humanized CD20 binding antibody of the invention. SLEDAI scores provide a numerical quantitation of disease activity. The SLEDAI is a weighted index of 24 clinical and laboratory parameters known to correlate with disease activity, with a numerical range of 0-103. see Bryan Gescuk & John Davis, "Novel therapeutic agent for systemic lupus erythematosus" in Current Opinion in Rheumatology 2002, 14:515-521. Antibodies to double-stranded DNA are believed to cause
15 renal flares and other manifestations of lupus. Patients undergoing antibody treatment can be monitored for time to renal flare, which is defined as a significant, reproducible increase in serum creatinine, urine protein or blood in the urine. Alternatively or in addition, patients can be monitored for levels of antinuclear antibodies and antibodies to double-stranded DNA. Treatments for SLE include high-dose corticosteroids and/or cyclophosphamide (HDCC).

20 Spondyloarthropathies are a group of disorders of the joints, including ankylosing spondylitis, psoriatic arthritis and Crohn's disease. Treatment success can be determined by validated patient and physician global assessment measuring tools.

For systemic lupus erythematosus, patients can be monitored for levels of antinuclear antibodies and antibodies to double-stranded DNA.

25 Various medications are used to treat psoriasis; treatment differs directly in relation to disease severity. Patients with a more mild form of psoriasis typically utilize topical treatments, such as topical steroids, anthralin, calcipotriene, clobetasol, and tazarotene, to manage the disease while patients with moderate and severe psoriasis are more likely to employ systemic (methotrexate, retinoids, cyclosporine, PUVA and UVB) therapies. Tars are also used. These therapies have a combination of safety concerns, time consuming regimens,
30 or inconvenient processes of treatment. Furthermore, some require expensive equipment and dedicated space in the office setting. Systemic medications can produce serious side effects, including hypertension, hyperlipidemia, bone marrow suppression, liver disease, kidney disease and gastrointestinal upset. Also, the use of phototherapy can increase the incidence of skin cancers. In addition to the inconvenience and discomfort associated with the use of topical therapies, phototherapy and systemic treatments require cycling patients on
35 and off therapy and monitoring lifetime exposure due to their side effects.

Treatment efficacy for psoriasis is assessed by monitoring changes in clinical signs and symptoms of the disease including Physician's Global Assessment (PGA) changes and Psoriasis Area and Severity Index (PASI) scores, Psoriasis Symptom Assessment (PSA), compared with the baseline condition. The patient can be measured periodically throughout treatment on the Visual analog scale used to indicate the degree of itching
40 experienced at specific time points.

Dosing

Depending on the indication to be treated and factors relevant to the dosing that a physician of skill in the field would be familiar with, the BAFF antagonists and CD20 binding antibodies of the invention will be

5 administered at a dosage that is efficacious for the treatment of that indication while minimizing toxicity and side effects. For the treatment of patients suffering from B-cell neoplasm such as non-Hodgkins lymphoma, in a specific embodiment, the anti-CD20 antibodies of the invention will be administered to a human patient at a dosage range of 1mg/kg to 20mg/kg body weight, preferably at 2.5mg/kg to 10mg/kg. In a preferred
10 embodiment, the anti-CD20 antibody is administered at a dosage of 10mg/kg or 375mg/m². For treating NHL, one dosing regimen would be to administer 375mg/m² of anti-CD20 antibody every other week for 2-4 doses, or one dose of the antibody composition in the first week of treatment, followed by a 2 week interval, then a second dose of the same amount of antibody is administered. Generally, NHL patients receive such treatment once during a year but upon recurrence of the lymphoma, such treatment can be repeated. In the treatment of
15 NHL, the anti-CD20 antibody plus BAFF antagonist therapy can be combined with chemotherapy such as with CHOP. In another embodiment, for the treatment of B cell neoplasms such as CLL or SLL, patients may receive four weekly doses of Rituxan at 375 mg/m² after or before administration with BR3-Fc. with relapsed CLL. For CLL, treatment with the anti-CD20 antibody and BAFF antagonists can be combined with chemotherapy, for example, with fludarabine and cytoxan.

For treating rheumatoid arthritis, in one embodiment, RituxanTM which is a chimeric antibody is
20 administered at 500mg per dose every other week for a total of 2 doses. A humanized anti-CD20 antibody, e.g., hu2H7v.16 or any other variant of hu 2H7 as disclosed herein, can be administered at less than 500mg per dose such as at between about 200-500mg per dose, between about 250mg-450mg, or 300-400mg per dose, for 2-4 doses every other week or every 3rd week.

BR3-Fc can be administered at a dosage range of 0.5mg/kg to 10mg/kg, preferably 1mg/kg to 5mg/kg,
25 more preferably, 1.5mg/kg to 2.5mg/kg. In one embodiment, BR3-Fc is administered at 5mg/kg every other day from day 1 to day 12 of treatment. Also contemplated is dosing at about 2-5mg/kg every 2-3 days for a total of 2-5 doses.

The treatment methods of the invention comprises a combination of concurrently and sequentially
30 administering the anti-CD20 antibody and the BAFF antagonist (both referred to herein as the drugs). In sequential administration, the drugs can be administered in either order, i.e., BAFF antagonist first followed by anti-CD20 antibody. The patient is treated with one drug and monitored for efficacy before treatment with the one drug. For example, if the BAFF antagonist produces a partial response, treatment can be followed with the anti-CD20 antibody to achieve a full response, and vice versa. The BR3-Fc which is an immunoadhesin, has a shorter half compared to a full length anti-CD20 antibody. For the treatment of autoimmune diseases such as
35 rheumatoid arthritis, if the anti-CD20 antibody is Rituxan and the BAFF antagonist is BR3-Fc, in a preferred embodiment, the patient in need thereof receives BR3-Fc prior to treatment with Rituxan. Alternatively, the patient can be initially administered both drugs and subsequent dosing can be with only one or the other drug.

To condition the patient to tolerate the drugs and/or to reduce the occurrence of adverse effects such as
40 infusion-related symptoms which arise from the initial and subsequent administrations of the therapeutic compound, the mammal in need thereof can be administered a first or initial conditioning dose of one or both drugs and then administered at least a second therapeutically effective dose of one or both drugs wherein the second and any subsequent doses are higher than the first dose. The first dose serves to condition the mammal to tolerate the higher second therapeutic dose. In this way, the mammal is able to tolerate higher doses of the therapeutic compound than could be administered initially. A "conditioning dose" is a dose which attenuates or

5 reduces the frequency or the severity of first dose adverse side effects associated with administration of a therapeutic compound. The conditioning dose may be a therapeutic dose, a sub-therapeutic dose, a symptomatic dose or a sub-symptomatic dose. A therapeutic dose is a dose which exhibits a therapeutic effect on the patient and a sub-therapeutic dose is a dose which does not exhibit a therapeutic effect on the patient treated. A symptomatic dose is a dose which induces at least one adverse effect on administration and a sub-symptomatic dose is a dose which does not induce an adverse effect. Some adverse effects are fever, headache, nausea, vomiting, breathing difficulties, myalgia, and chills.

Route of administration

15 The BAFF antagonists and the anti-CD20 antibodies are administered to a human patient in accord with known methods, such as by intravenous administration, *e.g.*, as a bolus or by continuous infusion over a period of time, by subcutaneous, intramuscular, intraperitoneal, intracerebrospinal, intra-articular, intrasynovial, intrathecal, or inhalation routes. The anti-CD20 antibody will generally be administered by intravenous or subcutaneous administration. The drugs can be administered by the same or different route.

20

Articles of Manufacture and Kits

Another embodiment of the invention is an article of manufacture comprising a BAFF antagonist and an anti-CD20 antibody useful for the treatment of a B cell based malignancy or a B-cell regulated autoimmune disorder disclosed above. In a specific embodiment, the article of manufacture contains the BR3-Fc of SEQ ID 2, and the hu2H7v.16 antibody, for the treatment of non-Hodgkin's lymphoma.

25 The article of manufacture comprises at least one container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, *etc.* The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition of the invention which is effective for treating the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is a CD20 binding antibody of the invention such as RituxanTM or hu2H7v.16, and the other active agent is a BAFF antagonist such as BR3-Fc. The label or package insert indicates that the composition is used for treating the particular condition, *e.g.*, non-Hodgkin's lymphoma or rheumatoid arthritis. The label or package insert will further comprise instructions for administering the composition to the patient. Additionally, the article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

40 Kits are also provided that are useful for various purposes, *e.g.*, for B-cell killing assays. As with the article of manufacture, the kit comprises a container and a label or package insert on or associated with the container. The container holds a composition comprising at least one anti-CD20 antibody and one BAFF antagonist of the invention. Additional containers may be included that contain, *e.g.*, diluents and buffers,

- 5 control antibodies. The label or package insert may provide a description of the composition as well as instructions for the intended in vitro or diagnostic use.

10 Experimental Examples

Example 1

Development of monoclonal antibodies to CD20

- Six Balb/c mice, three Lewis rats (Charles River Laboratories, Hollister, CA) and three Armenian hamsters (Cytogen Research and Development, Inc., Boston, MA) were hyperimmunized with adenovirus-infected human 293 cells transiently expressing murine CD20 (Genentech, Inc., South San Francisco, CA), in phosphate buffered saline (PBS). Pre-fusion boosts, consisting of murine cells expressing endogenous levels of murine CD20 and purified, purified recombinant mCD20 expressed in *E. coli* (Genentech, Inc., South San Francisco, CA), were administered three days prior to fusion. B-cells from all animals were fused with mouse myeloma cells (X63.Ag8.653; American Type Culture Collection, Manassas, VA) using a modified protocol analogous to one previously described (Kohler and Milstein, 1975; Hongo et al., 1995). After 10-14 days, the supernatants from the mouse and hamster fusions were harvested and screened for anti-murine CD20 and anti-human CD20 antibody production by direct enzyme-linked immunosorbent assay (ELISA). The mCD20 ELISA screen identified 59 positive hybridomas from the hamster fusion (2 crossreactive with hCD20) and 1 positive hybridoma from the mouse fusion (crossreactive with hCD20). Positive clones, showing the highest immunobinding after subcloning by limiting dilution, are either injected into Pristane-primed mice (Freund and Blair, 1982) for *in vivo* production of Mab or cultured in vitro. The ascites fluids and/or supernatants are pooled and purified by affinity chromatography (Pharmacia fast protein liquid chromatography [FPLC]; Pharmacia, Uppsala, Sweden) as previously described (Hongo et al., 1995) or using a modified version of the protocol previously described. The purified antibody preparations are sterile filtered (0.2- μ m pore size; Nalgene, Rochester NY) and stored at 4°C in PBS.

Direct ELISA for the evaluation of immune sera

- Microtiter plates (NUNC) were coated with 100 μ l/well of murine or human CD20 (1 μ g/ml; Genentech, Inc., South San Francisco, CA) in 0.05 M carbonate buffer, pH 9.6. Coated plates were washed three times with ELISA wash buffer (PBS/0.05% Tween 20) and blocked for at least 1 hr with PBS containing 0.5% bovine serum albumin and 0.05% Tween 20 (PBS/BSA/T20). Blocking buffer was then removed and 100 μ l of diluted samples and controls were added and incubated for 1 hr at ambient temperature. The plates were then washed and horseradish peroxidase conjugated species specific anti-IgG conjugate (Sigma, St. Louis, MO or ICN Cappel, Durham, NC) was added (100 μ l/well) and incubated for 1 hr at ambient temperature. The plates were washed and incubated with tetramethylbenzidine substrate (BioFX Laboratories, Owings Mills, MD) for 5-10 minutes followed by the addition of Stop Solution (100 μ l/well; BioFX Laboratories). The plates were then read using an automated plate reader (EL808, BioTek Instruments, Inc., Winooski, VT).

5 **References:**

Hongo, J.S., Mora-Worms, M., Lucas, C. and Fendly, B.M.: Development and characterization of murine monoclonal antibodies to the latency-associated peptide of transforming growth factor β 1. *Hybridoma* 1995; 14:253-260.

10 Kohler, G. and Milstein, C.: Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature* 1975; 256: 495-497.

Freund YR and Blair PB: Depression of natural killer activity and mitogen responsiveness in mice treated with pristane. *J Immunol* 1982; 129:2826-2830.

Example 2**Humanization of 2H7 anti-CD20 murine monoclonal antibody**

15 Humanization of the murine anti-human CD20 antibody, 2H7 (also referred to herein as m2H7, m for murine), was carried out in a series of site-directed mutagenesis steps. The CDR residues of 2H7 were identified by comparing the amino acid sequence of the murine 2H7 variable domains (disclosed in U.S. 5,846,818) with the sequences of known antibodies (Kabat et al., *Sequences of proteins of immunological interest*, Ed. 5. Public Health Service, National Institutes of Health, Bethesda, MD (1991)). The CDR regions
20 for the light and heavy chains were defined based on sequence hypervariability (Kabat et al., *supra*) and are shown in Fig. 12 and Fig. 13, respectively. Using synthetic oligonucleotides (Table 2), site-directed mutagenesis (Kunkel, *Proc. Natl. Acad. Sci.* 82:488-492 (1985)) was used to introduce all six of the murine 2H7 CDR regions into a complete human Fab framework corresponding to a consensus sequence $V_{\kappa I}, V_{H III}$ (V_L kappa subgroup I, V_H subgroup III) contained on plasmid pVX4. The phagemid pVX4 was used for
25 mutagenesis as well as for expression of F(ab)s in *E. coli*. Based on the phagemid pb0720, a derivative of pB0475 (Cunningham et al., *Science* 243: 1330-1336 (1989)), pVX4 contains a DNA fragment encoding a humanized consensus κ -subgroup I light chain ($V_L \kappa I-C_L$) and a humanized consensus subgroup III heavy chain ($V_H III-C_H1$) anti-IFN- α (interferon α) antibody. pVX4 also has an alkaline phosphatase promoter and Shine-Dalgarno sequence both derived from another previously described pUC119-based plasmid, pAK2 (Carter et al.,
30 *Proc. Natl. Acad. Sci. USA* 89: 4285 (1992)). A unique *SpeI* restriction site was introduced between the DNA encoding for the F(ab) light and heavy chains. The first 23 amino acids in both anti-IFN- α heavy and light chains are the *SvII* secretion signal sequence (Chang et al., *Gene* 55: 189-196 (1987)).

To construct the CDR-swap version of 2H7 (2H7.v2), site-directed mutagenesis was performed on a deoxyuridine-containing template of pVX4; all six CDRs of anti-IFN- α were changed to the murine 2H7 CDRs.
35 The resulting molecule is referred to as humanized 2H7 version 2 (2H7.v2), or the "CDR-swap version" of 2H7; it has the m2H7 CDR residues with the consensus human FR residues shown in Figures 12 and 13. Humanized 2H7.v2 was used for further humanization.

Table 2 shows the oligonucleotide sequence used to create each of the murine 2H7 (m2H7) CDRs in the H and L chain. For example, the CDR-H1 oligonucleotide was used to recreate the m2H7 H chain CDR1.
40 CDR-H1, CDR-H2 and CDR-H3 refers to the H chain CDR1, CDR2 and CDR3, respectively; similarly, CDR-L1, CDR-L2 and CDR-L3 refers to each of the L chain CDRs. The substitutions in CDR-H2 were done in two steps with two oligonucleotides, CDR-H2A and CDR-H2B.

5 **Table 2.** Oligonucleotide sequences used for construction of the CDR-swap of murine 2H7 CDRs into a human framework in pVX4. Residues changed by each oligonucleotide are underlined.

Substitution	Oligonucleotide sequence
CDR-H1	C TAC ACC TTC ACG <u>AGC</u> TAT <u>AAC</u> <u>ATG</u> CAC TGG GTC CG (SEQ ID NO. 54)
CDR-H2A	G ATT AAT CCT GAC <u>AAC</u> <u>GGC</u> <u>GAC</u> ACG <u>AGC</u> TAT AAC CAG <u>AAG</u> TTC AAG GGC CG (SEQ ID NO. 55)
CDR-H2B	GAA TGG GTT GCA <u>GCG</u> ATC <u>TAT</u> CCT <u>GGC</u> AAC GGC GAC AC (SEQ ID NO. 56)
CDR-H3	AT TAT TGT GCT CGA GTG <u>GTC</u> <u>TAC</u> TAT <u>AGC</u> <u>AAC</u> <u>AGC</u> <u>TAC</u> <u>TGG</u> <u>TAC</u> <u>TTC</u> GAC <u>GTC</u> TGG GGT CAA GGA (SEQ ID NO. 57)
CDR-L1	C TGC ACA GCC AGC <u>TCT</u> TCT <u>GTC</u> AGC TAT ATG CAT TG (SEQ ID NO. 58)
CDR-L2	AA CTA CTG ATT TAC <u>GCT</u> <u>CCA</u> <u>TGC</u> AAC CTC <u>GCG</u> TCT GGA GTC C (SEQ ID NO. 59)
CDR-L3	TAT TAC TGT CAA CAG <u>TGG</u> <u>AGC</u> <u>TTC</u> <u>AAT</u> CCG <u>CCC</u> ACA TTT GGA CAG (SEQ ID NO. 60)

Based on a sequence comparison of the murine 2H7 framework residues with the human V_HL_HV_HIII
 10 consensus framework (Figures 12 and 13) and previously humanized antibodies (Carter et al., *Proc. Natl. Acad. Sci. USA* 89:4285-4289 (1992)), several framework mutations were introduced into the 2H7.v2 Fab construct by site-directed mutagenesis. These mutations result in a change of certain human consensus framework residues to those found in the murine 2H7 framework, at sites that might affect CDR conformations or antigen contacts. Version 3 contained V_H(R71V, N73K), version 4 contained V_H(R71V), version 5 contained V_H(R71V, N73K)
 15 and V_L(L46P), and version 6 contained V_H(R71V, N73K) and V_L(L46P, L47W).

Humanized and chimeric Fab versions of m2H7 antibody were expressed in *E. coli* and purified as follows. Plasmids were transformed into *E. coli* strain XL-1 Blue (Stratagene, San Diego, CA) for preparation of double-and single-stranded DNA. For each variant, both light and heavy chains were completely sequenced using the dideoxynucleotide method (Sequenase, U.S. Biochemical Corp.). Plasmids were transformed into *E.*
 20 *coli* strain 16C9, a derivative of MM294, plated onto LB plates containing 5 µg/ml carbenicillin, and a single colony selected for protein expression. The single colony was grown in 5 ml LB-100 µg/ml carbenicillin for 5-8 h at 37° C. The 5 ml culture was added to 500 ml AP5-100 µg/ml carbenicillin and allowed to grow for 16 h in a 4 L baffled shake flask at 37°C. AP5 media consists of: 1.5g glucose, 11.0 Hycase SF, 0.6g yeast extract (certified), 0.19g anhydrous MgSO₄, 1.07g NH₄Cl, 3.73g KCl, 1.2g NaCl, 120 ml 1 M triethanolamine, pH 7.4,
 25 to 1 L water and then sterile filtered through 0.1 µm Sealkeen filter.

Cells were harvested by centrifugation in a 1 L centrifuge bottle (Nalgene) at 3000xg and the supernatant removed. After freezing for 1 h, the pellet was resuspended in 25 ml cold 10 mM MES-10 mM EDTA, pH 5.0 (buffer A). 250 µl of 0.1M PMSF (Sigma) was added to inhibit proteolysis and 3.5 ml of stock 10 mg/ml hen egg white lysozyme (Sigma) was added to aid lysis of the bacterial cell wall. After gentle shaking
 30 on ice for 1 h, the sample was centrifuged at 40,000xg for 15 min. The supernatant was brought to 50 ml with buffer A and loaded onto a 2 ml DEAE column equilibrated with buffer A. The flow-through was then applied to a protein G-Sepharose CL-4B (Pharmacia) column (0.5 ml bed volume) equilibrated with buffer A. The column was washed with 10 ml buffer A and eluted with 3 ml 0.3 M glycine, pH 3.0, into 1.25 ml 1 M Tris, pH

8.0. The F(ab) was then buffer exchanged into PBS using a Centricon-30 (Amicon) and concentrated to a final volume of 0.5 ml. SDS-PAGE gels of all F(ab)s were run to ascertain purity and the molecular weight of each variant was verified by electrospray mass spectrometry.

Plasmids for expression of full-length IgG's were constructed by subcloning the V_L and V_H domains of chimeric Fab as well as humanized Fab of hu2H7 antibodies into previously described pRK vectors for mammalian cell expression (Gorman et al., *DNA Prot. Eng. Tech.* 2:3-10 (1990)). Briefly, each Fab construct was digested with *EcoRV* and *BlnI* to excise a V_L fragment, which was cloned into the *EcoRV/BlnI* sites of plasmid pDR1 for expression of the complete light chain (V_L-C_L domains). Additionally, each Fab construct was digested with *PvuII* and *ApaI* to excise a V_H fragment, which was cloned into the *PvuII/ApaI* sites of plasmid pDR2 for expression of the complete heavy chain (V_H-CH₁-CH₂-CH₃ domains). For each IgG variant, transient transfections were performed by cotransfecting a light-chain expressing plasmid and a heavy-chain expressing plasmid into an adenovirus-transformed human embryonic kidney cell line, 293 (Graham et al., *J. Gen. Virol.*, 36:59-74, (1977)). Briefly, 293 cells were split on the day prior to transfection, and plated in serum-containing medium. On the following day, double-stranded DNA prepared as a calcium phosphate precipitate was added, followed by pAdVantage™ DNA (Promega, Madison, WI), and cells were incubated overnight at 37°C. Cells were cultured in serum-free medium and harvested after 4 days. Antibodies were purified from culture supernatants using protein A-Sepharose CL-4B, then buffer exchanged into 10 mM sodium succinate, 140 mM NaCl, pH 6.0, and concentrated using a Centricon-10 (Amicon). Protein concentrations were determined by quantitative amino acid analysis.

Example 3

This example describes generation of human CD20 BAC transgenic (Tg) mice and experiments to study the effects of anti-CD20 antibody or BAFF antagonist alone in the hCD20+ mice.

Human CD20 transgenic mice were generated from human CD20 BAC DNA (Invitrogen, Carlsbad, CA). Mice were screened based on the FACS analysis of human CD20 expression. As can be seen from the FACS plots in Figure 14, mice hemizygous (Tg+/-) and homozygous (Tg+/+) for the transgene express human CD20 on their B220+ B cells. Figure 15 shows the expression of various cell surface markers (CD43, IgM, IgD) during B cell differentiation and maturation. In the Tg+ mice, hCD20 is expressed on pre-B and immature B cells and mostly on mature B cells. The Tg+ mice were screened for human CD20 expression in the B cells of the bone marrow, spleen, mesenteric LN and Peyer's patches; the results are shown in Figures 16-19. Gating the cells on B220 and CD43 allows delineation into the various populations of B cells. Tg+ mice were then treated with anti-CD20 mAb (1mg total = 50mg/kg, equivalent to 3.5mg for a 70kg man) to see the effects on the B cells as outlined in the schematic in Figure 20. FACS analyses were done on peripheral blood, spleen, lymph node, bone marrow, and Peyer's Patches. Serum levels of anti-CD20 mAb were monitored. In mice, B cell depletion occurs within 3-4 days of treatment with anti-CD20 antibody. Not to be bound by any theory, B cell death appears to be mediated by ADCC or apoptosis or both. Treatment of Tg+ mice with anti-hCD20 mAb (m2H7) alone results in depletion of B cells in peripheral blood, mature peripheral lymph node B cells, T2 and follicular B cells in the spleen (see Figures 21-24). However, it was observed that certain B cell subsets are resistant to killing by anti-CD20 antibody despite very high, likely saturating levels of antibody on the cell surface. These resistant B cells are the marginal zone B cells in the spleen (Fig. 23), and the germinal center B

5 cells in both the Peyer's patches (Fig. 25) and spleen (Fig. 27). In one experiment (Fig. 27), mice were injected with a first dose of anti-CD20 mAb at 100ug on day 1, followed by a second, 100ug dose on day 3 (it is likely that a single dose at 50ug was sufficient to saturate the B cells); T2/follicular B cells were depleted but the germinal center B cells from the Peyer's patches were shown to be bound with anti-CD20 mAb but were resistant to killing.

10 The recovery of B cells following anti-CD20 antibody treatment was followed. Mice were administered antibody at day 1. Figure 26 shows that at day 6 post antibody treatment, B cells in the peripheral blood were not detectable. At week 6, upon clearance of the antibody, hCD20+ cells begin to be detected and by week 14, B cells appeared to have recovered to normal levels. Recovery stems from precursor B cells which do not express CD20 developing into CD20+ mature B cells.

15 Figure 27 shows FACS plots demonstrating resistance of splenic germinal center B cells to short-term (single injection) anti-CD20 mAb treatment. Mice were unimmunized or immunized with sheep red blood cells (SRBC) by intraperitoneal injection at day 1 to induce germinal centers in the spleen. The germinal centers appear by day 7. At day 8, one group of mice was treated with the m2H7 mAb to human CD20. The control set of mice was treated with mIgG2a isotype control antibody. Spleen cells from the mice were analyzed at day 12. PNA (peanut agglutinin) stains for germinal center. No detectable germinal center cells were seen in the spleens of mice not immunized with SRBC whereas the spleens of immunized mice show 0.3% PNA staining cells. While T2/Follicular B cells are depleted with anti-CD20 antibody treatment, marginal center B cells in the spleen are resistant to the antibody.

25 Next, it was determined whether upon B cell depletion, the mice were able to develop T independent immune response. Mice were treated with m2H7 or isotype control antibody mIgG2a at day 0. At days 3-7, B cell depletion has occurred. At day 7, the mice were injected i.v. with *Streptococcus pneumoniae* IV to induce a response to the polysaccharide. A T cell independent response was mounted on day 11. The results shown in Figure 28 demonstrated that treatment with anti-CD20 (2H7 or Rituxan) did not affect the B cell response from the marginal zone and germinal centers of the spleen, i.e., the non-depleted MZ and B1 B cells confer protection to T-independent antigens. This data demonstrates that some aspects of humoral immunity-specifically T-independent B cell responses (in this case) are preserved despite treatment with anti-CD20 mAb.

Example 4

35 This example demonstrates the synergy between anti-CD20 mAb and BAFF antagonist treatments for B cell modulation/depletion.

BAFF/BLyS/TALL-1 (member of the TNF superfamily) plays an important role in the survival and maturation of immature T2, FO and MZ B cells and enhances competitive survival of autoreactive B cells (S. Mandal et al., *Science* 296, 346-9 (2002); F. Mackay, P. Schneider, P. Rennert, J. Browning, *Annu Rev Immunol* 21, 231-64 (2003); P. A. Moore et al., *Science* 285, 260-3 (1999)). Overexpression of a soluble form of BAFF/BLyS/TALL-1 in mice results in B cell hyperplasia, hypergammaglobulinemia and autoimmune lupus-like syndrome (S. A. Masters et al., *Curr Biol* 10, 785-8 (2000)). Conversely, treatment of lupus-prone mice with a BAFFR/BR3-Fc fusion protein, which neutralizes BAFF/BLyS, results in improved autoimmune serologies, renal pathology and mortality (R. Lesley et al., *Immunity* 20, 441-53 (2004)).

5

Materials and methods:

In the Experimental Examples, the BR3-Fc or BAFFR/BR3-Fc used is hBR3-Fc of SEQ ID. NO. 2. For the experiment shown in Fig. 29, FVB mice expressing a bacterial artificial chromosome encoding human CD20 (designated as hCD20⁺ mice) were treated with intraperitoneal injections of anti-CD20 mAb (single
 10 injection of 100 micrograms on Day 9), BR3-Fc (100 micrograms every other day from Days 1 through 12), or the combination of anti-CD20 mAb and BR3-Fc. Each group consisted of 4 mice. Two days following the last injection, the mice were sacrificed and analyzed for hCD20⁺ B cells. FACS analysis of spleen, blood, lymph node and Peyer's Patches were analyzed for B cell markers (CD21⁺CD23⁺).

For the experiment shown in Fig. 30, hCD20 Tg⁺ mice were treated with control IgG_{2a}, BAFFR/BR3-
 15 Fc (100 µg/mouse IP daily for 12 days), anti-hCD20 mAb (100 µg/mouse IP on day 9) or the combination of BAFFR/BR3-Fc and anti-hCD20 mAb (same dosing as single treatment groups). B220⁺ splenocytes were isolated on day 13 and stained for CD21 and CD23. N=5 mice/group. Figure 30 shows the synergistic effects on B cell depletion of the combination of anti-hCD20 mAb and BR3-Fc in the human CD20 Tg⁺ mice. Figure 31 shows quantitation of depletion of B220⁺ total spleen B cells, marginal zone (MZ) and follicular (FO) B cells
 20 from hCD20 Tg⁺ mice. The mice were treated with single doses of 0.1 mg control IgG_{2a}, BAFF/BR3-Fc or anti-hCD20 mAb. Splenocytes were analyzed on day 4. N=5 mice/group.

Results:

These results are shown in Fig. 29, Fig. 30, and Fig. 31.

- 25
1. Anti-CD20 mAb therapy depletes >99% of mature circulating B cells in the blood and lymph nodes.
 2. BR3-Fc decreases mature circulating B cells in the blood and lymph nodes.
 3. Anti-CD20 mAb therapy depletes T2 and follicular B cells, but not marginal zone B cells in the spleen.
 4. BR3-Fc decreases T2/follicular and marginal zone B cells in the spleen.
 5. The combination of anti-CD20 mAb and BR3-Fc synergizes to deplete all populations of B cells in the
 30 spleen.

Treatment of hCD20⁺ mice with BAFFR/BR3-Fc for ~2 weeks resulted in a marked decrease in MZ and T2/FO B cells (Fig. 30, panel 3). Combined treatment of BAFFR/BR3-Fc and anti-hCD20 mAb, surprisingly, resulted in the depletion of all splenic B cell subsets (Fig. 30, panel 4). To further explore the
 35 potential synergy of BAFF neutralization and anti-hCD20 mAb, the extent of B cell loss four days following treatment with single doses of anti-hCD20 mAb and BAFFR/BR3-Fc was quantified. While treatment with single doses of anti-hCD20 mAb or BAFFR/BR3-Fc resulted in ~40-50% loss of MZ B cells and ~33-70% loss of FO B cells, the combination anti-hCD20 mAb and BAFFR/BR3-Fc resulted in >90% loss of MZ and FO B cells (Fig. 31). Hence, survival factors also play an important role in determining susceptibility to anti-hCD20
 40 mAb mediated B cell depletion.

Example 5**Production of BAFF antagonists**

BAFF₈₂₋₂₈₃ production.

5 A DNA fragment encoding human BAFF (residues 82-285) was cloned into the pET15b (Novagen) expression vector, creating a fusion with an N-terminal His-tag followed by a thrombin cleavage site. *E. coli* BL21(DE3) (Novagen) cultures were grown to mid-log phase at 37°C in LB medium with 50 mg/L carbenicillin and then cooled to 16 °C prior to induction with 1.0 mM IPTG. Cells were harvested by centrifugation after 12 h of further growth and stored at -80°C. The cell pellet was resuspended in 50 mM Tris, pH 8.0, and 500 mM
10 NaCl and sonicated on ice. After centrifugation, the supernatant was loaded onto a Ni-NTA agarose column (Qiagen). The column was washed with 50 mM Tris, pH 8.0, 500 mM NaCl, and 20 mM imidazole and then eluted with a step gradient in the same buffer with 250 mM imidazole. BAFF-containing fractions were pooled, thrombin was added, and the sample was dialyzed overnight against 20 mM Tris, pH 8.0, and 5 mM CaCl₂ at 4°C. The protein was further purified on a monoQ (Pharmacia) column and finally on an S-200 size exclusion
15 column in 20 mM Tris, 150 mM NaCl, and 5 mM MgCl₂. The resulting BAFF protein was used as described below.

BR3 extracellular domain production

The extracellular domain of human BR3 (residues 1 to 61) was subcloned into the pET32a expression vector (Novagen), creating a fusion with an N-terminal thioredoxin (TRX)-His-tag followed by an enterokinase
20 protease site. *E. coli* BL21(DE3) cells (Novagen) were grown at 30°C and protein expression induced with IPTG. TRX-BR3 was purified over a Ni-NTA column (Qiagen), eluted with an imidazole gradient, and cleaved with enterokinase (Novagen). BR3 was then purified over an S-Sepharose column, refolded overnight in PBS, pH 7.8, in the presence of 3 mM oxidized and 1 mM reduced glutathione, dialyzed against PBS, repurified over a MonoS column, concentrated, and dialyzed into PBS.

25

Peptide synthesis

MiniBR3 was synthesized as a C-terminal amide on a Pioneer peptide synthesizer (PE Biosystems) using standard Fmoc chemistry. The side chain thiols of cysteines 19 and 32 were protected as trifluoroacetic acid (TFA)-stable acetamidomethyl (Acm) derivatives. Peptides were cleaved from the resin by treatment with
30 5% triisopropyl silane in TFA for 1.5–4 hr at room temperature. After removal of TFA by rotary evaporation, peptides were precipitated by addition of ethyl ether, then purified by reversed-phase HPLC (acetonitrile/H₂O/0.1% TFA). Peptide identity was confirmed by electrospray mass spectrometry. After lyophilization, the oxidized peptide was purified by HPLC. HPLC fractions containing reduced miniBR3 were adjusted to a pH of ~ 9 with NH₄OH; the disulfide between cysteines 24 and 35 was then formed by addition of
35 a small excess of K₃Fe(CN)₆, and the oxidized peptide purified by HPLC. Acm groups were removed (with concomitant formation of the second disulfide) by treatment of the HPLC eluate with a small excess of I₂ over ~ 4 h. The progress of the oxidation was monitored by analytical HPLC, and the final product was again purified by HPLC. MiniBR3 was amino-terminally biotinylated on the resin by reaction with a 10-fold molar excess of sulfo-NHS-biotin (Pierce Chemical, Co.). The biotinylated miniBR3 was then cleaved from the resin and
40 purified as described above for the unbiotinylated miniBR3.

The following peptides ECFDLLVRHWVACGLLR (BLyS0027) (SEQ ID NO:9), ECFDLLVRHWVPCGLLR (BLyS0048) (SEQ ID NO:6) and ECFDLLVRAWVPCSVLK (BLyS0051) (SEQ ID NO:5) were synthesized generally as follows. Peptides were synthesized on a Rainin Symphony peptide

5 synthesizer system using Rink amide resin and a threefold excess of 9-fluorenylmethoxycarbonyl (Fmoc) protected amino acid activated with 2-(1 H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) in the presence of a fivefold excess of diisopropylethylamine (DIPEA). Amino acids were coupled twice at each position before deprotecting with a 20% solution of piperidine in dimethylformamide (DMF) and moving to the next residue. Washes between coupling steps were performed
 10 using dimethylacetamide (DMA). Following coupling of the final amino acid onto the peptide and its deprotection with 20% piperidine in DMF, the peptides were acylated at their amino terminus using 3 equivalents of acetic anhydride and 5 equivalents of DIPEA in DMA. Alternatively, the amino terminus was modified through acylation with 5-carboxyfluorescein, with (+)-biotin, or through reaction with another fluorophore or reporter molecule. The peptide was then cleaved from the resin through treatment with a solution
 15 of 95% trifluoroacetic acid (TFA) containing 2.5% water and 2.5% triisopropylsilane for 90 minutes. The volatiles were removed under reduced pressure, diethyl ether was added and the solids filtered off. The resulting precipitate was washed again with diethyl ether and the combined organics discarded. The washed solids were then washed successively with acetic acid, a 1:1 mixture of acetic acid and acetonitrile, a 1:1:1 mixture of acetic acid, acetonitrile and water, an 1:1:8 mixture of acetic acid, acetonitrile and water and finally
 20 with water. The combined washes were lyophilized and the resulting crude peptides purified using C18 reverse phase high performance liquid chromatography using a 30 minute 10% to 70% gradient of acetonitrile in water with 0.1% trifluoroacetic acid in each solvent at a flow rate of 15 milliliters per minute. Fractions containing the desired peptide were oxidized through addition of a saturated solution of iodine in acetic acid until the solution remained colored. This solution was then lyophilized. Finally, the lyophilized crude oxidized peptide was
 25 purified a second time under identical conditions and the fractions containing the desired peptide lyophilized. Some of the peptides were synthesized under identical conditions except that the synthesis was performed on a PerSeptive Pioneer automated synthesizer using a fourfold excess of amino acid, coupling only once per residue.

Example 6

Phage Display of 17mers

30 *Library construction.* A phagemid encoding the STII secretion signal sequence ("STII ss"), a linker (GGGSGGG, SEQ ID NO: 61), and a sequence encoding the C-terminal residues of minor protein III of M13 phage (e.g., residues 267-421) (hereinafter, "cP3") was used as a template for library construction. Two libraries were constructed using Kunkel mutagenesis techniques and oligonucleotides that introduced a fragment
 35 corresponding to residues 23-39 of human BR3 with a C32W mutation, also known as "17-mer C32W", and additionally encoded mutations within the 17-mer C32W region. Specifically, library 1 encoded replacement codons at residues numbered 31, 34 and 36-39 (replacement codon: NNS = any codon), and library 2 encoded replacement codons residues 27, 30, 31, 34 and 36-39 (replacement codon: VNC = encodes amino acids L, P, H, R, I, T, N, S, V, A, D and G). In the replacement codons: N is 25% A, 25% C, 25% G, 25% T; S is 50% G/50%
 40 C; V is 33% G/33% A/33% C; and C is 100% C. Library 1 encoded 1.1×10^9 members and Library 2 encoded 4.3×10^8 members.

Library Sorting. The phage were subject to four rounds of selection. In general, the phage input per round was 10^{14} phage for the 1st round (solid phase sorting) and 3×10^{12} phage for additional rounds (solution phase sorting).

5 *Phage Selection.* The first round of selection was a solid phase sorting method. Maxisorp immunoplates (96-well) were coated with BLyS₈₂₋₂₈₅ prepared as described above (100µl at 2µg/ml in 50mM carbonate buffer (pH 9.6)) overnight at 4°C. The wells were then blocked for one hour with 0.2% (w/v) BSA in phosphate-buffered saline (PBS) and washed 3-5 times with PBS, 0.05% Tween20. Phage particles ((100µl/well in ELISA buffer (PBS/0.5%BSA /0.05% Tween20)) were added to the wells. After two hours, the
10 wells were washed several times with PBS, 0.05% Tween20. The phage bound to the wells were eluted with 0.1N HCl for 10 min at RT. The eluted phage were neutralized by adding 1/20 volume 2M Tris pH 11.0.

 To titer the phage, log phase XL-1 (OD 600nm~0.3) was infected with eluted phage at 37°C for 30 minutes. Next, the infected cells were serially diluted in 10 fold increments in 2YT. 10µl aliquots of the infected cells were plated per carbenicillin plate. ~10⁸ phage from each library were obtained from the first
15 round of selection.

 To propagate the phage, eluted phage was used to infect log phase XL-1 (OD 600nm~0.3) at 37°C for 30 minutes. Helper phage, KO7, and carbenicillin were added to the infection at a final concentration of 1 x 10¹⁰ pfu/ml KO7 and 50ug/ml carbenicillin at 37°C for another 30 minutes. The culture was grown in 2YT media with carbenicillin 50ug/ml and 25ug/ml kanamycin to final volumes of 25ml at 37°C overnight.

20 The phage were purified by spinning down the cells at 10000 rpm for 10 minutes. The supernatant was collected. 20% PEG/2.5M NaCl was added at 1/5 of the supernatant volume, mixed and allowed to sit at room temperature for 5 minutes. The phage were spun down into a pellet at 10000 rpm for 10 minutes. The supernatant was discarded and the phage pellet spun again for 5 minutes at 5000 rpm. The pellets were resuspended in 0.7ml PBS and spun down at 13000 rpm for 10 minutes to clear debris. The OD of the
25 resuspended phage pellet was read at 268nm.

 The second to fourth rounds of selection utilized solution sorting methods. For the second round, Maxisorp Nunc 96-well plates were coated with 5ug/ml neutravidin (Pierce) at 4°C overnight. Next, the plate was blocked with 200 µl/ml Superblock (Pierce) in PBS for 30 min at room temperature. Tween20 was added to each well for a final concentration of 0.2% (v/w) and blocked for another 30 minutes at room temperature.
30 The amplified, purified phage from the first round of selection were incubated with 50nM biotinylated BAFF (final concentration) in 150ul buffer containing Superblock 0.5% and 0.1% Tween20 for 1 h at room temperature. The mixtures were then diluted 5-10X with PBS/0.05% Tween and applied at 100µl/well to the neutravidin coated plate. The plate was gently shaken for five minutes at room temperature to allow phage bound to biotinylated BAFF to be captured in the wells. The wells were then washed with PBS/0.05% Tween20
35 several times. Bound phage were eluted with 0.1N HCl for 10 min, neutralized, titered, propagated and purified as described above. ~3 x 10⁶ phage from each library were obtained from the second round of selection.

 The third round of selection was similar to the second round, except a concentration of 2nM biotinylated BAFF was incubated with the phage prior to dilution and addition to each well. Bound phage were eluted with 0.1N HCl for 10 min, neutralized, titered and propagated as described above. ~10⁴ phage from each
40 library were obtained from the third round of selection.

 Phage from the third round of selection were next subjected to two different selection methods in the fourth round. Method 4a was similar to the second and third rounds of selection except that the phage was incubated in the presence of 0.5nM biotinylated BAFF for 1h at room temperature. The mixture was then incubated for an additional 15 minutes at room temperature in the presence of 1000 fold excess (500nM) of

5 unbiotinylated BAFF prior to dilution and addition to the coated wells. Method 4b was also similar to the second and third rounds of selection except that 0.2nM BAFF was incubated with the phage before dilution and addition to each well. Bound phage from each round four selection were eluted with 0.1N HCl for 10 min, neutralized, titered and propagated as described above. $\sim 10^3$ phage were obtained for each library from each of the fourth rounds (4a and 4b) of selection.

10 *Clone Analysis.* After the fourth round of selection, individual clones were grown in a 96-well format in 400 μ L of 2YT medium supplemented with carbenicillin and KO7 helper phage. Supernatants from these cultures were used in phage ELISAs. For phage ELISAs, Nunc Maxisorp 96-well plates were coated overnight at 4 °C with 100 μ L of a 2 μ g/ml solution of BAFF in carbonate buffer, pH 9.6. The plate was washed with PBS and blocked with 0.5% BSA in PBS for two hours. Phage supernatant was diluted 1:4 in ELISA binding buffer
15 (PBS, 0.5%BSA, 0.05% Tween20) in the absence or presence of 50nM BAFF and incubated for 1h at RT. 100 μ L of the diluted phage supernatants were then transferred to the coated plates and allowed to shake gently to capture phage for 20 minutes. The plates were then washed with PBS/0.05% Tween20 several times. 100 μ L per well of HRP-conjugated anti-M13 antibody in PBS/0.05% Tween20 (1:5000) was then transferred to the plates and incubated for 20 min. After washing with PBS/0.05%Tween followed by PBS, the plate was incubated 5
20 min with 100 μ L PBS substrate solution containing 0.8 mg/ml OPD (Sigma) and 0.01% H_2O_2 . The reaction was quenched with 100 μ L/well 1M H_3PO_4 and the plate read at 490 nm. The clones tested were then sequenced as previously described (Weiss, G. A., Watanabe, C. K., Zhong, A., Goddard, A., and Sidhu, S. S. (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97, 8950-8954). Sequences of acceptable quality were translated and aligned. The amino acid sequences of the 17mers are shown in FIG.32.

25 Fourteen clones were further analyzed in a BAFF binding assay to determine their IC50 value. Clones 2 and 7 had a high number of siblings (clones with an identical sequence) in the fourth round. According to the phage ELISA assay, clones 13, 19, 22, 26, 32, 39 and 44 were greatly inhibited from binding to the plate by 50nM BAFF (FIG.11). The binding of clones 35, 45, 68, 82 and 90 was also greatly inhibited in the phage ELISA assay (FIG.11). Phage supernatants from these 14 clones were used to infect log phase XL-1 which
30 were propagated and purified as described above.

To normalize for display and phage yield and determine the appropriate dilution of phage for IC50 measurement, serial dilutions of purified phage from each clone were incubated in ELISA binding buffer (PBS, 0.5%BSA, 0.05% Tween20) for 1 h at room temperature. 100 μ L of each dilution were transferred to BAFF coated plates and allowed to shake gently to capture phage for 20 minutes as described above. Bound phage
35 was detected by HRP-conjugated anti-M13 antibody, followed by OPD/ H_2O_2 substrate reaction, quenched and read at 490nm as described above. By this process, the dilution of each clone that yielded ~ 1 O.D. at 490nm was determined and used in the IC50 assay.

To determine the IC50 value of each of the 14 clones, Nunc Maxisorp 96-well plates were coated overnight at 4 °C with 100 μ L of a 2 μ g/ml solution of BAFF in carbonate buffer, pH 9.6, and washed and
40 blocked as described above. A dilution of amplified, purified phage for each of the 14 clones was incubated in the presence of a concentration series of BAFF ranging from 0.003-1000 nM in 130 μ L ELISA binding buffer (PBS, 0.5%BSA, 0.05% Tween20) for 1 h at room temperature. 100 μ L of each of these concentration series were transferred to BAFF coated plates and captured, washed, detected with HRP-conjugated anti-M13 antibody

5 and processed as described above. IC₅₀ values were determined by a four-parameter fit of the ELISA signal for each of the 14 clones. The IC₅₀ values ranged from 0.4 (clone 44) to 11 nM (clone 22).

Competitive Displacement ELISA. The following 17-mers, Ac-ECFDLLVRHWVACGLLR-NH₂ (SEQ ID NO: 9) ("BLyS0027"), Ac-ECFDLLVRHWVPCGLLR-NH₂ (SEQ ID NO: 6) ("BLyS0048"), Ac-ECFDLLVRAWVPCSVLK-NH₂ (SEQ ID NO: 5) ("BLyS0051") were synthesized as described above. Nunc
10 Maxisorp 96-well plates were coated overnight at 4°C with 100 µl of a 2 µg/ml solution of BAFF in carbonate buffer, pH 9.6. The plate was washed with PBS and blocked with 1% skim milk in PBS. Serial dilutions of the BR3 ECD (residues 1-61) and the above 17-mer peptides were prepared in PBS/0.05% Tween 20 containing 3 ng/ml biotinylated miniBR3. After washing with PBS/0.05% Tween, 100 µl/well of each dilution was transferred and incubated for 1 hour at room temperature. The plate was washed with PBS/0.05% Tween and
15 incubated 15 min with 100 µl/well of 0.1 U/ml Streptavidin-POD (Boehringer Mannheim) in PBS/0.05% Tween. After washing with PBS/0.05% Tween followed by PBS, the plate was incubated 5 min with 100 µl PBS substrate solution containing 0.8 mg/ml OPD (Sigma) and 0.01% H₂O₂. The reaction was quenched with 100 µl/well 1M H₃PO₄ and the plate read at 490 nm. IC₅₀ values were determined by a four-parameter fit of the competitive displacement ELISA signal. The concentrations of initial stock solutions of miniBR3 and BR3
20 extracellular domain were determined by quantitative amino acid analysis.

The IC₅₀ values were determined for BR3 ECD, BLyS0027, BLyS0048 and BLyS0051 using this assay. The 17-mer peptides all had greater affinity for BAFF than the 62-mer BR3 ECD.

25 Example 4

Peptide-PEG conjugates

BLyS₈₂₋₂₈₅ production and Peptide synthesis. As described in Example 5 above.

Conjugation of Polymers to Peptides. PEGylated 17-mer peptides were generated by using
30 linear PEGs modified with N-hydroxysuccinimide chemistry (NHS) to react with primary amines (lysines and N-terminus). All PEG-NHS (PEG-SPA) reagents were purchased from Nektar Therapeutics, San Carlos, CA and stored under nitrogen at -70°C. The peptide was dissolved at 1 mg/mL in phosphate-buffer saline (PBS). To 0.4 mL aliquots of the peptide solution was added solid 2KPEG-SPA, 5KPEG-SPA, or 20KPEG-SPA. Enough solid was added to obtain a 3:1 molar ratio of PEG-SPA to peptide. These solutions were incubated at
35 room temperature for 1 hour and then the progress of the reaction was analyzed by reverse phase analytical HPLC on a 50 µL portion of the solution. The PEG addition and incubation was repeated 2 times until all of the peptide had been modified. The PEGylated peptides were tested for BAFF binding without further purification. The ratio of PEG:peptide in the purified conjugated product is approximately 1:1.

Competitive Displacement ELISA. A 17-mer, Ac-ECFDLLVRHWVPCGLLR-NH₂ (SEQ ID NO: 6) ("blys0048") was synthesized as described above. ECFDLLVRHWVPCGLL K (blys0095) (SEQ ID NO: 62)
40 was synthesized and coupled to each of 2K, 5K and 20K PEG-NHS as described above. Nunc Maxisorp 96-well plates were coated overnight at 4°C with 100 µl of a 2 µg/ml solution of BLyS in carbonate buffer, pH 9.6. The plate was washed with PBS and blocked with 1% skim milk in PBS. Serial dilutions of mini-BR3 (SEQ. ID. 50) and the above 17-mer peptide and PEG-peptide conjugate were prepared in PBS/0.05% Tween 20

5 containing 3 ng/ml biotinylated miniBR3. After washing with PBS/0.05% Tween, 100 µl/well of each dilution was transferred and incubated for 1 hour at room temperature. The plate was washed with PBS/0.05% Tween and incubated 15 min with 100 µl/well of 0.1 U/ml Streptavidin-POD (Boehringer Mannheim) in PBS/0.05% Tween. After washing with PBS/0.05% Tween followed by PBS, the plate was incubated 5 min with 100 µl PBS substrate solution containing 0.8 mg/ml OPD (Sigma) and 0.01% H₂O₂. The reaction was quenched
10 with 100 µl/well 1M H₃PO₄ and the plate read at 490 nm. IC₅₀ values were determined by a four-parameter fit of the competitive displacement ELISA signal. The equation is: $y = m1 + (m2 - m1) / (1 + m0/m4)^{m3}$, where m1 is the absorbance at infinite competitor concentration, m2 is the absorbance for no added competitor, m3 is the slope of the curve near the midpoint, m4 is the IC50 value and m0 is the concentration of competitor, peptide in this case. The concentration of biotinylated miniBR3 was about 10 pM. The concentration of initial stock
15 solution of miniBR3 was determined by quantitative amino acid analysis.

Results

The four-parameter fit of the competitive displacement ELISA signals provided IC50 values for: blys0095 of 19nM, blys0048 of 14nM and blys0095-2kPEG conjugate of 43nM, and blys0095-5kPEG
20 conjugate of 51nM using this assay. Similarly, the fit of the competitive displacement ELISA signals for a separate experiment provided IC50 values for blys0095-20kPEG conjugate of 99nM and blys0048 of 15nM.

The 17-mer peptide-PEG conjugates (2k, 5k and 20k) demonstrated binding ability for BAFF. The conjugation of PEG to blys0095 did not significantly reduce its binding affinity as compared to similar
25 unconjugated peptides.

Conclusion

The experiments herein demonstrated surprising results in that the combination of anti-CD20 mAb and BR3-Fc resulted in great synergy in depletion of all subsets of B cells.

References

30 References cited within this application, including patents, published applications and other publications, are hereby incorporated by reference.

5 What is claimed is:

1. A method of depleting B cells from a mixed population of cells comprising contacting the mixed population of cells with a BAFF antagonist and a CD20 binding antibody.
2. The method of claim 1 wherein the B cells are human B cells and the mixed population of cells are contacted with a BAFF antagonist and the CD20 binding antibody in vivo.
- 10 3. The method of claim 1 wherein the BAFF antagonist is an immunoadhesin.
4. The method of claim 3, wherein the immunoadhesin is selected from the group consisting of BR3 immunoadhesin comprising the extracellular domain of BR3, TACI immunoadhesin comprising the extracellular domain of TACI, and BCMA immunoadhesin comprising the extracellular domain of BCMA.
5. The method of claim 4, wherein the BR3 immunoadhesin is BR3-Fc of SEQ ID No. 2.
- 15 6. The method of claim 1, wherein the BAFF antagonist is an anti-BAFF antibody.
7. The method of claim 6, wherein the anti-BAFF antibody binds BAFF within a region of BAFF comprising residues 162-275.
8. The method of claim 1, wherein the BAFF antagonist is an anti-BR3 antibody.
9. The method of claim 8, wherein the anti-BR3 antibody binds BR3 in a region comprising
20 residues 23-38 of human BR3.
10. The method of claim 1, wherein the anti-CD20 antibody is RituxanTM.
11. The method of claim 1, wherein the anti-CD20 antibody is hu2H7v.16 having the light and heavy chain sequence of SEQ ID NO.15 and SEQ ID NO.16, respectively.
12. The method of claim 1 wherein the BAFF antagonist and the anti-CD20 antibody act
25 synergistically to deplete the B cells.
13. A method of treating a B cell neoplasm or malignancy characterized by B cells expressing CD20, comprising administering to a patient suffering from the neoplasm or malignancy, a therapeutically effective amount of a CD20 binding antibody and of a BAFF antagonist.
14. The method of claim 13, wherein the CD20 binding antibody and BAFF antagonist are
30 administered concurrently.
15. The method of claim 13, wherein the CD20 binding antibody and BAFF antagonist are administered sequentially.
16. The method of claim 13, wherein the BAFF antagonist is administered before the CD20 binding antibody.
- 35 17. The method of claim 13, wherein the B cell neoplasm is non-Hodgkin's lymphoma (NHL), small lymphocytic (SL) NHL, lymphocyte predominant Hodgkin's disease (LPHD), follicular center cell (FCC) lymphomas, acute lymphocytic leukemia (ALL), chronic lymphocytic leukemia (CLL), diffuse large B cell lymphoma, and Hairy cell leukemia.
18. The method of claim 17, wherein the B cell neoplasm is non-Hodgkin's lymphoma (NHL) or
40 chronic lymphocytic leukemia.
19. The method of claim 13, wherein the BAFF antagonist is an immunoadhesin.
20. The method of claim 19, wherein the immunoadhesin is selected from the group consisting of BR3 immunoadhesin comprising the extracellular domain of BR3, TACI immunoadhesin comprising the extracellular domain of TACI, and BCMA immunoadhesin comprising the extracellular domain of BCMA.

- 5 21. The method of claim 20, wherein the BR3 immunoadhesin is hBR3-Fc of SEQ ID NO. 2.
22. The method of claim 13, wherein the BAFF antagonist is an anti-BAFF antibody.
23. The method of claim 14, wherein the anti-BAFF antibody binds BAFF within a region of
BAFF comprising residues 162-275.
24. The method of claim 13, wherein the BAFF antagonist is an anti-BR3 antibody.
- 10 25. The method of claim 8, wherein the anti-BR3 antibody binds BR3 in a region comprising
residues 23-38 of human BR3.
26. The method of claim 13, wherein the CD20 binding antibody is a chimeric antibody
comprising the variable regions from a murine antibody fused to the constant regions of a human antibody.
27. The method of claim 26, wherein the chimeric antibody is RituxanTM.
- 15 28. The method of claim 13, wherein the CD20 binding antibody is a humanized antibody.
29. The method of claim 28, wherein the humanized antibody is hu2H7v.16 having the light and
heavy chain sequence of SEQ ID NO.15 and SEQ ID NO.16, respectively.
30. The method of claim 21, wherein the CD20 binding antibody is RituxanTM or hu2H7v.16
having the light and heavy chain sequence of SEQ ID NO.15 and SEQ ID NO.16, respectively.
- 20 31. The method of claim 30, wherein BR3-Fc is administered at a dosage of about 2-5mg/kg and
Rituxan is administered at a dosage of about 375mg/m².
32. The method of claim 13, wherein administration of the BAFF antagonist and the CD20
binding antibody produces a synergistic effect to deplete the B cells.
33. The method of claim 13, wherein the BAFF antagonist and the CD20 binding antibody are
25 administered in conjunction with chemotherapy.
34. A method of alleviating a B-cell regulated autoimmune disorder comprising administering to a
patient suffering from the disorder, a therapeutically effective amount of a CD20 binding antibody and of a
BAFF antagonist.
35. The method of claim 34, wherein the CD20 binding antibody and BAFF antagonist are
30 administered sequentially.
36. The method of claim 34, wherein the BAFF antagonist is administered before the CD20
binding antibody.
37. The method of claim 34, wherein the autoimmune disorder is selected from the group
consisting of rheumatoid arthritis, juvenile rheumatoid arthritis, systemic lupus erythematosus (SLE), lupus
35 nephritis, Wegener's disease, inflammatory bowel disease, idiopathic thrombocytopenic purpura (ITP),
thrombotic thrombocytopenic purpura (TTP), autoimmune thrombocytopenia, multiple sclerosis, psoriasis, IgA
nephropathy, IgM polyneuropathies, myasthenia gravis, vasculitis, diabetes mellitus, Reynaud's syndrome,
Sjorgen's syndrome and glomerulonephritis.
38. The method of claim 37, wherein the autoimmune disorder is rheumatoid arthritis and
40 systemic lupus erythematosus.
39. The method of claim 34, wherein the BAFF antagonist is an immunoadhesin.
40. The method of claim 39, wherein the immunoadhesin is selected from the group consisting of
BR3 immunoadhesin comprising the extracellular domain of BR3, TACI immunoadhesin comprising the
extracellular domain of TACI, and BCMA immunoadhesin comprising the extracellular domain of BCMA.

- 5 41. The method of claim 40, wherein the BR3 immunoadhesin is BR3-Fc of SEQ ID No. 2.
42. The method of claim 34, wherein the BAFF antagonist is an anti-BAFF antibody.
43. The method of claim 34, wherein the BAFF antagonist is an anti-BR3 antibody.
44. The method of claim 34, wherein the CD20 binding antibody is a chimeric antibody
comprising the variable regions from a murine antibody fused to the constant regions of a human antibody.
- 10 45. The method of claim 44, wherein the chimeric antibody is RituxanTM.
46. The method of claim 34, wherein the CD20 binding antibody is a humanized antibody.
47. The method of claim 28, wherein the humanized antibody is hu2H7v.16 having the light and
heavy chain sequence of SEQ ID NO.15 and SEQ ID NO.16, respectively.
48. The method of claim 41, wherein the CD20 binding antibody is RituxanTM or hu2H7v.16
15 having the light and heavy chain sequence of SEQ ID NO.15 and SEQ ID NO.16, respectively.
49. The method of claim 48, wherein BR3-Fc is administered at a dosage of about 2-5mg/kg and
Rituxan is administered at a dosage of about 2.5-10mg/kg.
50. The method of claim 34, wherein administration of the BAFF antagonist and the CD20
binding antibody produces a synergistic effect to deplete the B cells.
- 20 51. The method of claim 38, wherein the BAFF antagonist and the CD20 binding antibody is
administered in conjunction with therapy using a drug selected from nonsteroidal anti-inflammatory drugs
(NSAIDs), glucocorticoid, prednisone, and disease-modifying antirheumatic drug (DMARD).
52. A method of depleting marginal zone or germinal center B cells in a patient suffering from a B
cell neoplasm or a B-cell regulated autoimmune disorder, comprising administering to a patient in need thereof,
25 a therapeutically effective amount of a CD20 binding antibody and of a BAFF antagonist.
53. The method of any one of claims 1, 13, 34 and 52, wherein the BAFF antagonist is selected
from the group consisting of a polypeptide having the sequence of SEQ ID NO. 5, SEQ ID NO. 6, SEQ ID
NO.7 SEQ ID NO. 8, SEQ ID NO. 9, or SEQ ID NO. 10.
54. A composition comprising a CD20 binding antibody and a BAFF antagonist.
- 30 55. An article of manufacture comprising CD20 binding antibody, a BAFF antagonist, and a label
wherein the label indicates that the composition is for treating a B cell neoplasm or a B cell regulated
autoimmune disorder.
56. The method of claim 4 wherein the BR3 immunoadhesin comprises the sequence of any one of
SEQ ID NO.145-165 in FIG. 33 or a minimal region thereof.
- 35 57. The method of claim 20, wherein the BR3 immunoadhesin comprises the sequence of any one of
SEQ ID NO.145-165 in FIG. 33 or a minimal region thereof.
58. The method of claim 40, wherein the BR3 immunoadhesin comprises the sequence of any one of
SEQ ID NO.145-165 in FIG. 33 or a minimal region thereof.

1 AGCATCTGA GTAATAGTG GCTGGGCGG GAGCAGGCA GGTGGCGGA GCGTGTGA CCAGGAGG AGCTTTCCAC AGGCCCTGTG GAGGGGGTG
 1 TCGTAGGACT CATTACTAC OGGACCGGC CTGCTCCGT CCACGCGCT CGGCACACT GGTCTCTCTC GCGAAGGTG TCCGGACAC CTGCCCCAC
 M S G L G R S R R G G R S R V D Q E E R F P Q G L W T G V
 101 GCTATGAGT CTTGCCCGA AGAGCAGTAC TGGATCTTC TGTGGTAC CTGATGTCC TGCAAAACA TTTCARCCA TCAGAGCCAG GCAACTGTG
 CGTACTCTA GGAAGGGCT TCTGTCTAG ACCCTAGGAG AGGACCATG GAGTACAGG ACCTTTTGT AAGCTTGT AAGCTTGT AGCTGTGT GCTGGACAC
 30 A M R S C P E E Q Y W D P L L G T C M S C K T I C N H Q S Q R T C A
 201 CAGCCTCTG CAGTCACTC AGCTGCCGA AGAGCAAG CAGTTCTAT GACCATCTCC TGAGGACTG CATCAGCTGT GCTCCATCT GTGACAGCA
 GTCCGAGAC GTCCAGTCAG TCGAGCGCT TCTCGTTCC GTTCAGATA CTGTAGAGG ACTCCCTGAC GTAGTCGACA CGAGGTAGA CACTGTCTGT
 64 A F C R S L S C R K E Q G K F Y D H L L R D C I S C A S I C G Q H
 301 CCTBAGCAA TGTGCTACT TCTGTAGAA CAGCTCAG AGCCCTAGT ACCTCCACC AGAGCTCAG AGACAGGGA GTGAGGAGT TGAAGACAT
 GGAATCTGT ACACGTATGA AGACATCTT GTTCAGTCC TGGGTCACT TGGAGGTG TCTCGAGTCC TCTGTGCT CACTCTTCA ACTTTGTTA
 97 P K Q C A Y F C E N K L R S P V N L P P E L R R Q R S G E V E N N
 401 TCAGCAACT CGGAGGTA CCAGGATG GAGCAGAG GCTCAGAGC AGTCCAGCT CTCCGGGCG TGAGCTCAG TCAGTCTAG GTGGCCTGG
 AGTCTGTGA GCCCTTCACT GGTCTTCACT CTGTCTCTC CAGTCTCTG TTCTAGTCA GAGGCGCGG ACTTCACTC ACTCTAGT CACCGGACC
 130 S D N S G R Y Q G L E H R G S E A S P A L P G L K L S A D Q V A L V
 501 TCTACAGCAC GCTGGGCTC TGCCTCTG CTGCTCTG GTGGGGTG CCTGCTTCT CAGAGAGG GGGATCCCT GCTCTGACA
 AGATGTCTG OGAOCCGAG AGGACACAC GGCAGAGAC GAGGAGAC CACGCCAC GGTCTCTCTC CACTCTCTC CACTCTCTC CACTCTCTC
 164 Y S T L G L C L C A V L C C F L V A V A C F L K K R G D P C S C Q
 601 GCGCGCTCA AGGCCCGTC AAGTCCGCG CAGTCTTCC CAGGATCAG CGATGAGG CCGGAGCCCT GTAGCAGAT CCGCGAGCC AGTGAAGAT
 CCGGGCGGT TCGGGGCGG TTTCAGGCG GTTCAAGG GTCTAGTCT GCTACTCTG GCGTGGGA CACTCTCTA GGGGCTGG TCACTCTG
 197 P R S R P R Q S P A K S S Q D H A M E A G S P V S T S P E P V E T
 701 TCAGCTTCT GCTTCTCTA GTGAGGCG CCGAGGCG AGAGGCGAGT CAGCTGAGT AGCCCGAC CACTCTCTC TGAAGTGG GGGTCCACA
 AGCTGAGGA CCAAGGAGT CAGTCTCTC GGTGCTCTC TCTGCTCA GTGGGCTG GGTGAGAGG ACTTCTCAC CCACTCTC
 230 C S F C F P E C R A P T Q E S A V T P G T P D P T C A G R W G C H T
 801 CCAGGACCAC AGTCTCTG CTTTCCAC ACATCCGGA CAGTCTCTC GGTCTCTCT GGTCTCTCT CAGAGGAG GGGTCCAGT CATTAAGG
 GGTCTCTG TCAAGCTC GGAAGGCT TGTAGGCT GTACCGGA CCGTACACA CACAGGAG GGTCTCTCT CCGGTCTCT GTATTATCC
 264 R T T V L Q P C P H I P D S G L G I V C V P A Q E G G P G A O

FIG. 1A

901 GGTGAGGAG GGAAGAGAG AGGAGAGAG ATGGAGAGGA GGGAGAGAG AAGAGAGGT GGGAGAGGG GAGAGAGATA TGAGAGAGA GAGACAGAGG
CCAGTCCCTC CCTTCCCTCC TCCCTCTCTC TACCTCTCCT CCGCTCTCTC TTCTCTCCA CCGCTCTCC CTCTCTCTAT ACTCTCTCT CTCTGCTCTCC

1001 AGGCAGAAAG GCAGAGAAAC AGAGAGACA GAGAGGGAGA GAGAGACAGA GGGAGAGAGA GAGAGAGGG AAGAGAGGCA GAGAGGGAAA GAGGCAGAGA
TCCGTCCTTC CCTCTCTTG TCTCTCTGT CTCTCCCTCT CTCTCTGTCT CCGCTCTCT CTCTCTCCC TTCTCTCGT CTCTCCCTTT CTCCGCTCTCT

1101 AGGAAGAGA CAGGCAGAGA AGGAGAGAG CAGAGAGGA GAGAGCAGA GAGGAGAGA GGCAGAGAGA CAGAGAGGA GAGAGGGACA GAGAGAGATA
TCCCTTCTCT GTCCGTCTCT TCCTCTCTCC GTCTCTCCCT CTCTCCGTCT CTCTCTCTCT CCGCTCTCT GTCTCTCGT CTCTCCCTGT CTCTCTCTAT

1201 GAGCAGAGG TCGGGGCACT CTGAGTCCCA GTTCCAGTG CAGCTGTAGG TCGTCTATCAC CTAACCAAC CTAACCAAC GTCTCTCTCT GTCTCTCTCA
CTCGTCTCTCC AGCCCGCTGA GACTCAGGT CAAGGCTCAC GTCGACATCC AGCAGTAGTG GATTGCTGT CACGTTATTT CAGGAGCAG GACGAGAGT

1301 CAGCCCCCGA GAGCCCCCTCC TCCTGGAGAA TAAACCTTT GGCAGCTGCC CTTCCTCATA AAAAAAAA AAAAAA
GTCCGGGGCT CTCGGGGAGG AGGACCTCTT ATTTGGAAA CCGTCGACGG GAGGAGCTTT TTTTTTTT TTTTTT

FIG. 1B

1 AAGACTCATA CTATGAGACT TGAATTTAGT GATGATATTA AATCTTTAG TGGGCGTAG ACACAGACAG CCCCCTGTAAG AACCCACGAA GCAGGCGAAG
 TTCTGAGTTT GAATCTTTGA ACTTAATCTA CACCATTAAGT TTAGGAATGC ACGGCGCTTC TGTGTCTCTC EGGGCGATTC TTGGGTGCTT CGTCCGCTTC
 101 TTCAATGTTT TCACATCTT AGCTGCTCTT GCTGCATTTG GCTCGAATT CTGTAGAGT TATTACTTGT CCTCCAGGC TGTCTTTTCT GTAGCTOCTT
 AAGTAACAAG AGTTGTAGA TCGACGAGAA CGAGCTAAC GAGACCTTAA GAGACCTTCT AATATGAACA GGAAGGTCCG ACAGAGAGA CATCGAGGGA
 201 TGTCTTTCTT TTGTGATCAT GTTCAGATG GCTGGGCAGT GCTCCAGAA TGAATATTTT GACAGTTTGT TGCATGCTTG CATACCTTGT CAATCTOCTT
 ACAAAAGAAA AACACTAGTA CAACGTCTAC CGACCCGCTA CGAGGCTTTT ACTTATAAA CTGTCAAACA ACCTACGAC GTATGGAACA GTTGAAGCTA
 301 GTTCTTTCTA TACTOCTCTT CTAAACATGC AGCTTATTG TAATGCAAGT GTGACCAATT CAGTGAAGG AACGAATCG ATTCTCTGGA CCTGTTTGGG
 CAAGAAGATT ATGAGGAGGA GATGTACAG TCGCAATAAC ATTACGTTCA CACTGTTAA GTCACTTTCC TTGCTTACGC TAAGAGACCT GACCAAAACC
 29 ysSerSerAs nThrProPro LeuThrCysG InArgfyrCy sAsnAlaSer ValThrAsnS erVallySgl yThrAsnAla IleLeuThrP rCysLeuGl
 401 ACTGAGCTTA ATAATTCTT TGGCAGTTT CGTGCTAATG TTTTGTCTAA GGAAGATAAG CTCTGAACCA TTAAAGGACG AGTTTAAATA CACAGGATCA
 TGAATCGAAT TATTAAGAA ACCGTCAAAA GCACGATTAC AAAACAGATT CCTTCTATTG GAGACTTGGT AATTCTCTGC TCAATTTTTT GTGTCTTAGT
 62 yLeuSerLeu IleLeSerL euAlaValPh eValLeuMet PheLeuLeuA rglYsIkeSe rSerGluPro LeuYsAspG lUpheYsAs nThrGlySer
 501 GGTCTCTGG GATGGCTAA CATTGACCTG GAAAGAGCA GGACTGCTGA TGAATTTATT CTTCGAGAG GCTTCAGTA CACGGTGGAA GAATGCACTT
 CCAGAGGACC CGTACCGATT GTAACTGAC CTTTCTCTGT CCTGACACT ACTTAATAA GAAGCTCTC CGAGCTCAT GTGCCACCTT CTACGTGGA
 95 GlyLeuLeuG lYMetAlaAs nIleAspLeu GlulYsSera rglYsGlyAs pGluIlelle LeuProArgG lYLeuGluTy rThrValGlu GluCysThrC
 601 GTGAAGACTG CATCAAGAGG AACCAGAGG TCGACTCTGA CCATTGCTTT CCACCTCCAG CTATGGAGGA AGCGCAACC ATTCTTGTCA CCACGAAAGC
 CACTTCTGAC GTAGTTCTCG TTTGGCTTCC AGCTGAGACT GGTAACGAAA GGTCAGGCTC GATACCTCTT TCCGCTTGG TAAGAACAGT GGTGCTTTTG
 129 yGluAspCy sIleYsSer LysProYsV alAspSerAs pHisCysPhe ProLeuProA laMetGluGl uGlyAlaThr IleLeuValt hrThrIysTh
 701 GAATGACTAT TGCAGAGCC TGCAGCTGC TTTGAGTCTT ACGGAGATAG AGAATCAAT TTCTGCTAGG TAATTAACCA TTTCGACTCG AGCAGTGCCT
 CTTACTGATA ACCTTCTCGG ACGGTGACG ARACTCAGA TGCCTCTATC TCCTTAGTTA AAGACGATCC ATTAAATGGT AAGCTGAGC TCGTCAAGGT
 162 AsnAspTy rCysYsSerL euProAlaAl aLeuSerAla ThrGluIleG lYlySerIl eSerAlaArg OC*
 801 CTTTAAATAAT CTTTGTGTCAG AATAGATGAT GTGTGATATC TCCTTAGGAT GACTGTATTT TTCAAGTTGCC GATACAGCTT TTTGTCTCTT AACTGAGAA
 GAATTTTITA GAAACACATC TTATCTACTA CACAGCTTAG AGAATCTCTA CTGACATAAA AAGTCAACGG CTATGCTGAA AACACAGAGA TTGCACTCTT
 901 ACTCTTTATG TTAGATATAT TTCTCTAGGT TACTGTTGGG AGCTTAATGG TAGAACTTC CTGCTTTTCA TGAATAAGT CTTTTTTTTT CCTCA
 TGAGAAATAC AATCTATATA AAGAGATCCA ATGACAACCC TCGAATTACC ATCTTTGAG GAACCAAGT ACTAATTTCA GAAAAAAGAA GGAAT

FIG. 2

1 ATGATGACT CCACAGAAAG GAGCAGTCA GGCCTTACTT CTTCGCTTAA GAAAGAGAA GAAATGAAC TGAAGGATG TGTTCATC CTCCACGGA
 TACTACTGA GGTGCTTTC CCTCGTCAGT GCGAATGHA GAAAGGATT CTTTCTCTT CTTTACTTG ACTTCCTCAC ACAAGGTAG GAGGTGCGT
 1 M D D S T E R E Q S R L T S C L K K R E E M K L K E C V S I L P R K
 101 AGGAAGCCC CTTCTGCCA TCCTCCHAAG ACGGAAGCT GCTGGCTGCA ACCTGCTGC TGGCACTGCT GTCTTGCTGC CTCACGGTGG TGTCTTCTA
 TCCTTTCGGG GAGACAGCT AGGAGGTTC TCCCTTTCGA CGACCGAGCT TGGACGAGG ACGGTGAQA CAGAAGCAAG GAGTCCACAC ACGGAAGAT
 35 E S P S V R S S K D G K L L A A T L L L A L L S C C L T V V S F Y
 201 CCAGGTGGC GGCCTGCAAG GGGACCTGG CAGCCTCCG GCAGAGCTGC AGGGCACCA CGGGAGAG CTGCCAGCAG GACGAGGAGC CCCAGAGGC
 GGTCCACCG GGGGACGCT CCTGGACCG GTGGAGGEC CGTCTCGACG TCCGGTGGT GCGCTCTTC GAGGTGCTC CTGTCCTCG GGGGTTCGG
 68 Q V A A L Q G D L A S L R A E L Q G H H A E K L P A G A G A P K A
 301 GGCTTGAGG AAGCTCCAGC TGTACCGCG GACTGAA A TCTTGAAAC ACCAGTCCA GGAAGAGCA ACTCCAGTCA GAACAGCAGA AATAAGCGTG
 CCGAACCTCC TTCGAGGTG ACAGTGGCG CCTGACTTTT AGAAGCTTG TGGTCAGGT CCTCTTCGT TGAAGTCAGT CTGTGCTCT TTATTCGAC
 101 G L E E A P A V T A G L K I F E P P A P G E G N S S Q N S R N K R A
 401 CGTTTCAGG TCCAGAGAA ACAGTCATC AAGCTGCTT GCACTGAT GCAGACAGT GAACACCAAC TATACAAA GATCTTACA CATTTGTTCC
 GGCAAGTCC AGGCTTCTT TGTGATG TGTGACGAA CGTTGACTAA CGTCTCTAC TTGTGGTGG ATATGTTTT CTAGAATGT GTAACACAGG
 135 V Q G P E E T V T Q D C L Q L I A D S E T P T I Q K G S Y T F V P
 501 ATGGCTTCT AGCTTAAAG GGGAGAGTC CCTAGAGAA AAGAGATA AATATGTT GATGAACT CAAGAACT GTTACTTTT TTATATATGG TCAGSTTTA
 TACCGAGAG TCGAATTTT CCCCTTCAG GATCTTCTT TTCTCTTAT TTATTAACA GTTCTTTGA CCAATGAAGA AATATATACC AGTCCAAAT
 168 W L L S F K R G S A L E E K E N K I L V K E T G Y F F I Y G Q V L
 601 TATAGTATA AGACCTACG CATGGACAT CTAATCAGA GGAAGAGGT CCATGCTTT GGGATGAT TCACTCTGGT GACTTGTGT OGATGATTC
 ATATGACTAT TCTGGATCG GTACCTGTA GATTAAGTCT CCTCTTCCA GGTACAGAA CCCCTACTTA ACTCAGACCA CTGAACAAA GCTACATAG
 201 Y T D K T Y A M G H L I Q R K K V H V F G D E L S L V T L F R C I Q
 701 AAAATATGCC TGAACACTA CCAATATTT CCTGCTATTC AGCTGGCAT GCAATCTGG AAGAAGAGA TGAATCCAA CTTCATATAC CAAGAGAAA
 TTTATACGG ACTTGTGAT GGGTATTA GGAAGATAG TCGACCGTAA CGTTTGACC TTCTCTCTT ACTTGAGGT GAACGTATG GTTCTCTTT
 235 N M P E T L P N N S C Y S A G I A K L E E G D E L Q L A I P R E N
 801 TGCACAAATA TCACCTGATG GAGATGTCAC ATTTTGTG GCAATGAAC TGCCTGTA
 ACGTGTAT ATGACCTAC CTCACAGT GTCACAGT TAAACACCA CGTACTTG ACACACT
 268 A Q I S L D G D V T F F G A L K L L O

FIG. 3

1 GGTACGAGGC TTCCTAGAGG GACTGGAACC TAATCTCCT GAGGCTGAG GAGGTGGAG GGTCTCAGG CACGCTGGC CCACGACGG AGTGCCAGGA
 CCATGCTCCG AAGGATCTCC CTGACCTTGG ATTAAGAGGA CTCGACTCC CTCCACCTC CCAGAGTTC CCAGAGTCC GGTGCTGCC TCAOGGTCTC

 101 GCACCTAACAG TACCTTTAGC TTGCTTTCTT CTTCCCTCCT TTTTATTTT AAGTTCTTT TTATTCTTC TTGCTAACAC ACCTTCTTCC CTTCTGCACC
 CGTGATTGTC ATGGGATCG AACGAAGGA GAGGGGAGGA AAAATAAAG TTAAGGAAA AATAAAGAGG AACCATTTGT TGAAGAAGG GAAGACGTGG

 201 ACTGCCCGTA CCTTACCG CCGCGCCACC TCGTCTGAC CCACCTCTTG AACCCACAGC TGTGGCAGG GTCCCCAGCT CATGCCAGCC TCATCTCCTT
 TGACGGGCAT GGAATGGGC GGGCGGTGG AGGAACGATG GGTGAGAAC TTGGTGTGG ACAACGCTCC CAGGGTCCA M P A S S P F
 1

 301 TGTGCTAGC CCCAAAGG CCTCCAGGA ACATGGGGG CCTAGTCAGA GAGCGGCAC TCTAGTTGC CTTCTGGTTG AGTTGGGGG CAGCTCTGGG
 AGAACGATCG GGGGTTTCCC GGAGGTCCTG TGTACCCCCC GGTCTAGTCT CTGCGCGTG AGAGTCACG GGAHCCAC TCAACCCCCC GTCCAGACCC
 8 L L A P K G P P G N M G G P V R E P A L S V A L W L S W G A A L G

 401 GCGGTGGCT TGTGCCATGG CTCTGCTGAC CCAACAAACA GAGCTCAGA GCTCAGGAG AGAGTGAAG CCGCTGCAGG GACAGAGG CCGCTCCACG
 CCGCACCGA ACACGGTACC GAGACGACTG GGTGTTTGT CTGAGCTCT CCGAGTCTC TCTCCTCTG GCGACCTCC CTTGCTCTCC GGGAGGCTC
 41 A V A C A M A L L T Q Q T E L Q S L R R E V S R L Q G T G G P S Q

 501 AATGGGAAG GGTATCCCTG GCGAGTCTC CCGAGCAGA GTTCCGATGC CTGGAAGCC TGGAGAATG GGGAGAGATC CCGAAAGG AGAGCAGTGC
 TTACCCCTTC CCATAGGAG CGTCTCAGAG GGCCTGCTCT CAAGGCTACG GACCTTGG ACCCTCTAC CCGCTCTCAG GGCCTTTTCC TCTCGTCAAG
 74 N G E G Y P W Q S L P E Q S S D A L E A W E N G E R S R K R R A V L

 601 TCACCCAAA ACAGAAGAAG CAGCAGTCTG TCGTCCACTT GGTCCACTT AACGCCACT CCAAGGAGA CTCGATGTG ACAGAGGTGA TGTGCGAACC
 AGTGGGTTT TGTCTTCTTC GTGCTGAGAC AGGACGTGGA CCAAGGGTAA TTGCGGTGA GGTTCCTACT GAGGCTACAC TGTCTCCACT ACACGCTTGG
 108 T Q K Q K K Q H S V L H L V P I N A T S K D D S D V T E V M W Q P

 701 AGCTCTTAGG CGTGGAGAG GCTACAGGC CCAAGGATAT GGTGTCCGA TCCAGGATG TGAAGTTAT CTGCTGTATA GCGAGTCTT GTTTCAGAC
 TCGAGAAACC GCACCTCTC CGGATGTCG GGTCTCTATA CCACAGGCTT AGGTCTTACG ACCTCAATA GACGACATAT CGGTCCAGG CAAAGTTCTG
 141 A L R R G R G L Q A Q G Y G V R I Q D A G V Y L L Y S Q V L F Q D

 801 GTGACTTTCA CCATGGGTCA GGTGGTGTCT CCGAAGGCC AAGGAAGCA GGAGACTCTA TTCCGATGA TAAGAATAT GCCTCCAC CCGGACCGG
 CACTGAAGT GGTACCCAGT CCACACAGA GTCTTCCG GTCTTCCG TCTCTGAT AGGCTACAT ATTCTTCATA CGGAGGCTG GGCCTGGCC
 174 V T F T M G Q V V S R E G Q G R Q E T L F R C I R S M P S H P D R F

 901 CCTACACAG CTGCTATAGC GCAGGTGTCT TCCATTACA CCAAGGGAT ATTCTGATG TCATAATTC CCGGGCAGG GCGAACTTA ACCTCTCTC
 GGATGTGTC GACGATATCG CGTCCACAGA AGGTAAATGT GGTCCCTTA TAAGACTCAC AGTATTAGG GGGCGTTC CGCTTCTAT TGGAGAGG
 208 Y N S C Y S A G V F H L H Q G D I L S V I I P R A R A K L N L S P

FIG. 4A

1001 ACATGGAACC TTCTGGGGT TTGTGAACT GTGATTGTGT TATAAAGT GGCTCCAGC TTGGAAGACC AGGTTGGGTA CATACTGGAG ACAGCCAAGA
TGTACCTTGG AAGGACCCCA AACACTTTGA CACTAACACA ATATTTTCA CCGAGGTCG AACCTTCTGG TCCACCCAT GTATGACCTC TGTGGTTCT
241 H G T F L G F V K L O
1101 GCTGAGTATA TAAAGGAGAG GGAATGTGCA GGAACAGAGG CATCTTCTG GATTGGCTC CCGTTCTC ACTTTTCCCT TTTCATTCCC ACCCCCTAGA
CGACTCATAT ATTTCTCTC CTTACACGT CCTGTCTCC GTAGAAGGAC CCAACCGAG GGCRAAGGAG TGAAGAAGGA AAGTAAGG TGGGGGATCT
1201 CTTTGATTTT ACGGATATCT TGTCTCTGTT CCCCATGGAG CTCGGAATTC TTGGTGTGT GTAGATGAGG GGGGGGGGAC GGGGGCCAGG CATTTGTTAG
GAACTAAA TGCCTATAGA ACGAAGACAA GGGGTACCTC GAGGCTAAG AACGCACACA CATCTACTCC CCGCCCCCTG CCGCGGTCC GTACAGTC
1301 ACCTGGTCGG GGGCCACTGG AAGCATCCAG AACAGCACCA CCATCTTA
TGGACCAACC CCGGTGACC TTCGTAGGTC TTGTCGTGGT GGTAGAT

FIG. 4B

TACIs

agcatcctgagtaATGAGTGGCCTGGGCCGGAGCAGGCGAGGTGGCCGGAGCCGTGTGGACCAGG
 AGGAGCGCTGGTCACTCAGCTGCCGCAAGGAGCAAGGCAAGTTCTATGACCATCTCCTGAGGGAC
 TGCATCAGCTGTGCCTCCATCTGTGGACAGCACCCCTAAGCAATGTGCATACTTCTGTGAGAACA
 GCTCAGGAGCCCAGTGAACCTTCCACCAGAGCTCAGGAGACAGCGGAGTGGAGAAGTTGAAAACA
 ATTCAGACAACCTCGGGAAGGTACCAAGGATTTGGAGCACAGAGGCTCAGAAGCAAGTCCAGCTCTC
 CCGGGGCTGAAGCTGAGTGCAGATCAGGTGGCCCTGGTCTACAGCACGCTGGGGCTCTGCCTGTG
 TGCCGTCTCTGTGCTTCTGGTGGCGGTGGCCTGCTTCTCAAGAAGAGGGGGGATCCCTGCT
 CCTGCCAGCCCCGCTCAAGGCCCGTCAAAGTCCGGCCAAGTCTTCCAGGATCACCGGATGGAA
 GCGGAGAGCCCTGTGAGCACATCCCCGAGCCAGTGGAGACCTGCAGCTTCTGCTTCCCTGAGTG
 CAGGGCGCCACGCAGGAGAGCGCAGTCACGCCTGGGACCCCCGACCCCACTTGTGCTGGAAGGT
 GGGGGTGGCACACCAGGACCACAGTCTGCAGCCTTGCCACACATCCAGACAGTGGCCTTGGC
 ATTGTGTGTGTGCTGCCAGGAGGGGGGCCAGGTGCATAAatgggggtcagggagggaaagga
 ggagggagagagatggagaggagggagagagaaagagaggtggggagaggggagagagataatga
 ggagagagagacagaggaggcagaaaggagagaaacagaggagacagagaggagagagagaca
 gagggagagagagacagagggaagagaggcagagagggaaagaggcagagaaggaaagagacag
 gcagagaaggagagaggcagagaggagagaggcagagagggagagaggcagagagacagagagg
 gagagagggacagagagagatagagcaggaggtcggggcactotgagtccagttccagtgag
 ctgtaggtcgtcatcacctaaccacagtgcaataaagtcctcgtgctgctcacagcccc
 gagagccccctcctcctggagaataaaacctttggcagctgcccctcctcaaaaaaaaaaaaaaa
 aaaa

FIG._5A**TACIs :**

MSGLGRSRRGGRSRVDQERWLSRKEQGFYDHLRLDCISCASI¹CGQHPKQCA²YFCENKLRSP
 VNLPELRRQRSGEVENNSDNSGRYQGLEHRGSEASPALPGLKLSADQVALVYSTLGLCLCAVL
 CFLVAVACFLKRGDPCSCQPRSRPQSPAKSSQDHAMEAGSPVSTSPFPVETCSFCFPECRAFT
 QESAVTPGTPDPTCAGRNGCHTRTTVLQPCPHIPDSGLGIVCVPAQEGGPG

FIG._5B**human BR3:**

cgtcggcaccATGAGGCGAGGGCCCCGGAGCCTGCGGGGCGAGGACGCGCCAGCCCCAGGCCCT
 GCGTCCCGGCGAGTGTCTCGACCTGCTGGTCCGCCACTGCGTGGCCTGCGGGCTCCTGCGCACG
 CCGCGGCGGAAACCGGCGGGGGCCAGCAGCCCTGCGCCCGAGGACGGCGCTGCAGCCGAGGAGTC
 GGTGGGCGCGGGGGCCGGCGAGGCGCGCTGCCCTGCGCGGGCTGCTCTTTGGCGCCCCGCGC
 TGCTGGGCTGGCACTGGTCTGCGCTGGTCTGCGTGGGTCTGGTGGAGCTGGAGGCGGCGACAG
 CGGCGGCTTGGCGGCGCTCTCCGAGAGGCCCCGACGGAGACAAGGACGCCCCAGAGCCCCCT
 GGACAAGGTATCATTTCTGTCTCCGGGAATCTCTGATGCCACAGCTCCTGCCTGGCCTCCTCTG
 GGAAGACCCAGGAACCAACCCACCTGGCCACAGTGTCCCTGTGCCAGCCACAGAGCTGGGCTCC
 ACTGAACTGGTGACCACCAAGACGGCCGGCCCTGAGCAACAATAGcagggagccggcaggaggtg
 gccccctgccc

FIG._6A**human BR3:**

MRRGPRSLRGRDAPAPTPCVPAECFDLLVRHCVACGLLRTPRPKPAGASSPAPRTALQFQESVGA
 GAGEAALPLPGLLFGAPALLGLALVLALVLVGLVSWRRRQRRLRGASSAEAPDGDKDAPEFLDKV
 IILSPGISDATAPAWPPPGEDPGTTPGHSVPVPATELGSTELVTTKTAGPEQQ

FIG._6B

murine BR3

ATGGGCGCCAGGAGACTCCGGGTCCGAAGCCAGAGGAGCCGGGACAGCTCGGTGCCACCCAGTG
CAATCAGACCCGAGTGCTTCGACCCTCTGGTGAGAACTGCGTGTCTGTGAGCTCTTCACACGC
CGGACACTGGACATACAAGCAGCCTGGAGCCTGGGACAGCTCTGCAGCCTCAGGAGGGCTCCGCG
CTGAGACCCGACGTGGCGCTGCTCGTGGTGCCCCCGCACTCCTGGGACTGATACTGGCGCTGAC
CCTGGTGGGTCTAGTGAGTCTGGTGAGCTGGAGGTGGCGTCAACAGCTCAGGACGGCTCCCCAG
ACACTTCAGAAGGAGTCCAGCAAGAGTCCCTGGAAAATGTCTTTGTACCTCCTCAGAAACCCCT
CATGCCTCAGCTCCTACCTGGCCTCCGCTCAAAGAAGATGCAGACAGCGCCCTGCCAGCCACAG
CGTCCCGGTGCCCGCCACAGAAGTGGGCTCCACCGAGCTGGTGACCACCAAGACAGCTGGCCAG
AGCAATAGcagcagtgagggtggaacccaggatctctactgggcttgtggacttcacccaaca
gcttgggaaagaacttggcccttcagtgaaggagtcctttgcctggggggcgaacccggcagaac
cagacactacaggocacatgagattgcttttgtgttagctcttgacttgagaacgttccatttct
gagatgggttttaagcctgtgtgccttcagatggttggatagacttgagggttgcatatttaac
tctgtagtgaagcggagactggaacttaactctcggttctaaaaattttggattactgggctggag
gtatggctcagcagttcggtttgtgtgctgttctagccgaggactccagttgttcagcttcccg
aactcagatctggcagcttaagaccacctgtcactccagccctggaacatccttgctccaaag
gcaccagcactgatttgccttagagca
cacatatgcatgcatgcacacttaaaaatgtcaaaattagcggctggagaaattcatggtcaaca
gcgcttactgtgattccagaggatgagagtttgattcccagaatgcactgcgggtggctcattac
tgagcataacttttgccttcaggggacctgatgcctctggacttcatgggcatctgtattcacgtg
cacatectactcttttacaatg
ataaaatataagatgggcatggtggtacacacotttaatcccaacattggggaagcaaaggcagg
caggtaaatgagttggaggccatcctggtctacatagcaagttccaggctaaccagagctaaatg
gtgagaccaagtctcaaaaataatactcccccccccaaaaaaaaacttttaaattttgattttt
tcttttattattttttttatattaatttcatggtgtttagaagtggatatacttagatgggtgact
aagaggaggtaaagccatcaggactgagcccctaacaatacaaggagaaagcagagacaatgaaca
cgccctctcctgctgtgtgcccagctctggaccaccagccagagggcaatcatcagatgtgggccc
ctagaaccttcagagccgaaagctaaatcaatctcatttctttgtaaagctatttagccttaggt
gttttggttacggtgatataaaatggactaacacaggcactatgagtaagaagcttttctttgagc
tgggaaagggtactgtttaaaccaaaaattaatctgaataaaaaaaggctaaggggaagacactt

FIG. 6C

PRO XXXXXXXXXXXXXXXX (Length = 15 amino acids)

Comparison Protein XXXXXYYYYYYY (Length = 12 amino acids)

% amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

5 divided by 15 = 33.3%

FIG. 7A

PRO XXXXXXXXXX (Length = 10 amino acids)

Comparison Protein XXXXXYYYYYYZZY (Length = 15 amino acids)

% amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

5 divided by 10 = 50%

FIG. 7B

	10	20	30	40
hBR3	M---	RRGPRSLRGRDAPAPT	PCVPAECFDLLVRHCVACGL	
mBR3	MGARRLRVRSQSRDSSVPTQCNQTECFDPLVRNCVSCEL			
	M...R...RS.R.RD+...PT.C...+ECFD.LVR+CV+C.L			

	50	60	70	80
hBR3	LRTPRPKFAGASSPAPRTALQPQESVGAGAGEAALPLPLGL			
mBR3	FHTP--DTGHTSSLEPGTALQPQE-----GSALRPDVAL			
	..TP.....+SS..P.TALQPQE.....G.A..P...L			

	90	100	110	120
hBR3	LFGAPALLGLALVLALV.LVGLVSWRRRQRRLRGASSAEA			
mBR3	LVGAPALLGLLILALTIVGLVSLVSWRWRRQQ-LRTASP-DT			
	L.GAPALLGL.L.L+LV.LV.LVSWR.RQ+.LR.AS...+			

	130	140	150	160
hBR3	PDGDKDAPEPLDKVIILSPGISDATAPAWPPPGEDPGTTP			
mBR3	SEGVQQ--ESLENVFPSSSETPHASAPTWPPLKEDADSAL			
	..+G...E.L+.V...S.....A+AP.WPP..ND...++.			

	170	180
hBR3	PGHSVPVPATELGSTELVTTKTAGPEQQ	
mBR3	PRHSVPVPATELGSTELVTTKTAGPEQ	
	P.HSVPVPATELGSTELVTTKTAGPEQ.	

FIG. 9

Sequence of human CD20 Showing Predicted Transmembrane (boxed) and Extracellular (underlined) Regions

CD20.hu	10	20	30	40	50
	MTTPRNSVNGTFFPAEPMKGP	IA MQSGPKPLFRMSSLVGPTQ	SFFMR	ESK	
CD20.hu	60	70	80	90	100
	TLGAVQIMNGLFHLALGGLLMIPAGIYAPICVT	WYPLWGGIMYII	SGSL		
CD20.hu	110	120	130	140	150
	LAATEKNSRKCLVKGMIMN	SLSLFAISGMILSIMDILN	KISHFLKME		
CD20.hu	160	170	180	190	200
	SLNFI RAHTPYININCEPAN	SEKNSPSTOYCYSIOS	FLGILSVMLIF		
CD20.hu	210	220	230	240	250
	AFFOELVIA	GIVENEWKRTCSRPKSNIVLLSAEKK	QETBIKEEVVGLT		
CD20.hu	260	270	280	290	297
	ETSSQPKNEEDIEIPIQEEEEETETNFPEPPQ	QESSPLENDSSP			

FIG. 10

Figure 11

Human CD20 cDNA sequence

ATGACAACACCCAGAAATTCAGTAAATGGGACTTTCCCGGCAGAGCCAATGAAAGGCCCTAT
TGCTATGCAATCTGGTCCAAAACCACTCTTCAGGAGGATGTCTTCACTGGTGGGCCCCACG
CAAAGCTTCTTCATGAGGGAATCTAAGACTTTGGGGGCTGTCCAGATTATGAATGGGCTCTT
CCACATTGCCCTGGGGGGTCTTCTGATGATCCAGCAGGGATCTATGCACCCATCTGTGTG
ACTGTGTGGTACCCCTCTCTGGGGAGGCATTATGTATATTATTTCCGGATCACTCCTGGCAGC
AACGGAGAAAACTCCAGGAAGTGTTTGGTCAAAGGAAAAATGATAATGAATTCATTGAGCC
TCTTTGCTGCCATTTCTGGAATGATTCTTTCAATCATGGACATACTTAATATTAATTTCCCA
TTTTTAAAAATGGAGAGTCTGAATTTTATTAGAGCTCACACACCATATATTAACATATACAAC
TGTGAACCAGCTAATCCCTCTGAGAAAACTCCCATCTACCCAATACTGTTACAGCATACAA
TCTCTGTTCTTGGGCATTTTGTGAGTGATGCTGATCTTGCCTTCTTCCAGGAACCTTGTAATA
GCTGGCATCGTTGAGAATGAATGGAAAAGAACGTGCTCCAGACCCAAATCTAACATAGTTCT
CCTGTCAGCAGAAGAAAAAAGAACAGACTATTGAAATAAAGAAAGAGTGGTTGGGCTAA
CTGAAACATCTTCCCAACCAAAGAATGAAGAAGACATTGAAATTATTCCAATCCAAGAAGAGG
AAGAAGAAGAAACAGAGACGAACCTTCCAGAACCTCCCAAGATCAGGAATCCTCACCAATA
GAAAATGACAGCTCTCCT

FIG. 12

Sequence alignment of variable light-chain domain

	10	20	30	40
2H7	QIVLSQSPAIL	SASPGEKVTITC	[RASSSVS-YMH]	WYQOKP
	* * * *	* * *	* * *	*
hu2H7.v16	DIQMTQSPSSL	SASVGDRVTITC	[RASSSVS-YMH]	WYQOKP
			* * * *	
hum KI	DIQMTQSPSSL	SASVGDRVTITC	[RASQISNYLA]	WYQOKP
	50	60	70	80
2H7	GSSPKPWIY	[APSNLAS]	GVPARFSGSGSGT	SYSLTISRVEA
	** *	*	***	****
hu2H7.v16	GKAPKPLIY	[APSNLAS]	GVPSRFSGSGSGT	DFTLTISLQP
	*	* * *		
hum KI	GKAPKLLIY	[AASSLES]	GVPSRFSGSGSGT	DFTLTISLQP
	90	100		
2H7	EDAATYYC	[QQWSENPPT]	FGAGTKLELKR	
	*		* * *	
hu2H7.v16	EDFATYYC	[QQWSENPPT]	FGQGTKVEIKR	
		**** *		
hum KI	EDFATYYC	[QQYNSLPWT]	FGQGTKVEIKR	

FIG. 13
Sequence alignment of variable heavy-chain domain

	10	20	30	40
2H7	QAYLQQSGAELVRPGASVKMSCKAS [GYTFTSYNMH] WVRQT			
	*** ** * * * * *			
hu2H7.v16	EVQLVESGGGLVQPGGSLRLSCAAS [GYTFTSYNMH] WVRQA			
	* * * *			
hum III	EVQLVESGGGLVQPGGSLRLSCAAS [GFTFSSYAMS] WVRQA			

	50	a	60	70	80
2H7	PRQGLEWIG [AIYPGNGDTSYNQKFKG] KATLTVDKSSSTAYM				
	** * * * * *				
hu2H7.v16	PGKGLEWVG [AIYPGNGDTSYNQKFKG] RFTISVDKSKNTLYL				
	* * * * * * * *				
hum III	PGKGLEWVA [VISGDGGSTYYADSVKG] RFTISRDNKNTLYL				

	abc	90	100abcde	110
2H7	QLSSLTSEDSAVYFCAR [VYYYSNSYWFYFDV] WGTGTLTVTVSS			
	** * * *			
hu2H7.v16	QMNSLRAEDTAVYYCAR [VYYYSNSYWFYFDV] WQGTLTVTVSS			
	***** * * *			
hum III	QMNSLRAEDTAVYYCAR [GRVGYSLY---DY] WQGTLTVTVSS			

Human CD20 Transgene expression in mouse B220⁺ cells




FIG. 14




FIG. 15

Expression of human CD20 in Tg⁺ mouse bone marrow





FIG. 16

Expression of human CD20 in Tg⁺ mouse splenic B cells




FIG. 17

Expression of human CD20 in Tg⁺ mesenteric LNs




FIG. 18

Expression of human CD20 in Tg⁺ Peyer's Patches




FIG. 19

Effects of anti-CD20 mAb in mice




FIG. 20 Serum levels of anti-CD20 mAb

Depletion of peripheral B cells with anti-CD20 mAbs




FIG. 21

Depletion of mature peripheral LN B cells by anti-CD20 mAb




FIG. 22

Depletion of splenic T2 B cells by anti-CD20 mAbs




FIG. 23

Depletion of re-circulating mature B cells by anti-CD20 mAbs




FIG. 24

Resistance of Peyer's Patches Germinal Center B cells to anti-CD20 mAbs




FIG. 25

Depletion & Recovery of B cells following anti-CD20 mAb Rx




FIG. 26

Resistance of Splenic Germinal Center B cells to short-term anti-CD20 mAb Rx




FIG. 27

Non-depleted MZ and B1 B cells confer protection
to T-Independent antigens




FIG. 28

Distinct biologic effects of BR3 and anti-CD20 mAb treatment




FIG. 29




FIG. 30




FIG. 31

SEQ ID NO:	SEQUENCE	CLONE NUMBER
(SEQ ID NO: 1)	E C F D L L V R Q W V P C C E R I B G	1
(SEQ ID NO: 2)	E C F D L L V R R B W W V P C C E R I B G	2, 6, 11, 14, 24, 28, 34, 36, 40, 42, 46, 47
(SEQ ID NO: 3)	E C F D L L V R R B W W V P C C E R I B G	3
(SEQ ID NO: 4)	E C F D L L V R R B W W V P C C E R I B G	4
(SEQ ID NO: 5)	E C F D L L V R R B W W V P C C E R I B G	5
(SEQ ID NO: 6)	E C F D L L V R R B W W V P C C E R I B G	7, 9, 10, 20, 25, 29
(SEQ ID NO: 7)	E C F D L L V R R B W W V P C C E R I B G	8
(SEQ ID NO: 8)	E C F D L L V R R B W W V P C C E R I B G	12
(SEQ ID NO: 9)	E C F D L L V R R B W W V P C C E R I B G	13
(SEQ ID NO: 10)	E C F D L L V R R B W W V P C C E R I B G	15
(SEQ ID NO: 11)	E C F D L L V R R B W W V P C C E R I B G	16
(SEQ ID NO: 12)	E C F D L L V R R B W W V P C C E R I B G	17
(SEQ ID NO: 13)	E C F D L L V R R B W W V P C C E R I B G	18
(SEQ ID NO: 14)	E C F D L L V R R B W W V P C C E R I B G	19
(SEQ ID NO: 15)	E C F D L L V R R B W W V P C C E R I B G	21
(SEQ ID NO: 16)	E C F D L L V R R B W W V P C C E R I B G	22
(SEQ ID NO: 17)	E C F D L L V R R B W W V P C C E R I B G	23
(SEQ ID NO: 18)	E C F D L L V R R B W W V P C C E R I B G	26
(SEQ ID NO: 19)	E C F D L L V R R B W W V P C C E R I B G	27
(SEQ ID NO: 20)	E C F D L L V R R B W W V P C C E R I B G	30
(SEQ ID NO: 21)	E C F D L L V R R B W W V P C C E R I B G	31
(SEQ ID NO: 22)	E C F D L L V R R B W W V P C C E R I B G	32
(SEQ ID NO: 23)	E C F D L L V R R B W W V P C C E R I B G	33
(SEQ ID NO: 24)	E C F D L L V R R B W W V P C C E R I B G	35
(SEQ ID NO: 25)	E C F D L L V R R B W W V P C C E R I B G	37
(SEQ ID NO: 26)	E C F D L L V R R B W W V P C C E R I B G	38

FIG. 32A

SEQ ID NO:	SEQUENCE	CLONE NUMBER
(SEQ ID NO:)	S H Q K S L H I N D R I A H D T H N N R R G V I I H I	39
(SEQ ID NO:)	L F L L L L L L L I L L I L L L L I L L L L L I I L I I L	41
(SEQ ID NO:)	L T V V L L I A L I A L R L L V A L R L L V L A L L I	43
(SEQ ID NO:)	I S S S E S A A H S S R D I H S V A A N N R N I H G D	44
(SEQ ID NO:)	C C	45
(SEQ ID NO:)	P P P P S D P P P D P D A P P P D D D P P P P P P P R	48
(SEQ ID NO:)	V V	49
(SEQ ID NO:)	W W	50
(SEQ ID NO:)	S K G A Q G R H B V A R I R A H R L A L H A R A R R D	52, 71
(SEQ ID NO:)	R R R R R R D G A R N N R G R G N N S B R N N R R G R	53
(SEQ ID NO:)	V V	54
(SEQ ID NO:)	L L	55
(SEQ ID NO:)	L L L L L L I R P P S L P P L P P R V S P I R R L I P	56
(SEQ ID NO:)	D D	57
(SEQ ID NO:)	F F	58
(SEQ ID NO:)	C C	59
(SEQ ID NO:)	E E	60
(SEQ ID NO:)	(SEQ ID NO:)	61
(SEQ ID NO:)	(SEQ ID NO:)	62
(SEQ ID NO:)	(SEQ ID NO:)	63
(SEQ ID NO:)	(SEQ ID NO:)	64
(SEQ ID NO:)	(SEQ ID NO:)	65
(SEQ ID NO:)	(SEQ ID NO:)	66
(SEQ ID NO:)	(SEQ ID NO:)	67
(SEQ ID NO:)	(SEQ ID NO:)	68
(SEQ ID NO:)	(SEQ ID NO:)	69
(SEQ ID NO:)	(SEQ ID NO:)	70

GAGSP

FIG. 32B

CLONE NUMBER

SEQUENCE

SEQ ID NO:

72	R	S	R	A	D	H	A	I	R	G	R	R	D	I	I	I	N	H	H	R	H	A	I	D	I
73	L	L	I	L	L	L	L	I	L	L	L	L	L	L	L	L	L	L	L	L	I	I	L	L	
74	V	V	L	A	L	V	L	A	L	V	V	L	A	L	A	A	L	V	A	L	L	V	H	L	V
75	I	G	G	I	N	A	R	G	S	A	A	G	R	D	R	G	A	A	S	D	D	S	S	G	H
76	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
77	P	P	P	P	P	P	P	D	P	D	D	D	D	P	P	P	P	P	P	P	P	P	P	P	
78	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	
79	W	W	W	W	W	W	W	W	W	W	W	W	W	W	W	W	W	W	W	W	W	W	W	W	
80	V	A	R	I	A	G	A	D	R	B	A	A	H	H	A	R	N	A	R	B	D	S	A	I	A
81	R	R	H	R	N	N	N	N	R	H	N	N	R	R	R	N	R	N	L	H	R	N	R	B	N
82	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	V	
83	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	
84	P	S	V	H	I	R	B	L	A	A	V	G	P	D	V	I	D	P	P	A	P	L	I	S	S
85	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	
86	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	
87	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
88	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	E	
89))))))))))))))))))))))))	
90	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	
91	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	
92	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	
93	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	
94	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	
95	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	
96	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	(SEQ ID NO:)	

FIG.32C




FIG. 33

Sequence Listing

<110> Genentech, Inc.

<120> COMBINATION THERAPY FOR B CELL DISORDERS

<130> 11669.258WO11

<140> New Filing

<141> 2005-12-12

<150> US 11/021,874

<151> 2004-12-22

<150> US 10/861,049

<151> 2004-06-04

<150> US 60/476,531

<151> 2003-06-06

<150> US 60/476,481

<151> 2003-06-05

<150> US 60/476,414

<151> 2003-06-05

<160> 165

<210> 1

<211> 314

<212> PRT

<213> Mus musculus

<400> 1

Met	Ser	Ala	Leu	Leu	Ile	Leu	Ala	Leu	Val	Gly	Ala	Ala	Val	Ala
1				5						10				15

Ser	Thr	Gly	Ala	Arg	Arg	Leu	Arg	Val	Arg	Ser	Gln	Arg	Ser	Arg
				20					25					30

Asp	Ser	Ser	Val	Pro	Thr	Gln	Cys	Asn	Gln	Thr	Glu	Cys	Phe	Asp
				35					40					45

Pro	Leu	Val	Arg	Asn	Cys	Val	Ser	Cys	Glu	Leu	Phe	His	Thr	Pro
				50					55					60

Asp	Thr	Gly	His	Thr	Ser	Ser	Leu	Glu	Pro	Gly	Thr	Ala	Leu	Gln
				65					70					75

Pro	Gln	Glu	Gly	Gln	Val	Thr	Gly	Asp	Lys	Lys	Ile	Val	Pro	Arg
				80					85					90

Asp	Cys	Gly	Cys	Lys	Pro	Cys	Ile	Cys	Thr	Val	Pro	Glu	Val	Ser
				95					100					105

PCT/US05/44926

Ser Val Phe Ile Phe Pro Pro Lys Pro Lys Asp Val Leu Thr Ile
 110 115 120
 Thr Leu Thr Pro Lys Val Thr Cys Val Val Val Asp Ile Ser Lys
 125 130 135
 Asp Asp Pro Glu Val Gln Phe Ser Trp Phe Val Asp Asp Val Glu
 140 145 150
 Val His Thr Ala Gln Thr Gln Pro Arg Glu Glu Gln Phe Asn Ser
 155 160 165
 Thr Phe Arg Ser Val Ser Glu Leu Pro Ile Met His Gln Asp Trp
 170 175 180
 Leu Asn Gly Lys Glu Phe Lys Cys Arg Val Asn Ser Ala Ala Phe
 185 190 195
 Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Arg Pro
 200 205 210
 Lys Ala Pro Gln Val Tyr Thr Ile Pro Pro Pro Lys Glu Gln Met
 215 220 225
 Ala Lys Asp Lys Val Ser Leu Thr Cys Met Ile Thr Asp Phe Phe
 230 235 240
 Pro Glu Asp Ile Thr Val Glu Trp Gln Trp Asn Gly Gln Pro Ala
 245 250 255
 Glu Asn Tyr Lys Asn Thr Gln Pro Ile Met Asn Thr Asn Gly Ser
 260 265 270
 Tyr Phe Val Tyr Ser Lys Leu Asn Val Gln Lys Ser Asn Trp Glu
 275 280 285
 Ala Gly Asn Thr Phe Thr Cys Ser Val Leu His Glu Gly Leu His
 290 295 300
 Asn His His Thr Glu Lys Ser Leu Ser His Ser Pro Gly Lys
 305 310

<210> 2

<211> 311

<212> PRT

<213> Homo sapiens

<400> 2

Met Ser Ala Leu Leu Ile Leu Ala Leu Val Gly Ala Ala Val Ala
 1 5 10 15
 Ser Thr Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro
 20 25 30
 Ala Pro Thr Pro Cys Val Pro Ala Glu Cys Phe Asp Leu Leu Val

					35						40						45
Arg	His	Cys	Val	Ala	Cys	Gly	Leu	Leu	Arg	Thr	Pro	Arg	Pro	Lys			
				50					55					60			
Pro	Ala	Gly	Ala	Ser	Ser	Pro	Ala	Pro	Arg	Thr	Ala	Leu	Gln	Pro			
				65					70					75			
Gln	Glu	Ser	Gln	Val	Thr	Asp	Lys	Ala	Ala	His	Tyr	Thr	Leu	Cys			
				80					85					90			
Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe			
				95					100					105			
Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr			
				110					115					120			
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Ala	Val	Ser	His	Glu	Asp	Pro			
				125					130					135			
Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn			
				140					145					150			
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg			
				155					160					165			
Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly			
				170					175					180			
Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro			
				185					190					195			
Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro			
				200					205					210			
Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn			
				215					220					225			
Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp			
				230					235					240			
Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr			
				245					250					255			
Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu			
				260					265					270			
Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn			
				275					280					285			
Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr			
				290					295					300			
Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys							
				305					310								

<210> 3
 <211> 232
 <212> PRT
 <213> Artificial sequence

<220>
 <223> sequence is synthesized

<400> 3
 Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr
 1 5 10 15
 Gly Val His Ser Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu
 20 25 30
 Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
 35 40 45
 Ser Ser Val Ser Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Lys
 50 55 60
 Ala Pro Lys Pro Leu Ile Tyr Ala Pro Ser Asn Leu Ala Ser Gly
 65 70 75
 Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 80 85 90
 Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr
 95 100 105
 Cys Gln Gln Trp Ser Phe Asn Pro Pro Thr Phe Gly Gln Gly Thr
 110 115 120
 Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile
 125 130 135
 Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val
 140 145 150
 Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln
 155 160 165
 Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser
 170 175 180
 Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
 185 190 195
 Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 200 205 210
 Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys
 215 220 225

Ser Phe Asn Arg Gly Glu Cys
230

<210> 4

<211> 471

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 4

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr
1 5 10 15

Gly Val His Ser Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu
20 25 30

Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly
35 40 45

Tyr Thr Phe Thr Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro
50 55 60

Gly Lys Gly Leu Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly
65 70 75

Asp Thr Ser Tyr Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser
80 85 90

Val Asp Lys Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu
95 100 105

Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr
110 115 120

Tyr Ser Asn Ser Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr
125 130 135

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
140 145 150

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
155 160 165

Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
170 175 180

Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
185 190 195

Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val
200 205 210

Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn

215	220	225
Val Asn His Lys Pro Ser Asn Thr Lys	Val Asp Lys Lys Val	Glu
230	235	240
Pro Lys Ser Cys Asp Lys Thr His Thr	Cys Pro Pro Cys Pro	Ala
245	250	255
Pro Glu Leu Leu Gly Gly Pro Ser Val	Phe Leu Phe Pro Pro	Lys
260	265	270
Pro Lys Asp Thr Leu Met Ile Ser Arg	Thr Pro Glu Val Thr	Cys
275	280	285
Val Val Val Asp Val Ser His Glu Asp	Pro Glu Val Lys Phe	Asn
290	295	300
Trp Tyr Val Asp Gly Val Glu Val His	Asn Ala Lys Thr Lys	Pro
305	310	315
Arg Glu Glu Gln Tyr Asn Ser Thr Tyr	Arg Val Val Ser Val	Leu
320	325	330
Thr Val Leu His Gln Asp Trp Leu Asn	Gly Lys Glu Tyr Lys	Cys
335	340	345
Lys Val Ser Asn Lys Ala Leu Pro Ala	Pro Ile Glu Lys Thr	Ile
350	355	360
Ser Lys Ala Lys Gly Gln Pro Arg Glu	Pro Gln Val Tyr Thr	Leu
365	370	375
Pro Pro Ser Arg Glu Glu Met Thr Lys	Asn Gln Val Ser Leu	Thr
380	385	390
Cys Leu Val Lys Gly Phe Tyr Pro Ser	Asp Ile Ala Val Glu	Trp
395	400	405
Glu Ser Asn Gly Gln Pro Glu Asn Asn	Tyr Lys Thr Thr Pro	Pro
410	415	420
Val Leu Asp Ser Asp Gly Ser Phe Phe	Leu Tyr Ser Lys Leu	Thr
425	430	435
Val Asp Lys Ser Arg Trp Gln Gln Gly	Asn Val Phe Ser Cys	Ser
440	445	450
Val Met His Glu Ala Leu His Asn His	Tyr Thr Gln Lys Ser	Leu
455	460	465
Ser Leu Ser Pro Gly Lys		
470		

<210> 5
 <211> 17

<212> PRT
<213> Artificial sequence

<220>
<223> sequence is synthesized

<400> 5
Glu Cys Phe Asp Leu Leu Val Arg Ala Trp Val Pro Cys Ser Val
1 5 10 15

Leu Lys

<210> 6
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> sequence is synthesized

<400> 6
Glu Cys Phe Asp Leu Leu Val Arg His Trp Val Pro Cys Gly Leu
1 5 10 15

Leu Arg

<210> 7
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> sequence is synthesized

<400> 7
Glu Cys Phe Asp Leu Leu Val Arg Arg Trp Val Pro Cys Glu Met
1 5 10 15

Leu Gly

<210> 8
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> sequence is synthesized

<400> 8
Glu Cys Phe Asp Leu Leu Val Arg Ser Trp Val Pro Cys His Met
1 5 10 15

Leu Arg

<210> 9
 <211> 17
 <212> PRT
 <213> Artificial sequence

<220>
 <223> sequence is synthesized

<400> 9
 Glu Cys Phe Asp Leu Leu Val Arg His Trp Val Ala Cys Gly Leu
 1 5 10 15

Leu Arg

<210> 10
 <211> 291
 <212> PRT
 <213> Artificial sequence

<220>
 <223> sequence is synthesized

<400> 10
 Met Leu Pro Gly Cys Lys Trp Asp Leu Leu Ile Lys Gln Trp Val
 1 5 10 15
 Cys Asp Pro Leu Gly Ser Gly Ser Ala Thr Gly Gly Ser Gly Ser
 20 25 30
 Thr Ala Ser Ser Gly Ser Gly Ser Ala Thr His Met Leu Pro Gly
 35 40 45
 Cys Lys Trp Asp Leu Leu Ile Lys Gln Trp Val Cys Asp Pro Leu
 50 55 60
 Gly Gly Gly Gly Gly Val Asp Lys Thr His Thr Cys Pro Pro Cys
 65 70 75
 Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro
 80 85 90
 Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val
 95 100 105
 Thr Cys Trp Trp Asp Val Ser His Glu Asp Pro Glu Val Lys Phe
 110 115 120
 Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys
 125 130 135
 Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Trp Ser Val Leu

140	145	150
Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys		
155	160	165
Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile		
170	175	180
Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu		
185	190	195
Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr		
200	205	210
Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp		
215	220	225
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro		
230	235	240
Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr		
245	250	255
Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser		
260	265	270
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu		
275	280	285
Ser Leu Ser Pro Gly Lys		
290		

<210> 11

<211> 471

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 11

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr		
1	5	10
15		
Gly Val His Ser Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu		
20	25	30
Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly		
35	40	45
Tyr Thr Phe Thr Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro		
50	55	60
Gly Lys Gly Leu Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly		
65	70	75

Asp Thr Ser Tyr Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser	80	85	90
Val Asp Lys Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu	95	100	105
Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr	110	115	120
Tyr Ser Asn Ser Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr	125	130	135
Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe	140	145	150
Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala	155	160	165
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val	170	175	180
Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro	185	190	195
Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val	200	205	210
Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn	215	220	225
Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu	230	235	240
Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala	245	250	255
Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys	260	265	270
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys	275	280	285
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn	290	295	300
Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro	305	310	315
Arg Glu Glu Gln Tyr Asn Ala Thr Tyr Arg Val Val Ser Val Leu	320	325	330
Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys	335	340	345

Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Ala Ala Thr Ile
 350 355 360
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
 365 370 375
 Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr
 380 385 390
 Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 395 400 405
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
 410 415 420
 Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
 425 430 435
 Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser
 440 445 450
 Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
 455 460 465
 Ser Leu Ser Pro Gly Lys
 470

<210> 12

<211> 1377

<212> DNA

<213> Homo sapiens

<400> 12

agcatcctga gtaatgagtg gcctgggccc gagcaggcca ggtggccgga 50
 gccgtgtgga ccaggaggag cgctttccac agggcctgtg gacgggggtg 100
 gctatgagat cctgccccga agagcagtag tgggatcctc tgctgggtac 150
 ctgcatgtcc tgcaaaacca ttgcaacca tcagagccag cgcacctgtg 200
 cagccttctg caggtcactc agctgccgca aggagcaagg caagttctat 250
 gaccatctcc tgagggactg catcagctgt gcctccatct gtggacagca 300
 ccctaagcaa tgtgcatact tctgtgagaa caagctcagg agcccagtga 350
 accttccacc agagctcagg agacagcgga gtggagaagt tgaaaacaat 400
 tcagacaact cggaaggta ccaaggattg gagcacagag gctcagaagc 450
 aagtcagct ctcccggggc tgaagctgag tgcagatcag gtggccctgg 500
 tctacagcac gctggggctc tgctgtgtg ccgtcctctg ctgcttctg 550

gtggcggtgg cctgcttcct caagaagagg ggggatccct gtccttgcca 600
 gccccgctca aggccccgtc aaagtccggc caagtcttcc caggatcacg 650
 cgatggaagc cggcagccct gtgagcacat cccccgagcc agtggagacc 700
 tgcagcttct gcttcctga gtgcagggcg ccacgcagg agagcgagt 750
 cagcctggg acccccgacc ccacttggtg tggaagggtg ggtgcccaca 800
 ccaggaccac agtcctgcag ccttgcccac acatcccaga cagtggcctt 850
 ggcattgtgt gtgtgcctgc ccaggagggg ggcccagggtg cataaatggg 900
 ggtcagggag ggaaaggagg agggagagag atggagagga ggggagagag 950
 aaagagaggt ggggagaggg gagagagata tgaggagaga gagacagagg 1000
 aggcagaaag ggagagaaac agaggagaca gagagggaga gagagacaga 1050
 gggagagaga gacagagggg aagagaggca gagagggaaa gaggcagaga 1100
 aggaaagaga caggcagaga aggagagagg cagagaggga gagaggcaga 1150
 gagggagaga ggcagagaga cagagaggga gagagggaca gagagagata 1200
 gagcaggagg tcggggcact ctgagtccca gttcccagtg cagctgtagg 1250
 tcgtcatcac ctaaccacac gtgcaataaa gtctctgtgc ctgctgctca 1300
 cagccccga gagcccctcc tcctggagaa taaaaccttt ggcagctgcc 1350
 cttcctcaaa aaaaaaaaaa aaaaaaa 1377

<210> 13

<211> 107

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 13

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val
1				5					10					15

Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Ser	Ser	Val	Ser
				20					25					30

Tyr	Met	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Pro
				35					40					45

Leu	Ile	Tyr	Ala	Pro	Ser	Asn	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg
				50					55					60

Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75

Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp
80 85 90

Ser Phe Asn Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
95 100 105

Lys Arg

<210> 14

<211> 122

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 14

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr
20 25 30

Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45

Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr
50 55 60

Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser Val Asp Lys Ser
65 70 75

Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90

Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr Tyr Ser Asn Ser
95 100 105

Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val
110 115 120

Ser Ser

<210> 15

<211> 213

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 15

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val
1				5					10					15
Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Ser	Ser	Val	Ser
				20					25					30
Tyr	Met	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Pro
				35					40					45
Leu	Ile	Tyr	Ala	Pro	Ser	Asn	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg
				50					55					60
Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser
				65					70					75
Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Trp
				80					85					90
Ser	Phe	Asn	Pro	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile
				95					100					105
Lys	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser
				110					115					120
Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu
				125					130					135
Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp
				140					145					150
Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln
				155					160					165
Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu
				170					175					180
Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val
				185					190					195
Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg
				200					205					210

Gly Glu Cys

<210> 16

<211> 452

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 16

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	1	5	10	15
Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	20	25	30	
Ser	Tyr	Asn	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	35	40	45	
Glu	Trp	Val	Gly	Ala	Ile	Tyr	Pro	Gly	Asn	Gly	Asp	Thr	Ser	Tyr	50	55	60	
Asn	Gln	Lys	Phe	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Val	Asp	Lys	Ser	65	70	75	
Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	80	85	90	
Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Val	Val	Tyr	Tyr	Ser	Asn	Ser	95	100	105	
Tyr	Trp	Tyr	Phe	Asp	Val	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	110	115	120	
Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	125	130	135	
Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	140	145	150	
Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	155	160	165	
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	170	175	180	
Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	185	190	195	
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	200	205	210	
Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	215	220	225	
Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	230	235	240	
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	245	250	255	
Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	260	265	270	

Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
 275 280 285
 Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
 290 295 300
 Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
 305 310 315
 Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 320 325 330
 Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys
 335 340 345
 Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
 350 355 360
 Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
 365 370 375
 Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
 380 385 390
 Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 395 400 405
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 410 415 420
 Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
 425 430 435
 Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 440 445 450

Gly Lys

<210> 17

<211> 452

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 17

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr
 20 25 30

Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu

	35		40		45
Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr					
	50		55		60
Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser Val Asp Lys Ser					
	65		70		75
Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp					
	80		85		90
Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr Tyr Ser Asn Ser					
	95		100		105
Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val					
	110		115		120
Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro					
	125		130		135
Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu					
	140		145		150
Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser					
	155		160		165
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln					
	170		175		180
Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser					
	185		190		195
Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys					
	200		205		210
Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys					
	215		220		225
Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu					
	230		235		240
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr					
	245		250		255
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp					
	260		265		270
Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp					
	275		280		285
Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln					
	290		295		300
Tyr Asn Ala Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His					
	305		310		315

Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 320 325 330
 Lys Ala Leu Pro Ala Pro Ile Ala Ala Thr Ile Ser Lys Ala Lys
 335 340 345
 Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
 350 355 360
 Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
 365 370 375
 Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
 380 385 390
 Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 395 400 405
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 410 415 420
 Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
 425 430 435
 Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 440 445 450

Gly Lys

<210> 18

<211> 213

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 18

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15
 Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Ser Val Ser
 20 25 30
 Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro
 35 40 45
 Leu Ile Tyr Ala Pro Ser Asn Leu Ala Ser Gly Val Pro Ser Arg
 50 55 60
 Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
 65 70 75

Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp
 80 85 90
 Ala Phe Asn Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
 95 100 105
 Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser
 110 115 120
 Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu
 125 130 135
 Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp
 140 145 150
 Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln
 155 160 165
 Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu
 170 175 180
 Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val
 185 190 195
 Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg
 200 205 210

Gly Glu Cys

<210> 19

<211> 213

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 19

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15
 Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Ser Val Ser
 20 25 30
 Tyr Leu His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro
 35 40 45
 Leu Ile Tyr Ala Pro Ser Asn Leu Ala Ser Gly Val Pro Ser Arg
 50 55 60
 Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
 65 70 75
 Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp

	80		85		90
Ala Phe Asn Pro	Pro Thr Phe Gly Gln	Gly Thr Lys Val	Glu Ile		
	95		100		105
Lys Arg Thr Val	Ala Ala Pro Ser Val	Phe Ile Phe Pro	Pro Ser		
	110		115		120
Asp Glu Gln Leu	Lys Ser Gly Thr Ala	Ser Val Val Cys	Leu Leu		
	125		130		135
Asn Asn Phe Tyr	Pro Arg Glu Ala Lys	Val Gln Trp Lys	Val Asp		
	140		145		150
Asn Ala Leu Gln	Ser Gly Asn Ser Gln	Glu Ser Val Thr	Glu Gln		
	155		160		165
Asp Ser Lys Asp	Ser Thr Tyr Ser Leu	Ser Ser Thr Leu	Thr Leu		
	170		175		180
Ser Lys Ala Asp	Tyr Glu Lys His Lys	Val Tyr Ala Cys	Glu Val		
	185		190		195
Thr His Gln Gly	Leu Ser Ser Pro Val	Thr Lys Ser Phe	Asn Arg		
	200		205		210

Gly Glu Cys

<210> 20

<211> 452

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 20

Glu Val Gln Leu Val	Glu Ser Gly Gly Gly	Leu Val Gln Pro	Gly
1	5	10	15
Gly Ser Leu Arg Leu	Ser Cys Ala Ala Ser	Gly Tyr Thr Phe	Thr
20	25	30	
Ser Tyr Asn Met His	Trp Val Arg Gln Ala	Pro Gly Lys Gly	Leu
35	40	45	
Glu Trp Val Gly Ala	Ile Tyr Pro Gly Asn	Gly Asp Thr Ser	Tyr
50	55	60	
Asn Gln Lys Phe Lys	Gly Arg Phe Thr Ile	Ser Val Asp Lys	Ser
65	70	75	
Lys Asn Thr Leu Tyr	Leu Gln Met Asn Ser	Leu Arg Ala Glu	Asp
80	85	90	

Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Val	Val	Tyr	Tyr	Ser	Asn	Ser		95	100	105
Tyr	Trp	Tyr	Phe	Asp	Val	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val		110	115	120
Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro		125	130	135
Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu		140	145	150
Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser		155	160	165
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln		170	175	180
Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser		185	190	195
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys		200	205	210
Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys		215	220	225
Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu		230	235	240
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr		245	250	255
Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp		260	265	270
Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp		275	280	285
Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln		290	295	300
Tyr	Asn	Ala	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His		305	310	315
Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn		320	325	330
Lys	Ala	Leu	Pro	Ala	Pro	Ile	Ala	Ala	Thr	Ile	Ser	Lys	Ala	Lys		335	340	345
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg		350	355	360

Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
 365 370 375
 Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
 380 385 390
 Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 395 400 405
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 410 415 420
 Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
 425 430 435
 Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 440 445 450

Gly Lys

<210> 21
 <211> 213
 <212> PRT
 <213> Artificial sequence

<220>
 <223> sequence is synthesized

<400> 21
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15
 Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Ser Val Ser
 20 25 30
 Tyr Leu His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro
 35 40 45
 Leu Ile Tyr Ala Pro Ser Asn Leu Ala Ser Gly Val Pro Ser Arg
 50 55 60
 Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
 65 70 75
 Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp
 80 85 90
 Ala Phe Asn Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
 95 100 105
 Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser
 110 115 120
 Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu

	125		130		135
Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp					
	140		145		150
Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln					
	155		160		165
Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu					
	170		175		180
Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val					
	185		190		195
Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg					
	200		205		210

Gly Glu Cys

<210> 22

<211> 452

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 22

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly					
1	5		10		15
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr					
	20		25		30
Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu					
	35		40		45
Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr					
	50		55		60
Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser Val Asp Lys Ser					
	65		70		75
Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp					
	80		85		90
Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr Tyr Ser Asn Ser					
	95		100		105
Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val					
	110		115		120
Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro					
	125		130		135

Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu
				140					145					150
Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
				155					160					165
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln
				170					175					180
Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
				185					190					195
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys
				200					205					210
Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys
				215					220					225
Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu
				230					235					240
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr
				245					250					255
Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp
				260					265					270
Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp
				275					280					285
Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln
				290					295					300
Tyr	Asn	Ala	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His
				305					310					315
Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn
				320					325					330
Lys	Ala	Leu	Pro	Ala	Pro	Ile	Ala	Ala	Thr	Ile	Ser	Lys	Ala	Lys
				335					340					345
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg
				350					355					360
Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys
				365					370					375
Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly
				380					385					390
Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
				395					400					405

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 410 415 420

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
 425 430 435

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 440 445 450

Gly Lys

<210> 23
 <211> 121
 <212> PRT
 <213> Mus musculus

<400> 23
 Gln Ala Tyr Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Pro Gly
 1 5 10 15
 Ala Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr
 20 25 30
 Ser Tyr Asn Met His Trp Val Lys Gln Thr Pro Arg Gln Gly Leu
 35 40 45
 Glu Trp Ile Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr
 50 55 60
 Asn Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser
 65 70 75
 Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp
 80 85 90
 Ser Ala Val Tyr Phe Cys Ala Arg Val Val Tyr Tyr Ser Asn Ser
 95 100 105
 Tyr Trp Tyr Phe Asp Val Trp Gly Thr Gly Thr Thr Val Thr Val
 110 115 120

Ser

<210> 24
 <211> 106
 <212> PRT
 <213> Mus musculus

<400> 24
 Gln Ile Val Leu Ser Gln Ser Pro Ala Ile Leu Ser Ala Ser Pro
 1 5 10 15
 Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser Ser Ser Val Ser

	20		25		30
Tyr Met His Trp	Tyr Gln Gln Lys Pro	Gly Ser Ser Pro Lys Pro			
	35		40		45
Trp Ile Tyr Ala	Pro Ser Asn Leu Ala	Ser Gly Val Pro Ala Arg			
	50		55		60
Phe Ser Gly Ser	Gly Ser Gly Thr Ser	Tyr Ser Leu Thr Ile Ser			
	65		70		75
Arg Val Glu Ala	Glu Asp Ala Ala Thr	Tyr Tyr Cys Gln Gln Trp			
	80		85		90
Ser Phe Asn Pro	Pro Thr Phe Gly Ala	Gly Thr Lys Leu Glu Leu			
	95		100		105

Lys

<210> 25
 <211> 293
 <212> PRT
 <213> Homo sapiens

<400> 25
 Met Ser Gly Leu Gly Arg Ser Arg Arg Gly Gly Arg Ser Arg Val
 1 5 10 15
 Asp Gln Glu Glu Arg Phe Pro Gln Gly Leu Trp Thr Gly Val Ala
 20 25 30
 Met Arg Ser Cys Pro Glu Glu Gln Tyr Trp Asp Pro Leu Leu Gly
 35 40 45
 Thr Cys Met Ser Cys Lys Thr Ile Cys Asn His Gln Ser Gln Arg
 50 55 60
 Thr Cys Ala Ala Phe Cys Arg Ser Leu Ser Cys Arg Lys Glu Gln
 65 70 75
 Gly Lys Phe Tyr Asp His Leu Leu Arg Asp Cys Ile Ser Cys Ala
 80 85 90
 Ser Ile Cys Gly Gln His Pro Lys Gln Cys Ala Tyr Phe Cys Glu
 95 100 105
 Asn Lys Leu Arg Ser Pro Val Asn Leu Pro Pro Glu Leu Arg Arg
 110 115 120
 Gln Arg Ser Gly Glu Val Glu Asn Asn Ser Asp Asn Ser Gly Arg
 125 130 135
 Tyr Gln Gly Leu Glu His Arg Gly Ser Glu Ala Ser Pro Ala Leu
 140 145 150

Pro Gly Leu Lys Leu Ser Ala Asp Gln Val Ala Leu Val Tyr Ser
 155 160 165
 Thr Leu Gly Leu Cys Leu Cys Ala Val Leu Cys Cys Phe Leu Val
 170 175 180
 Ala Val Ala Cys Phe Leu Lys Lys Arg Gly Asp Pro Cys Ser Cys
 185 190 195
 Gln Pro Arg Ser Arg Pro Arg Gln Ser Pro Ala Lys Ser Ser Gln
 200 205 210
 Asp His Ala Met Glu Ala Gly Ser Pro Val Ser Thr Ser Pro Glu
 215 220 225
 Pro Val Glu Thr Cys Ser Phe Cys Phe Pro Glu Cys Arg Ala Pro
 230 235 240
 Thr Gln Glu Ser Ala Val Thr Pro Gly Thr Pro Asp Pro Thr Cys
 245 250 255
 Ala Gly Arg Trp Gly Cys His Thr Arg Thr Thr Val Leu Gln Pro
 260 265 270
 Cys Pro His Ile Pro Asp Ser Gly Leu Gly Ile Val Cys Val Pro
 275 280 285
 Ala Gln Glu Gly Gly Pro Gly Ala
 290

<210> 26

<211> 995

<212> DNA

<213> Homo sapiens

<400> 26

aagactcaaa cttagaaact tgaattagat gtggtattca aatccttacg 50
 tgccgcgaag acacagacag cccccgtaag aaccacgaa gcaggcgaag 100
 ttcattgttc tcaacattct agctgctctt gctgcatttg ctctggaatt 150
 cttgtagaga tattacttgt ccttccaggc tggtctttct gtagctccct 200
 tggtttcttt ttgtgatcat gttgcagatg gctgggcagt gctcccaaaa 250
 tgaatatttt gacagtttgt tgcattgctg cataccttgt caacttcgat 300
 gttctttctaa tactctctct ctaacatgac agcggtattg taatgcaagt 350
 gtgaccaatt cagtgaagg aacgaatgcg attctctgga cctggttggg 400
 actgagctta ataatttctt tggcagtttt cgtgctaattg tttttgctaa 450

ggaagataag ctctgaacca ttaaaggacg agtttaaaaa cacaggatca 500
 ggtctcctgg gcatggctaa cattgacctg gaaaagagca ggactgggtga 550
 tgaaattatt cttccgagag gcctcgagta cacggtggaa gaatgcacct 600
 gtgaagactg catcaagagc aaaccgaagg tcgactctga ccattgcttt 650
 ccactcccag ctatggagga aggcgcaacc attcttgtca ccacgaaaac 700
 gaatgactat tgcaagagcc tgccagctgc tttgagtgt acggagatag 750
 agaaatcaat ttctgctagg taattaacca tttcgactcg agcagtgcc 800
 ctttaaaaaat cttttgtcag aatagatgat gtgtcagatc tctttaggat 850
 gactgtatatt ttcatgtgcc gatacagctt tttgtcctct aactgtggaa 900
 actctttatg ttagatatat ttctctaggt tactgttggg agcttaatgg 950
 tagaaacttc cttggtttca tgattaaagt cttttttttt cctga 995

<210> 27

<211> 184

<212> PRT

<213> Homo sapiens

<400> 27

Met	Leu	Gln	Met	Ala	Gly	Gln	Cys	Ser	Gln	Asn	Glu	Tyr	Phe	Asp
1				5					10					15
Ser	Leu	Leu	His	Ala	Cys	Ile	Pro	Cys	Gln	Leu	Arg	Cys	Ser	Ser
				20					25					30
Asn	Thr	Pro	Pro	Leu	Thr	Cys	Gln	Arg	Tyr	Cys	Asn	Ala	Ser	Val
				35					40					45
Thr	Asn	Ser	Val	Lys	Gly	Thr	Asn	Ala	Ile	Leu	Trp	Thr	Cys	Leu
				50					55					60
Gly	Leu	Ser	Leu	Ile	Ile	Ser	Leu	Ala	Val	Phe	Val	Leu	Met	Phe
				65					70					75
Leu	Leu	Arg	Lys	Ile	Ser	Ser	Glu	Pro	Leu	Lys	Asp	Glu	Phe	Lys
				80					85					90
Asn	Thr	Gly	Ser	Gly	Leu	Leu	Gly	Met	Ala	Asn	Ile	Asp	Leu	Glu
				95					100					105
Lys	Ser	Arg	Thr	Gly	Asp	Glu	Ile	Ile	Leu	Pro	Arg	Gly	Leu	Glu
				110					115					120
Tyr	Thr	Val	Glu	Glu	Cys	Thr	Cys	Glu	Asp	Cys	Ile	Lys	Ser	Lys
				125					130					135

Pro Lys Val Asp Ser Asp His Cys Phe Pro Leu Pro Ala Met Glu
 140 145 150

Glu Gly Ala Thr Ile Leu Val Thr Thr Lys Thr Asn Asp Tyr Cys
 155 160 165

Lys Ser Leu Pro Ala Ala Leu Ser Ala Thr Glu Ile Glu Lys Ser
 170 175 180

Ile Ser Ala Arg

<210> 28
 <211> 858
 <212> DNA
 <213> Homo sapiens

<400> 28
 atggatgact ccacagaaag ggagcagtca cgccttactt cttgccttaa 50
 gaaaagagaa gaaatgaaac tgaaggagtg tgtttccatc ctcccacgga 100
 aggaaagccc ctctgtccga tcctccaaag acggaaagct gctggctgca 150
 accttgctgc tggcactgct gtcttgctgc ctcacggtgg tgtctttcta 200
 ccaggtggcc gccctgcaag gggacctggc cagcctccgg gcagagctgc 250
 agggccacca cgcgagagaag ctgccagcag gagcaggagc cccaaggcc 300
 ggcttgaggg aagctccagc tgtcaccgag ggactgaaaa tctttgaacc 350
 accagctcca ggagaaggca actccagtca gaacagcaga aataagcgtg 400
 ccgttcaggg tccagaagaa acagtcactc aagactgctt gcaactgatt 450
 gcagacagtg aaacaccaac tatacaaaaa ggatcttaca catttgttcc 500
 atggcttctc agctttaaaa ggggaagtgc cctagaagaa aaagagaata 550
 aaatattggg caaagaaaact ggttactttt ttatatatgg tcaggtttta 600
 tatactgata agacctacgc catgggacat ctaattcaga ggaagaagg 650
 ccatgtcttt ggggatgaat tgagtctggg gactttgttt cgatgtattc 700
 aaaatatgcc tgaacacta cccaataatt cctgctattc agctggcatt 750
 gcaaaaactgg aagaaggaga tgaactccaa cttgcaatac caagagaaaa 800
 tgcacaaata tactggatg gagatgtcac attttttggg gcattgaaac 850
 tgctgtga 858

<210> 29

<211> 285

<212> PRT

<213> Homo sapiens

<400> 29

Met	Asp	Asp	Ser	Thr	Glu	Arg	Glu	Gln	Ser	Arg	Leu	Thr	Ser	Cys	1	5	10	15
Leu	Lys	Lys	Arg	Glu	Glu	Met	Lys	Leu	Lys	Glu	Cys	Val	Ser	Ile	20	25	30	
Leu	Pro	Arg	Lys	Glu	Ser	Pro	Ser	Val	Arg	Ser	Ser	Lys	Asp	Gly	35	40	45	
Lys	Leu	Leu	Ala	Ala	Thr	Leu	Leu	Leu	Ala	Leu	Leu	Ser	Cys	Cys	50	55	60	
Leu	Thr	Val	Val	Ser	Phe	Tyr	Gln	Val	Ala	Ala	Leu	Gln	Gly	Asp	65	70	75	
Leu	Ala	Ser	Leu	Arg	Ala	Glu	Leu	Gln	Gly	His	His	Ala	Glu	Lys	80	85	90	
Leu	Pro	Ala	Gly	Ala	Gly	Ala	Pro	Lys	Ala	Gly	Leu	Glu	Glu	Ala	95	100	105	
Pro	Ala	Val	Thr	Ala	Gly	Leu	Lys	Ile	Phe	Glu	Pro	Pro	Ala	Pro	110	115	120	
Gly	Glu	Gly	Asn	Ser	Ser	Gln	Asn	Ser	Arg	Asn	Lys	Arg	Ala	Val	125	130	135	
Gln	Gly	Pro	Glu	Glu	Thr	Val	Thr	Gln	Asp	Cys	Leu	Gln	Leu	Ile	140	145	150	
Ala	Asp	Ser	Glu	Thr	Pro	Thr	Ile	Gln	Lys	Gly	Ser	Tyr	Thr	Phe	155	160	165	
Val	Pro	Trp	Leu	Leu	Ser	Phe	Lys	Arg	Gly	Ser	Ala	Leu	Glu	Glu	170	175	180	
Lys	Glu	Asn	Lys	Ile	Leu	Val	Lys	Glu	Thr	Gly	Tyr	Phe	Phe	Ile	185	190	195	
Tyr	Gly	Gln	Val	Leu	Tyr	Thr	Asp	Lys	Thr	Tyr	Ala	Met	Gly	His	200	205	210	
Leu	Ile	Gln	Arg	Lys	Lys	Val	His	Val	Phe	Gly	Asp	Glu	Leu	Ser	215	220	225	
Leu	Val	Thr	Leu	Phe	Arg	Cys	Ile	Gln	Asn	Met	Pro	Glu	Thr	Leu	230	235	240	
Pro	Asn	Asn	Ser	Cys	Tyr	Ser	Ala	Gly	Ile	Ala	Lys	Leu	Glu	Glu	245	250	255	

Gly	Asp	Glu	Leu	Gln	Leu	Ala	Ile	Pro	Arg	Glu	Asn	Ala	Gln	Ile
				260					265					270
Ser	Leu	Asp	Gly	Asp	Val	Thr	Phe	Phe	Gly	Ala	Leu	Lys	Leu	Leu
				275					280					285

<210> 30

<211> 1348

<212> DNA

<213> Homo sapiens

<400> 30

```

ggtagcaggc ttcctagagg gactggaacc taattctcct gaggctgagg 50
gaggggtggag ggtctcaagg caacgctggc cccacgacgg agtgccagga 100
gcactaacag tacccttagc ttgctttcct cctccctcct ttttattttc 150
aagttccttt ttattttctcc ttgcgtaaca accttcttcc cttctgcacc 200
actgcccgta cccttaccgg ccccgccacc tccttgctac cccactcttg 250
aaaccacagc tggtggcagg gtccccagct catgccagcc tcattctcctt 300
tcttgctagc ccccaaaggg cctccaggca acatgggggg cccagtcaga 350
gagccggcac tctcagttgc cctctggttg agttgggggg cagctctggg 400
ggccgtggct tgtgccatgg ctctgctgac ccaacaaaca gagctgcaga 450
gcctcaggag agaggtgagc cggctgcagg ggacaggagg cccctcccag 500
aatggggaag ggtatccctg gcagagtctc ccggagcaga gttccgatgc 550
cctggaagcc tgggagaatg gggagagatc ccggaaaagg agagcagtgc 600
tcacccaaaa acagaagaag cagcactctg tcctgcacct gggtcccatt 650
aacgccacct ccaaggatga ctccgatgtg acagaggtga tgtggcaacc 700
agctcttagg cgtgggagag gcctacaggc ccaaggatat ggtgtccgaa 750
tccaggatgc tggagtttat ctgctgtata gccaggtcct gtttcaagac 800
gtgactttca ccatgggtca ggtggtgtct cgagaaggcc aaggaaggca 850
ggagactcta ttccgatgta taagaagtat gccctccac ccggaccggg 900
cctacaacag ctgctatagc gcaggtgtct tcattttaca ccaaggggat 950
attctgagtg tcataattcc ccgggcaagg gcgaaactta acctctctcc 1000
acatggaacc ttcctggggg ttgtgaaact gtgattgtgt tataaaaagt 1050

```

ggctcccagc ttggaagacc aggggtgggta catactggag acagccaaga 1100
 gctgagtata taaaggagag ggaatgtgca ggaacagagg catcttcctg 1150
 ggtttggtc cccgttcctc acttttccct ttccattccc accccctaga 1200
 ctttgatttt acggatatct tgcttctgtt ccccatggag ctccgaattc 1250
 ttgctgtgtg ttagatgagg ggcggggggac gggcgccagg cattgttcag 1300
 acctggtcgg ggcccactgg aagcatccag aacagcacca ccatctta 1348

<210> 31

<211> 250

<212> PRT

<213> Homo sapiens

<400> 31

Met	Pro	Ala	Ser	Ser	Pro	Phe	Leu	Leu	Ala	Pro	Lys	Gly	Pro	Pro	1	5	10	15
Gly	Asn	Met	Gly	Gly	Pro	Val	Arg	Glu	Pro	Ala	Leu	Ser	Val	Ala	20	25	30	
Leu	Trp	Leu	Ser	Trp	Gly	Ala	Ala	Leu	Gly	Ala	Val	Ala	Cys	Ala	35	40	45	
Met	Ala	Leu	Leu	Thr	Gln	Gln	Thr	Glu	Leu	Gln	Ser	Leu	Arg	Arg	50	55	60	
Glu	Val	Ser	Arg	Leu	Gln	Gly	Thr	Gly	Gly	Pro	Ser	Gln	Asn	Gly	65	70	75	
Glu	Gly	Tyr	Pro	Trp	Gln	Ser	Leu	Pro	Glu	Gln	Ser	Ser	Asp	Ala	80	85	90	
Leu	Glu	Ala	Trp	Glu	Asn	Gly	Glu	Arg	Ser	Arg	Lys	Arg	Arg	Ala	95	100	105	
Val	Leu	Thr	Gln	Lys	Gln	Lys	Lys	Gln	His	Ser	Val	Leu	His	Leu	110	115	120	
Val	Pro	Ile	Asn	Ala	Thr	Ser	Lys	Asp	Asp	Ser	Asp	Val	Thr	Glu	125	130	135	
Val	Met	Trp	Gln	Pro	Ala	Leu	Arg	Arg	Gly	Arg	Gly	Leu	Gln	Ala	140	145	150	
Gln	Gly	Tyr	Gly	Val	Arg	Ile	Gln	Asp	Ala	Gly	Val	Tyr	Leu	Leu	155	160	165	
Tyr	Ser	Gln	Val	Leu	Phe	Gln	Asp	Val	Thr	Phe	Thr	Met	Gly	Gln	170	175	180	
Val	Val	Ser	Arg	Glu	Gly	Gln	Gly	Arg	Gln	Glu	Thr	Leu	Phe	Arg				

	185		190		195
Cys Ile Arg Ser Met Pro Ser His Pro Asp Arg Ala Tyr Asn Ser					
	200		205		210
Cys Tyr Ser Ala Gly Val Phe His Leu His Gln Gly Asp Ile Leu					
	215		220		225
Ser Val Ile Ile Pro Arg Ala Arg Ala Lys Leu Asn Leu Ser Pro					
	230		235		240
His Gly Thr Phe Leu Gly Phe Val Lys Leu					
	245		250		

<210> 32

<211> 595

<212> DNA

<213> Homo sapiens

<400> 32

cgtcggcacc atgaggcgag ggccccggag cctgcggggc agggacgcgc 50
 cagccccac gccctgcgtc ccggccgagt gcttcgacct gctggtccgc 100
 cactgcgtgg cctgcgggct cctgcgcacg ccgcggccga aaccggccgg 150
 ggccagcagc cctgcgcccc ggacggcgct gcagccgcag gagtgcgtgg 200
 gcgcggggggc cggcgaggcg gcgctgcccc tgcccgggct gctctttggc 250
 gccccgcgc tgctgggcct ggcactggtc ctggcgctgg tcttggtggg 300
 tctggtgagc tggaggcggc gacagcgcg gcttcgcggc gcgtcctccg 350
 cagaggcccc cgacggagac aaggacgccc cagagcccct ggacaaggtc 400
 atcattctgt ctccgggaat ctctgatgcc acagctcctg cctggcctcc 450
 tcttggggaa gaccaggaa ccacccacc tggccacagt gtccctgtgc 500
 cagccacaga gctgggctcc actgaactgg tgaccaccaa gacggccggc 550
 cctgagcaac aatagcaggg agccggcagg aggtggcccc tgccc 595

<210> 33

<211> 184

<212> PRT

<213> Homo sapiens

<400> 33

Met Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala					
1		5		10	15
Pro Thr Pro Cys Val Pro Ala Glu Cys Phe Asp Leu Leu Val Arg					
	20		25		30

His Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro
 35 40 45
 Ala Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln
 50 55 60
 Glu Ser Val Gly Ala Gly Ala Gly Glu Ala Ala Leu Pro Leu Pro
 65 70 75
 Gly Leu Leu Phe Gly Ala Pro Ala Leu Leu Gly Leu Ala Leu Val
 80 85 90
 Leu Ala Leu Val Leu Val Gly Leu Val Ser Trp Arg Arg Arg Gln
 95 100 105
 Arg Arg Leu Arg Gly Ala Ser Ser Ala Glu Ala Pro Asp Gly Asp
 110 115 120
 Lys Asp Ala Pro Glu Pro Leu Asp Lys Val Ile Ile Leu Ser Pro
 125 130 135
 Gly Ile Ser Asp Ala Thr Ala Pro Ala Trp Pro Pro Pro Gly Glu
 140 145 150
 Asp Pro Gly Thr Thr Pro Pro Gly His Ser Val Pro Val Pro Ala
 155 160 165
 Thr Glu Leu Gly Ser Thr Glu Leu Val Thr Thr Lys Thr Ala Gly
 170 175 180
 Pro Glu Gln Gln

<210> 34
 <211> 1881
 <212> DNA
 <213> Mus musculus

<400> 34
 atgggcgcca ggagactccg gttccgaagc cagaggagcc gggacagctc 50
 ggtgcccacc cagtgaatc agaccgagtg cttcgaccct ctggtgagaa 100
 actgcgtgtc ctgtgagctc ttccacacgc cggacactgg acatacaagc 150
 agcctggagc ctgggacagc tctgcagcct caggagggct ccgcgctgag 200
 acccgacgtg gcgctgctcg tcggtgcccc cgactcctg ggactgatac 250
 tggcgctgac cctggtgggt ctagtgagtc tggtagctg gaggtggcgt 300
 caacagctca ggacggcctc ccagacact tcagaaggag tccagcaaga 350
 gtccctggaa aatgtctttg taccctcctc agaaaccct catgcctcag 400

ctcctacctg gcctccgctc aaagaagatg cagacagcgc cctgccacgc 450
cacagcgtcc cgggtgcccgc cacagaactg ggctccaccg agctggtgac 500
caccaagaca gctggcccag agcaatagca gcagtggagg ctggaaccca 550
gggatctcta ctgggcttgt ggacttcacc caacagcttg ggaaagaact 600
tggcccttca gtgacggagt cctttgcctg gggggcgaac ccggcagaac 650
cagacactac aggccacatg agattgcttt tgtgttagct cttgacttga 700
gaacgttcca tttctgagat ggtttttaag cctgtgtgcc ttcagatggt 750
tggtatagact tgagggttgc atatttaatc tctgtagtga gtcggagact 800
ggaaacttaa tctcgttcta aaaatttttg attactgggc tggaggtatg 850
gctcagcagt tcggtttgtg tgctgttcta gccgaggact ccagttgttc 900
agcttcccgg aactcagatc tggcagctta agaccacctg tcaactccagc 950
ccctggaaca tccttgctc caaaggcacc agcactcatt tgctctagag 1000
cacacacaca cacacacaca cacacacaca cacacacaca catatgcatg 1050
catgcacact taaaaatgtc aaaattagcg gctggagaaa ttcattggtca 1100
acagcgctta ctgtgattcc agaggatgag agtttgattc ccagaatgca 1150
ctgcgggttg ctctattactg agcataactt ttgcttcagg ggacctgatg 1200
cctctggact tcatgggcat ctgtattcac gtgcacatcc tacacacaca 1250
cacacacaca cacacagaca tacacacaca cacactcttt taaaaatgat 1300
aaaatataag ataggcatgg tggtagacac ctttaatccc aacattgggg 1350
aagcaaaggc aggcaggtaa ctgagttgga ggccatcctg gtctacatag 1400
caagttccag gctaaccaga gctaaatggt gagaccaagt ctcaaaataa 1450
tactcccccc ccaaaaaaaaa aaaactttta aattttgatt tttttctttt 1500
attattatatt tttatattaa tttcatggtg tttagaagtg gtatacttag 1550
atggtgacta agaggaggta aagccatcag gactgagccc ctaacatata 1600
aggagaaagc agagacaatg aacacgcccc tctcctgctg tgtgccagct 1650
ctggaccacc agccagaggg caatcatcag atgtggggcc tagaaccttc 1700
agagccgaaa gctaaatcaa tctcatttct ttgtaaagct atttagcctt 1750

aggtgttttg ttacggtgat ataaaatgga ctaacacagg cactatgagt 1800
 aagaagcttt tctttgagct gggaaaggta ctgttaaacc aaaattaatc 1850
 tgaataaaaa aaggctaagg ggaagacact t 1881

<210> 35
 <211> 175
 <212> PRT
 <213> Mus musculus

<400> 35
 Met Gly Ala Arg Arg Leu Arg Val Arg Ser Gln Arg Ser Arg Asp
 1 5 10 15
 Ser Ser Val Pro Thr Gln Cys Asn Gln Thr Glu Cys Phe Asp Pro
 20 25 30
 Leu Val Arg Asn Cys Val Ser Cys Glu Leu Phe His Thr Pro Asp
 35 40 45
 Thr Gly His Thr Ser Ser Leu Glu Pro Gly Thr Ala Leu Gln Pro
 50 55 60
 Gln Glu Gly Ser Ala Leu Arg Pro Asp Val Ala Leu Leu Val Gly
 65 70 75
 Ala Pro Ala Leu Leu Gly Leu Ile Leu Ala Leu Thr Leu Val Gly
 80 85 90
 Leu Val Ser Leu Val Ser Trp Arg Trp Arg Gln Gln Leu Arg Thr
 95 100 105
 Ala Ser Pro Asp Thr Ser Glu Gly Val Gln Gln Glu Ser Leu Glu
 110 115 120
 Asn Val Phe Val Pro Ser Ser Glu Thr Pro His Ala Ser Ala Pro
 125 130 135
 Thr Trp Pro Pro Leu Lys Glu Asp Ala Asp Ser Ala Leu Pro Arg
 140 145 150
 His Ser Val Pro Val Pro Ala Thr Glu Leu Gly Ser Thr Glu Leu
 155 160 165
 Val Thr Thr Lys Thr Ala Gly Pro Glu Gln
 170 175

<210> 36
 <211> 265
 <212> PRT
 <213> Homo sapiens

<400> 36
 Met Ser Gly Leu Gly Arg Ser Arg Arg Gly Gly Arg Ser Arg Val

1	5	10	15
Asp Gln Glu Glu Arg Phe Pro Gln Gly Leu Trp Thr Gly Val Ala	20	25	30
Met Arg Ser Cys Pro Glu Glu Gln Tyr Trp Asp Pro Leu Leu Gly	35	40	45
Thr Cys Met Ser Cys Lys Thr Ile Cys Asn His Gln Ser Gln Arg	50	55	60
Thr Cys Ala Ala Phe Cys Arg Ser Leu Ser Cys Arg Lys Glu Gln	65	70	75
Gly Lys Phe Tyr Asp His Leu Leu Arg Asp Cys Ile Ser Cys Ala	80	85	90
Ser Ile Cys Gly Gln His Pro Lys Gln Cys Ala Tyr Phe Cys Glu	95	100	105
Asn Lys Leu Arg Ser Pro Val Asn Leu Pro Pro Glu Leu Arg Arg	110	115	120
Gln Arg Ser Gly Glu Val Glu Asn Asn Ser Asp Asn Ser Gly Arg	125	130	135
Tyr Gln Gly Leu Glu His Arg Gly Ser Glu Ala Ser Pro Ala Leu	140	145	150
Pro Gly Leu Lys Leu Ser Ala Asp Gln Val Ala Leu Val Tyr Ser	155	160	165
Thr Leu Gly Leu Cys Leu Cys Ala Val Leu Cys Cys Phe Leu Val	170	175	180
Ala Val Ala Cys Phe Leu Lys Lys Arg Gly Asp Pro Cys Ser Cys	185	190	195
Gln Pro Arg Ser Arg Pro Arg Gln Ser Pro Ala Lys Ser Ser Gln	200	205	210
Asp His Ala Met Glu Ala Gly Ser Pro Val Ser Thr Ser Pro Glu	215	220	225
Pro Val Glu Thr Cys Ser Phe Cys Phe Pro Glu Cys Arg Ala Pro	230	235	240
Thr Gln Glu Ser Ala Val Thr Pro Gly Thr Pro Asp Pro Thr Cys	245	250	255
Ala Gly Arg Thr Ala Pro Pro Arg Glu Gly	260	265	

<210> 37

<211> 107

<212> PRT

<213> Mus musculus

<400> 37

Gln	Ile	Val	Leu	Ser	Gln	Ser	Pro	Ala	Ile	Leu	Ser	Ala	Ser	Pro
1				5					10					15
Gly	Glu	Lys	Val	Thr	Met	Thr	Cys	Arg	Ala	Ser	Ser	Ser	Val	Ser
				20					25					30
Tyr	Met	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Ser	Ser	Pro	Lys	Pro
				35					40					45
Trp	Ile	Tyr	Ala	Pro	Ser	Asn	Leu	Ala	Ser	Gly	Val	Pro	Ala	Arg
				50					55					60
Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Ser	Tyr	Ser	Leu	Thr	Ile	Ser
				65					70					75
Arg	Val	Glu	Ala	Glu	Asp	Ala	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Trp
				80					85					90
Ser	Phe	Asn	Pro	Pro	Thr	Phe	Gly	Ala	Gly	Thr	Lys	Leu	Glu	Leu
				95					100					105

Lys Arg

<210> 38

<211> 108

<212> PRT

<213> Homo sapiens

<400> 38

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val
1				5					10					15
Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Ser	Ile	Ser
				20					25					30
Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys
				35					40					45
Leu	Leu	Ile	Tyr	Ala	Ala	Ser	Ser	Leu	Glu	Ser	Gly	Val	Pro	Ser
				50					55					60
Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile
				65					70					75
Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln
				80					85					90
Tyr	Asn	Ser	Leu	Pro	Trp	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu
				95					100					105

Ile Lys Arg

<210> 39
 <211> 10
 <212> PRT
 <213> Artificial sequence

<220>
 <223> sequence is synthesized

<400> 39
 Arg Ala Ser Ser Ser Val Ser Tyr Met His
 5 10

<210> 40
 <211> 7
 <212> PRT
 <213> Artificial sequence

<220>
 <223> sequence is synthesized

<400> 40
 Ala Pro Ser Asn Leu Ala Ser
 5

<210> 41
 <211> 9
 <212> PRT
 <213> Artificial sequence

<220>
 <223> sequence is synthesized

<400> 41
 Gln Gln Trp Ser Phe Asn Pro Pro Thr
 5

<210> 42
 <211> 119
 <212> PRT
 <213> Homo sapiens

<400> 42
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
 20 25 30

Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 35 40 45

Glu Trp Val Ala Val Ile Ser Gly Asp Gly Gly Ser Thr Tyr Tyr

	50		55		60
Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser					
	65		70		75
Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp					
	80		85		90
Thr Ala Val Tyr Tyr Cys Ala Arg Gly Arg Val Gly Tyr Ser Leu					
	95		100		105
Tyr Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser					
	110		115		

<210> 43

<211> 10

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 43

Gly Tyr Thr Phe Thr Ser Tyr Asn Met His
5 10

<210> 44

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 44

Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr Asn Gln Lys Phe
1 5 10 15

Lys Gly

<210> 45

<211> 13

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<400> 45

Val Val Tyr Tyr Ser Asn Ser Tyr Trp Tyr Phe Asp Val
5 10

<210> 46

<211> 452

<221> Xaa
<222> 17
<223> Unknown amino acid

<400> 78
Glu Cys Phe Asp Leu Leu Val Arg Thr Trp Val Pro Cys Asn Val
1 5 10 15

Leu Xaa

<210> 79
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 79
Glu Cys Phe Asp Leu Leu Val Arg Arg Trp Val Pro Cys Glu Leu
1 5 10 15

Leu Val

<210> 80
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 80
Glu Cys Phe Asp Leu Leu Val Arg Ser Trp Val Pro Cys Tyr Ser
1 5 10 15

Leu Lys

<210> 81
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 81
Glu Cys Phe Asp Leu Leu Val Arg Gln Trp Val Ser Cys Gln Val
1 5 10 15

Phe Ala

<210> 82
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 82
Glu Cys Phe Asp Leu Leu Val Arg Val Trp Val Pro Cys Ser Arg
1 5 10 15

Leu Tyr

<210> 83
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 83
Glu Cys Phe Asp Leu Leu Val Arg Gln Trp Val Pro Cys Gly Ala
1 5 10 15

Leu Gly

<210> 84
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 84
Glu Cys Phe Asp Leu Leu Val Arg Ala Trp Val Pro Cys Asn Glu
1 5 10 15

Leu Arg

<210> 85
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 85

Glu Cys Phe Asp Leu Leu Val Arg Glu Trp Val Pro Cys Arg Ile
1 5 10 15

Leu Gln

<210> 86
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 86
Glu Cys Phe Asp Leu Leu Val Arg Arg Trp Val Pro Cys Ser Trp
1 5 10 15

Leu Leu

<210> 87
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 87
Glu Cys Phe Asp Leu Leu Val Arg Arg Trp Val Pro Cys Ser Leu
1 5 10 15

Val Lys

<210> 88
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 88
Glu Cys Phe Asp Leu Leu Val Arg Gln Trp Val Pro Cys Arg Ala
1 5 10 15

Leu Met

<210> 89
<211> 17
<212> PRT
<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 89

Glu	Cys	Phe	Asp	Leu	Leu	Val	Arg	Ala	Trp	Val	Pro	Cys	Ser	Tyr
1				5					10					15

Leu Ser

<210> 90

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 90

Glu	Cys	Phe	Asp	Leu	Leu	Val	Arg	Asp	Trp	Val	Pro	Cys	Ser	Leu
1				5					10					15

Leu Phe

<210> 91

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 91

Glu	Cys	Phe	Asp	Leu	Leu	Val	Arg	Ser	Trp	Val	Pro	Cys	Thr	Leu
1				5					10					15

Leu Ser

<210> 92

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 92

Glu	Cys	Phe	Asp	Leu	Leu	Val	Arg	Lys	Trp	Val	Pro	Cys	Ser	Thr
1				5					10					15

Phe His

<210> 93
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 93
Glu Cys Phe Asp Leu Leu Val Arg Gly Trp Val Pro Cys Ser Val
1 5 10 15

Leu Gln

<210> 94
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 94
Glu Cys Phe Asp Leu Leu Val Arg Ala Trp Val Pro Cys Ser Val
1 5 10 15

Leu Lys

<210> 95
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 95
Glu Cys Phe Asp Leu Leu Val Arg Gln Trp Val Ser Cys Glu Leu
1 5 10 15

Leu Ser

<210> 96
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 96

Glu Cys Phe Asp Leu Leu Val Arg Gly Trp Val Asp Cys Ser Leu
1 5 10 15

Leu Leu

<210> 97
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 97
Glu Cys Phe Asp Ile Leu Val Asp Arg Trp Val Pro Cys Ala Ile
1 5 10 15

Leu His

<210> 98
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 98
Glu Cys Phe Asp Arg Leu Val Gly His Trp Val Pro Cys Ala Ala
1 5 10 15

Leu Ile

<210> 99
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 99
Glu Cys Phe Asp Pro Leu Val Ala Arg Trp Val Pro Cys His Leu
1 5 10 15

Ile Asn

<210> 100
<211> 17
<212> PRT
<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 100

Glu	Cys	Phe	Asp	Pro	Leu	Val	Arg	Val	Trp	Val	Asp	Cys	Ser	Ile
1				5					10					15

Leu Asp

<210> 101

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 101

Glu	Cys	Phe	Asp	Ser	Leu	Val	Asn	Ala	Trp	Val	Pro	Cys	Ser	Ala
1				5					10					15

Ile Arg

<210> 102

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 102

Glu	Cys	Phe	Asp	Leu	Leu	Val	Asn	Arg	Trp	Val	Asp	Cys	Arg	Leu
1				5					10					15

Leu Ile

<210> 103

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 103

Glu	Cys	Phe	Asp	Pro	Leu	Val	Arg	Ile	Trp	Val	Ala	Cys	Asp	Arg
1				5					10					15

Leu Ala

<210> 104
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 104
Glu Cys Phe Asp Pro Leu Val Gly Arg Trp Val Pro Cys Thr Leu
1 5 10 15

Leu His

<210> 105
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 105
Glu Cys Phe Asp Leu Leu Val Arg Ala Trp Val Pro Cys His Leu
1 5 10 15

Ile Asp

<210> 106
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 106
Glu Cys Phe Asp Pro Leu Val Gly His Trp Val Pro Cys Ser Val
1 5 10 15

Leu Thr

<210> 107
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 107

Glu Cys Phe Asp Pro Leu Val Asn Arg Trp Val Asp Cys Val Ala
1 5 10 15

Leu His

<210> 108

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 108

Glu Cys Phe Asp Arg Leu Val Asn Leu Trp Val Asp Cys Ala Leu
1 5 10 15

Leu Asn

<210> 109

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 109

Glu Cys Phe Asp Val Leu Val Ser Ala Trp Val Asp Cys Ala Arg
1 5 10 15

Leu Asn

<210> 110

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 110

Glu Cys Phe Asp Ser Leu Val Arg Leu Trp Val Pro Cys Asn Leu
1 5 10 15

Leu Arg

<210> 111

<211> 22

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 111

Glu	Cys	Phe	Asp	Pro	Leu	Val	Arg	His	Trp	Val	Pro	Cys	Asn	Leu
1				5					10				15	

Leu	Arg	Gly	Ala	Gly	Ser	Pro
				20		

<210> 112

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 112

Glu	Cys	Phe	Asp	Ile	Leu	Val	Asn	Ala	Trp	Val	Pro	Cys	Arg	Val
1				5					10				15	

Ile	Gly
-----	-----

<210> 113

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 113

Glu	Cys	Phe	Asp	Arg	Leu	Val	Asn	Arg	Trp	Val	Pro	Cys	Asn	Leu
1				5					10				15	

Ile	Val
-----	-----

<210> 114

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 114

Glu	Cys	Phe	Asp	Arg	Leu	Val	Arg	Ala	Trp	Val	Pro	Cys	Thr	Ala
1				5					10				15	

Leu	Thr
-----	-----

<210> 115
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 115
Glu Cys Phe Asp Leu Leu Val Arg Arg Trp Val Pro Cys His Leu
1 5 10 15

Ile Thr

<210> 116
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 116
Glu Cys Phe Asp Ile Leu Val Gly Arg Trp Val Pro Cys Gly Leu
1 5 10 15

Ile His

<210> 117
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 117
Glu Cys Phe Asp Pro Leu Val Arg Asp Trp Val Arg Cys Asp Ile
1 5 10 15

Leu Thr

<210> 118
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 118

Glu Cys Phe Asp Pro Leu Val Arg Val Trp Val Pro Cys Thr Val
1 5 10 15

Leu Arg

<210> 119

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 119

Glu Cys Phe Asp Ser Leu Val Arg Ala Trp Val Pro Cys Gly Val
1 5 10 15

Leu Ser

<210> 120

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 120

Glu Cys Phe Asp Val Leu Val His Arg Trp Val Pro Cys Gly Leu
1 5 10 15

Ile Arg

<210> 121

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 121

Glu Cys Phe Asp His Leu Val Arg Ile Trp Val Pro Cys Thr Ala
1 5 10 15

Leu Ala

<210> 122

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 122

Glu	Cys	Phe	Asp	Thr	Leu	Val	Asn	Ala	Trp	Val	Pro	Cys	Asn	Leu
1					5				10					15

Leu Asp

<210> 123

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 123

Glu	Cys	Phe	Asp	Arg	Leu	Val	Asn	Gly	Trp	Val	Pro	Cys	Ala	Val
1				5					10					15

Leu His

<210> 124

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 124

Glu	Cys	Phe	Asp	Arg	Leu	Val	Asn	Ala	Trp	Val	Asp	Cys	Arg	Leu
1				5					10					15

Leu Ala

<210> 125

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 125

Glu	Cys	Phe	Asp	Leu	Leu	Val	Asn	Asp	Trp	Val	Pro	Cys	Gly	Ala
1				5					10					15

Ile Thr

<210> 126
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 126
Glu Cys Phe Asp Ala Leu Val Arg Arg Trp Val Asp Cys Ser Leu
1 5 10 15

Leu Arg

<210> 127
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 127
Glu Cys Phe Asp Ala Leu Val His Arg Trp Val Asp Cys Ala Val
1 5 10 15

Leu Gly

<210> 128
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 128
Glu Cys Phe Asp Val Leu Val Asn Ala Trp Val Asp Cys Ala Val
1 5 10 15

Leu Arg

<210> 129
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 129

Glu Cys Phe Asp Gly Leu Val Asn Ala Trp Val Asp Cys Gly Leu
1 5 10 15

Leu Arg

<210> 130
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 130
Glu Cys Phe Asp Pro Leu Val Arg His Trp Val Pro Cys Arg Ala
1 5 10 15

Leu Asp

<210> 131
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 131
Glu Cys Phe Asp Asp Leu Val Arg His Trp Val Pro Cys Asp Leu
1 5 10 15

Leu Thr

<210> 132
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 132
Glu Cys Phe Asp Val Leu Val Arg Ala Trp Val Pro Cys Arg Ala
1 5 10 15

Leu Thr

<210> 133
<211> 17
<212> PRT
<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 133

Glu	Cys	Phe	Asp	Ile	Leu	Val	Asn	Arg	Trp	Val	Pro	Cys	Gly	Ala
1				5					10					15

Leu Thr

<210> 134

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 134

Glu	Cys	Phe	Asp	Asp	Leu	Val	Arg	Asn	Trp	Val	Pro	Cys	Ala	Leu
1				5					10					15

Leu Asn

<210> 135

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 135

Glu	Cys	Phe	Asp	Pro	Leu	Val	Asn	Ala	Trp	Val	Pro	Cys	Ala	Val
1				5					10					15

Leu His

<210> 136

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 136

Glu	Cys	Phe	Asp	Pro	Leu	Val	Leu	Arg	Trp	Val	Pro	Cys	Ser	Ala
1				5					10					15

Leu His

<210> 137
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 137
Glu Cys Phe Asp Ala Leu Val His Arg Trp Val Pro Cys Asp Leu
1 5 10 15

Leu Arg

<210> 138
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 138
Glu Cys Phe Asp Pro Leu Val Arg Asp Trp Val Pro Cys Asp Leu
1 5 10 15

Ile His

<210> 139
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 139
Glu Cys Phe Asp Leu Leu Val Asn Ser Trp Val Pro Cys Ser Val
1 5 10 15

Ile Ala

<210> 140
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 140

Glu Cys Phe Asp Thr Leu Val Arg Ala Trp Val Pro Cys Ser His
1 5 10 15

Leu Thr

<210> 141

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 141

Glu Cys Phe Asp Ser Leu Val Arg Ile Trp Val Pro Cys Gly Leu
1 5 10 15

Ile Asp

<210> 142

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 142

Glu Cys Phe Asp Ser Leu Val Asn Ala Trp Val Pro Cys His Val
1 5 10 15

Leu Thr

<210> 143

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<220>

<221> Unsure

<222> 1

<223> Unknown amino acid

<220>

<221> Unsure

<222> 3

<223> Unknown amino acid

<220>

<221> Unsure
<222> 5
<223> Unknown amino acid

<220>
<221> Unsure
<222> 7-12
<223> Unknown amino acid

<220>
<221> Unsure
<222> 14-17
<223> Unknown amino acid

<400> 143
Xaa Cys Xaa Asp Xaa Leu Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa
1 5 10 15

Xaa Xaa

<210> 144
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<220>
<221> Unsure
<222> 1
<223> Unknown amino acid

<220>
<221> Unsure
<222> 3
<223> Unknown amino acid

<220>
<221> Unsure
<222> 5
<223> Unknown amino acid

<220>
<221> Unsure
<222> 8-9
<223> Unknown amino acid

<220>
<221> Unsure
<222> 11-12
<223> Unknown amino acid

<220>

<221> Unsure
<222> 14-15
<223> Unknown amino acid

<220>
<221> Unsure
<222> 17
<223> Unknown amino acid

<400> 144
Xaa Cys Xaa Asp Xaa Leu Val Xaa Xaa Trp Xaa Xaa Cys Xaa Xaa
1 5 10 15

Leu Xaa

<210> 145
<211> 17
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<220>
<221> Unsure
<222> 5
<223> Unknown amino acid

<220>
<221> Unsure
<222> 8-9
<223> Unknown amino acid

<220>
<221> Unsure
<222> 12
<223> Unknown amino acid

<220>
<221> Unsure
<222> 14-17
<223> Unknown amino acid

<400> 145
Glu Cys Phe Asp Xaa Leu Val Xaa Xaa Trp Val Xaa Cys Xaa Xaa
1 5 10 15

Xaa Xaa

<210> 146
<211> 70
<212> PRT
<213> Homo sapiens

<400> 146

Arg	Arg	Gly	Pro	Arg	Ser	Leu	Arg	Gly	Arg	Asp	Ala	Pro	Ala	Pro
1				5					10					15

Thr	Pro	Cys	Val	Pro	Ala	Glu	Cys	Phe	Asp	Leu	Leu	Val	Arg	His
				20					25					30

Cys	Val	Ala	Cys	Gly	Leu	Leu	Arg	Thr	Pro	Arg	Pro	Lys	Pro	Ala
				35					40					45

Gly	Ala	Ser	Ser	Pro	Ala	Pro	Arg	Thr	Ala	Leu	Gln	Pro	Gln	Glu
				50					55					60

Ser	Val	Gly	Ala	Gly	Ala	Gly	Glu	Ala	Ala
				65				70	

<210> 147

<211> 63

<212> PRT

<213> Mus musculus

<400> 147

Arg	Arg	Leu	Arg	Val	Arg	Ser	Gln	Arg	Ser	Arg	Asp	Ser	Ser	Val
1				5					10					15

Pro	Thr	Gln	Cys	Asn	Gln	Thr	Glu	Cys	Phe	Asp	Pro	Leu	Val	Arg
				20					25					30

Asn	Cys	Val	Ser	Cys	Glu	Leu	Phe	His	Thr	Pro	Asp	Thr	Gly	His
				35					40					45

Thr	Ser	Ser	Leu	Glu	Pro	Gly	Thr	Ala	Leu	Gln	Pro	Gln	Glu	Gly
				50					55					60

Ser Ala Leu

<210> 148

<211> 71

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 148

Arg	Arg	Leu	Arg	Val	Arg	Ser	Gln	Arg	Ser	Arg	Asp	Ser	Pro	Ala
1				5					10					15

Pro	Thr	Pro	Cys	Val	Pro	Ala	Glu	Cys	Phe	Asp	Leu	Leu	Val	Arg
				20					25					30

His	Cys	Val	Ala	Cys	Gly	Leu	Leu	Arg	Thr	Pro	Arg	Pro	Lys	Pro
				35					40					45

Ala Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln
50 55 60

Glu Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
65 70

<210> 149

<211> 71

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 149

Arg Arg Leu Arg Val Arg Ser Gln Arg Ser Arg Asp Ser Ser Val
1 5 10 15

Pro Thr Gln Cys Val Pro Ala Glu Cys Phe Asp Leu Leu Val Arg
20 25 30

His Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro
35 40 45

Ala Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln
50 55 60

Glu Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
65 70

<210> 150

<211> 71

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 150

Arg Arg Leu Arg Val Arg Ser Gln Arg Ser Arg Asp Ser Ser Val
1 5 10 15

Pro Thr Gln Cys Asn Gln Thr Glu Cys Phe Asp Pro Leu Val Arg
20 25 30

His Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro
35 40 45

Ala Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln
50 55 60

Glu Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
65 70

<210> 151
 <211> 71
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Sequence is synthesized

<400> 151
 Arg Arg Leu Arg Val Arg Ser Gln Arg Ser Arg Asp Ser Ser Val
 1 5 10 15
 Pro Thr Gln Cys Asn Gln Thr Glu Cys Phe Asp Pro Leu Val Arg
 20 25 30
 Asn Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro
 35 40 45
 Ala Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln
 50 55 60
 Glu Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
 65 70

<210> 152
 <211> 70
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Sequence is synthesized

<400> 152
 Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
 1 5 10 15
 Thr Pro Cys Asn Gln Thr Glu Cys Phe Asp Pro Leu Val Arg His
 20 25 30
 Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
 35 40 45
 Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
 50 55 60
 Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
 65 70

<210> 153
 <211> 70
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Sequence is synthesized

<400> 153

```

Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
 1           5           10           15

Thr Pro Cys Asn Pro Thr Glu Cys Phe Asp Pro Leu Val Arg His
           20           25           30

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
           35           40           45

Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
           50           55           60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
           65           70

```

<210> 154

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 154

```

Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
 1           5           10           15

Thr Pro Cys Asn Gln Ala Glu Cys Phe Asp Pro Leu Val Arg His
           20           25           30

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
           35           40           45

Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
           50           55           60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
           65           70

```

<210> 155

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 155

```

Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
 1           5           10           15

Thr Pro Cys Asn Pro Ala Glu Cys Phe Asp Pro Leu Val Arg His
           20           25           30

```

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
35 40 45

Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
50 55 60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
65 70

<210> 156

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 156

Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
1 5 10 15

Thr Pro Cys Val Gln Thr Glu Cys Phe Asp Pro Leu Val Arg His
20 25 30

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
35 40 45

Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
50 55 60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
65 70

<210> 157

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 157

Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
1 5 10 15

Thr Pro Cys Val Gln Ala Glu Cys Phe Asp Pro Leu Val Arg His
20 25 30

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
35 40 45

Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
50 55 60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
 65 70

<210> 158

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 158

Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
 1 5 10 15

Thr Pro Cys Val Pro Ala Glu Cys Phe Asp Pro Leu Val Arg His
 20 25 30

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
 35 40 45

Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
 50 55 60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
 65 70

<210> 159

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 159

Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
 1 5 10 15

Thr Pro Cys Asn Pro Ala Glu Cys Phe Asp Ser Leu Val Arg His
 20 25 30

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
 35 40 45

Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
 50 55 60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
 65 70

<210> 160

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 160

Arg	Arg	Gly	Pro	Arg	Ser	Leu	Arg	Gly	Arg	Asp	Ala	Pro	Ala	Pro
1					5				10					15

Thr	Pro	Cys	Asn	Pro	Ala	Glu	Cys	Phe	Asp	Ala	Leu	Val	Arg	His
			20						25					30

Cys	Val	Ala	Cys	Gly	Leu	Leu	Arg	Thr	Pro	Arg	Pro	Lys	Pro	Ala
			35						40					45

Gly	Ala	Ser	Ser	Pro	Ala	Pro	Arg	Thr	Ala	Leu	Gln	Pro	Gln	Glu
				50					55					60

Ser	Val	Gly	Ala	Gly	Ala	Gly	Glu	Ala	Ala
			65						70

<210> 161

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 161

Arg	Arg	Gly	Pro	Arg	Ser	Leu	Arg	Gly	Arg	Asp	Ala	Pro	Ala	Pro
1					5				10					15

Thr	Pro	Cys	Asn	Gln	Thr	Glu	Cys	Phe	Asp	Leu	Leu	Val	Arg	His
			20						25					30

Cys	Val	Ala	Cys	Gly	Leu	Leu	Arg	Thr	Pro	Arg	Pro	Lys	Pro	Ala
			35						40					45

Gly	Ala	Ser	Ser	Pro	Ala	Pro	Arg	Thr	Ala	Leu	Gln	Pro	Gln	Glu
				50					55					60

Ser	Val	Gly	Ala	Gly	Ala	Gly	Glu	Ala	Ala
			65						70

<210> 162

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 162

Arg	Arg	Gly	Pro	Arg	Ser	Leu	Arg	Gly	Arg	Asp	Ala	Pro	Ala	Pro
1					5				10					15

```

Thr Pro Cys Asn Pro Thr Glu Cys Phe Asp Leu Leu Val Arg His
      20                      25                      30

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
      35                      40                      45

Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
      50                      55                      60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
      65                      70

```

<210> 163

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 163

```

Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
  1              5              10              15

Thr Pro Cys Asn Gln Ala Glu Cys Phe Asp Leu Leu Val Arg His
      20                      25                      30

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
      35                      40                      45

Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
      50                      55                      60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
      65                      70

```

<210> 164

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 164

```

Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
  1              5              10              15

Thr Pro Cys Val Gln Ala Glu Cys Phe Asp Leu Leu Val Arg His
      20                      25                      30

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
      35                      40                      45

```


Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
50 55 60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
65 70

<210> 165

<211> 70

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 165

Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
1 5 10 15

Thr Pro Cys Asn Pro Ala Glu Cys Phe Asp Leu Leu Val Arg His
20 25 30

Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala
35 40 45

Gly Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu
50 55 60

Ser Val Gly Ala Gly Ala Gly Glu Ala Ala
65 70

INTERNATIONAL SEARCH REPORT

 International application No
 PCT/US2005/044926

A. CLASSIFICATION OF SUBJECT MATTER

C07K14/705 C07K16/28 A61K47/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data, MEDLINE, EMBASE, Sequence Search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CRAXTON ANDREW ET AL: "A role for zTNF4/APRIL in the co-stimulation of B cell proliferation by macrophages" FASEB JOURNAL, vol. 16, no. 4, 20 March 2002 (2002-03-20), pages A704-A705, XP009064584 & ANNUAL MEETING OF THE PROFESSIONAL RESEARCH SCIENTISTS ON EXPERIMENTAL BIOLOGY; NEW ORLEANS, LOUISIANA, USA; APRIL 20-24, 2002 ISSN: 0892-6638 the whole document ----- -/--	1-58



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

5 Apr11 2006

Date of mailing of the international search report

03/05/2006

Name and mailing address of the ISA/

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Sommerfeld, T

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2005/044926

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>REFF M E ET AL: "Depletion of B cells in vivo by a chimeric mouse human monoclonal antibody to CD20" BLOOD, W.B. SAUNDERS, PHILADELPHIA, VA, US, vol. 83, no. 2, 15 January 1994 (1994-01-15), pages 435-445, XP002112549 ISSN: 0006-4971 the whole document</p>	
A	<p>GROSS J A ET AL: "TACI and BCMA are receptors for a TNF homologue implicated in B-cell autoimmune disease" NATURE, NATURE PUBLISHING GROUP, LONDON, GB, vol. 404, 27 April 2000 (2000-04-27), pages 995-999, XP002140939 ISSN: 0028-0836 the whole document</p>	
A	<p>WO 00/43032 A (BIOGEN, INC; APOTEC S.A; BROWNING, JEFFREY; AMBROSE, CHRISTINE; MACKA) 27 July 2000 (2000-07-27) the whole document</p>	
P,X	<p>WO 2005/000351 A (GENENTECH, INC; CHAN, ANDREW; GONG, QIAN; MARTIN, FLAVIUS) 6 January 2005 (2005-01-06) the whole document</p>	1-58
P,X	<p>ANOLIK ET AL: "New treatments for SLE: cell-depleting and anti-cytokine therapies" BAILLIERE'S BEST PRACTICE AND RESEARCH. CLINICAL RHEUMATOLOGY, BAILLIERE TINDALL, LONDON, GB, vol. 19, no. 5, October 2005 (2005-10), pages 859-878, XP005140891 ISSN: 1521-6942 the whole document</p>	1-58

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2005/044926

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: —
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-53 and 56-58 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the proscribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

 International application No
 PCT/US2005/044926

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0043032	A	27-07-2000	AT 247482 T	15-09-2003
			AU 762839 B2	03-07-2003
			AU 3214200 A	07-08-2000
			BR 0007719 A	13-11-2001
			CA 2360062 A1	27-07-2000
			CN 1347326 A	01-05-2002
			CZ 20012696 A3	17-10-2001
			DE 60004635 D1	25-09-2003
			DE 60004635 T2	09-06-2004
			DK 1146892 T3	24-11-2003
			EE 200100386 A	16-12-2002
			EP 1146892 A2	24-10-2001
			ES 2204528 T3	01-05-2004
			HK 1040628 A1	14-10-2005
			HU 0105283 A2	29-04-2002
			JP 2002535285 T	22-10-2002
			MX PA01007464 A	06-06-2003
			NO 20013641 A	25-09-2001
			NZ 513284 A	31-10-2003
			PL 349772 A1	09-09-2002
			PT 1146892 T	31-12-2003
			SK 10442001 A3	05-02-2002
			TR 200102147 T2	22-04-2002
			TR 200403498 T2	21-03-2005
WO 2005000351	A	06-01-2005	AU 2004251679 A1	06-01-2005
			AU 2004256042 A1	20-01-2005
			CA 2526402 A1	20-01-2005
			CA 2528434 A1	06-01-2005
			EP 1631313 A2	08-03-2006
			EP 1629001 A2	01-03-2006
			WO 2005005462 A2	20-01-2005